

ISSN 2278-4136

JPP 2014; 3 (2): 95-118

Received: 18-06-2014

Accepted: 23-06-2014

Mukul Chauhan

Department of Chemistry, College of
Natural and Computational Sciences,
Ambo University, Ethiopia

A review on Morphology, Phytochemistry and Pharmacological activities of medicinal herb *Plumbago Zeylanica* Linn.

Mukul Chauhan**Abstract**

Plumbagin is the most common and broad spectrum phytochemical of *P. zeylanica*. The leaves and root bark contains plumbagin. The root yield new pigments, viz, 3-chloroplumbagin, 3, 3- biplumbagin, binaphthoquinone identify as 3', 6'-biplumbagin, and four other pigments identify as isozeylanone, zeylanone, elliptinone, and droserone. The isolation of plumbagin, droserone, isoshinanone and a new naphthalenone i.e., 1, 2 (3)-tetrahydro-3, 3'- plumbagin is reported from the phenolic fraction of the light petrol extract of the roots. Two plumbagic acid glucosides; 3'-o-beta-glucopyranosyl plumbagic acid and 3'-o-beta- glucopyranosyl plumbagic acid methyl ester along with five naphthaquinones. All parts of the plant are used, but the roots have tremendous pharmacological properties. The pulped roots or aerial parts are reported abortifacient, while powdered bark, root or leaves are used to treat gonorrhoea, syphilis, tuberculosis, rheumatic pain, swellings, wound healing dyspepsia, piles, diarrhoea, skin diseases, leprosy and also reported to possess antibacterial, antifungal, and cantharides.

Keywords: Plumbagin, Anticancer activity, phytochemical, Antioxidant, Antibacterial drug, naphthoquinones, *Plumbago zeylanica*, Nonyl 8-methyl-dodec-7-enoate.

1. Introduction

From thousands of years and a remarkable number of modern drugs have been obtained from natural sources, particularly from the plants. Plant based medicines have played an important role in primary health care needs of human as well as animals. Variety of plants exhibit antimicrobial, larvicidal, anti- inflammatory and antioxidant activities due to the presence of some active compounds like essential oils, flavonoids, terpenoids, tri-terpenoids, glycosides, alkaloids and other natural phenolic compounds play a dominant role in the maintenance of human health since ancient times^[1]. Natural products play on important role in drug development programmes in the pharmaceutical industry^[2].

There are a few reports on the use of plants in traditional healing by either tribal people or indigenous community^[3-5]. Many reports show the effectiveness of traditional herbs against microorganisms; as a result, plants have become one of the bases of modern medicine^[6]. As an alternative form of health care and the development of microbial resistance to the available antibiotics has led researchers to investigate the antimicrobial activity of medicinal plants^[7-10]. Silver and Bostian^[11] have documented the use of natural products as new antibacterial drugs. There is an urgent need to identify novel substances active towards highly resistant pathogens^[12, 13]. It is thought that herbal remedies have the advantage of combining their active components with many other substances which appear to be inactive but which give the plant as a whole a level of safety and efficiency superior to that of its isolated, pure active components; moreover, in developing countries, synthetic drugs are presently too expensive and also are often adulterated^[14]. *P. zeylanica* Linn (Plumbaginaceae) is a perennial herb commonly distributed in forest of the Uttarakhand, India, and cultivated in the gardens throughout India. It grows wild as a garden plant in eastern, northern and southern India and *has* been reported to be used in variety of folk medicine in Africa and Asia.

It has been using in the treatment of refractory prostate cancer^[15] and shown anti fertility activity, antihyperlipidemic activity^[16], anti estrogenic activity^[17] to kill intestinal parasites, treat rheumatism, anemia due to "stagnant blood", external and internal trauma, toxic swelling and malignant furunculosis scabies^[18]. Antiplasmodial^[19], antimicrobial^[20], antifungal^[21], anti-inflammatory^[22], antibacterial^[23], hypolipidaemic and antiatherosclerotic activities^[24].

Correspondence:**Mukul Chauhan**

Department of Chemistry, College of
Natural and Computational
Sciences, Ambo University, Ethiopia

Plumbagin (5-hydroxy-2-methylnaphthalene-1,4-dione) is a naturally occurring yellow pigment produced by the members of *Plumbaginaceae*, accumulated mostly in root^[25]. Plumbagin showed anticancer^[26], antimicrobial and antibiotic^[27, 28], antibacterial and antifungal activities^[29]. Five coumarins – seselin^[30], 5-methoxyseselin^[31], suberosin^[32], xanthyletin and xanthoxyletin have been isolated from the roots of *Plumbago zeylanica*^[33].

The aim of this study was to emphasize the phytochemistry and pharmacological activities of *Plumbago zeylanica* herb.

2. Morphology

There is no consistency in the literature citing the classification of *P. zeylanica* as herb or shrub. Some authors have described it as a perennial dicot herb^[34, 35], while it has also been designated as a shrub by others^[36]. *P. zeylanica* plant attains a height of about 0.5–2 m (1.6–6.6 ft). The leaves are alternate, simple, ovate or ovate-lanceolate, elliptical or oblong, 0.5–12 cm in length with a tapered base 3 cm broad and often with a hairy margin. The stipules are absent and the petiole is narrow (0–5 mm long) with small auricles in young leaves. The inflorescence is of terminal raceme-type about 6–30 cm long and many-flowered. Flowers are white in colour^[34, 37] long, inodorous, inbracteate, axillary and terminal elongated spikes, bisexual regular, pentamerous, pedicellate and sweet-scented. Calyx densely covered with stalked, sticky glands. Corolla is white, very slender, and tubular and Stamens 5, free. Ovary superior, 5-gonous, one celled, ovule one basal.

The style is filiform with five elongated stigma lobes and the ovary is superior, single-celled. The flowers are also characterized by having a tubular calyx (7–11 mm long and 5-ribbed) with glandular trichomes (hair) secreting a sticky mucilage. The plant flowers round the year and pollination is primarily by insects. The mucilaginous glands aid in trapping insects and fruit dispersal by animals. The fruit of the plant is an oblong (7.5–8 mm long) five-furrowed capsule containing single seed. Each seed is oblong in structure, 5–6 mm long and reddish- brown to dark brown in colour. Roots are straight, smooth, branched or unbranched, with or without secondary roots and about 30 cm or more in length and 6 cm in diameter^[38]. They are light- yellow when fresh and become reddish-brown on drying. The roots have a strong and characteristic odour with acrid and bitter taste^[34].

3. Phytochemistry

Plumbagin is the most common and broad spectrum phytochemical of *P. zeylanica*. The leaves and root bark contains plumbagin. The root yield new pigment, viz, 3-chloroplumbagin, 3, 3- biplumbagin, binaphthoquinone identify as 3', 6'-biplumbagin, and four other pigments identify as isozeylanone, zeylanone, elliptinone, and droserone. The

isolation of plumbagin, droserone, isoshinanolone and a new naphthalenone i.e., 1, 2 (3)-tetrahydro-3, 3'- plumbagin is reported from the phenolic fraction of the light petrol extract of the roots. Two plumbagic acid glucosides; 3'-o-beta-glucopyranosyl plumbagic acid and 3'-o-beta- glucopyranosyl plumbagic acid methyl ester along with five naphthaquinones (plumbagin, chitranone, maritnone, elliptinone and isoshinanolone), and five coumarins (seselin, methoxyseselin, suberosine, xanthyletin and xanthoxyletin) were isolated from the roots isolated by Lin and coworkers^[33].


Some phytochemical from different parts of *P. zeylanica* are reported by different workers^[36, 39, 40, 41]. Like in stem plumbagin, zeylanone, isozeylanone, sitosterol, stigmasterol, campesterol, and dihydroflavonol plumbagin. In leaves plumbagin and chitanone. Flowers contain plumbagin, zeylanone, and glucose. Fruit contains plumbagin, glucopyranoside and sitosterol.










Seeds also contain plumbagin, and the root bark of *P. zeylanica* contains plumbagin. The root yield new pigment, viz, 3-chloroplumbagin, 3, 3- biplumbagin, binaphthoquinone identify as 3', 6'- biplumbagin, and four other pigments identify as isozeylanone, zeylanone, elliptinone, and droserone. The isolation of plumbagin, droserone, isoshinanolone and a new naphthalenone i.e., 1, 2 (3)-tetrahydro-3, 3'- plumbagin is reported from the phenolic fraction of the light petrol extract of the roots. Two plumbagic acid glucosides; 3'-o-beta-glucopyranosyl plumbagic acid and 3'-o-beta- glucopyranosyl plumbagic acid methyl ester along with five naphthaquinones (plumbagin, chitranone, maritnone, elliptinone and isoshinanolone), and five coumarins (seselin, methoxyseselin, suberosine, xanthyletin and xanthoxyletin) were isolated from the roots. Plumbagin (2-methyl-5-hydroxy-1,4-naphthoquinone) is a yellow crystalline bioactive phytoconstituent present in the roots isolated from *P. zeylanica* by soxhlet apparatus followed by silica gel column chromatography^[33, 42].








3.1 Phytochemicals isolated from *P. zeylanica*


Plumbagin is the major phytochemical isolated from *P. zeylanica* and it have tremendous medicinal values. Several other naphthoquinones, binaphthoquinones^[43-47], coumarins^[33, 48], di-phenyl sulfone^[49], carboxylic acids and esters^[50], meroterpenes^[51], triterpenoids^[52, 53], amino acids^[54], anthraquinones^[55], steroids^[56], steroid glucosides^[48], sugars^[57] and other compounds^[58-62]. Recently four other compounds one naphthoquinone and three difuranonaphthoquinones have been isolated and characterized^[63, 64] Table 1. Some qualitative phytochemicals results in leaves of *P. zeylanica* revealed the presence of alkaloids, glycoside, reducing sugars, simple phenolics, tannins, Lignin, saponins and flavonoids reported qualitatively^[41] and tested^[84].

Table 1: List of phytochemicals isolated from different parts of *p. zeylanica*


Compound names	References
Fatty acid esters:	
	
Linoleic acid	[67]


Palmitic acid [67]

Trilinolein [83]

Nonyl nonanoate [56]

Nonyl 8-methyl-dodec-7-enoate [56]

Hentriacontane [65]

1,2-Benzenedicarboxylic acid [50]

Diisooctyl phthalate [50]

Gugultetrol-18-ferrulate [65]

Vanillic acid [68]
Naphthoquinones:



Plumbagin [56,69,70]

3-chloroplumbagin [71]

3,8-dihydroxy-6-methoxy-2-isopropyl-1,4-naphthoquinone [44]

5,7-dihydroxy-8-methoxy-2-methyl-1,4-naphthoquinone [44]

2-methylnaphthazarin [43]

2-methyl-5-(3'-methyl-but-2'-enyloxy)-[1,4]naphthoquinone [63]

Droserone (3,5-Dihydroxy-2-methyl-1,4-naphthoquinone) [47,55]




Methylene-3,3'-diplumbagin [43]




Dihydrogesterone [72]




Isozeylanone [73]




Zeylanone [73]




Chitranone [45]




Maritinone [43]




Elliptinone [47]




3,3'-diplumbagin [74]



1,2(3)-tetrahydro-3,3'-biplumbagin [75]




9-hydroxy-2-isopropenyl-1,8-dioxa-dicyclopenta[b,g]naphthalene-4,10-dione [64]




2-(1-hydroxy-1-methyl-ethyl)-9-methoxy-1,8- [64]


dioxo-dicyclopenta [b,g]naphthalene-4,10-dione

Meroterpenes:


12- hydroxyisobakuchiol [51]




Bakuchiol [51]

Flavonoid and flavonoid glucosides:


2-(2,4-Dihydroxy-phenyl)-3,6,8-trihydroxy-chromen-4-one [76]




Saponaretin [51]




Isoorientin [51]




Isoaffinetin [51]

Coumarins:


Xanthyletin [51]




Xanthoxyletin [51]




Psoralen [51]




Seselin [33]




5-methoxyseselin [33]




Suberosin [33]




2,2- dimethyl- 5- hydroxy- 6- acetyl chromene [56]




2, 5- dimethyl- 7- hydroxychromone [77]

Amino acids and alkaloids:


Aspartic acid [54]




Tryptophan [54]




Tyrosine [54]




Threonine [54]




Alanine [54]




Histidine [54]




Glycine [54]




Methionine [54]




Hydroxyproline [54]




Indole- 3- carboxaldehyde [77]



Harman [78]











Neoechinulin A [78]



N - (N' - benzoyl- S - phenylalaninyl)- S - phenylalaninol [78]


Steroids:


<p>β-sitosterol [49,172]</p>

<p>Stigmasterol acetate [56]</p>

<p>Sitosterone [56]</p>

<p>Stigmasterol [56]</p>

<p>Ergostadiene-3β,5α,6β-triol [78]</p>

<p>α-amyrin [53]</p>

<p>β-amyrin [53]</p>




Taraxasterol

[53]




ψ -taraxasterol [79]


Terpenoids:




1-keto-3 β ,19 α -ainyaroxyurs-12-ene-24,28-dioic acid dimethyl ester [52]




3-O- β -D-arabinopyranosy derivative [52]




Friedelinol [56]



Lupeol [53]




Lupanone [56]




Lupeol acetate [83]


Glycosides:




β -sitosterol-3 β -glucopyranoside-6'-O-palmitate [60]






Hydroplumbagin glucoside [80]




β -sitosteryl glucoside [83]




3'-O- β -glucopyranosyl plumbagic acid [33]

1-hydroxy-3-methyl-6-methoxyanthraquinone-8-O- β -D-xylopyranoside [55]3'-O- β -glucopyranosyl plumbagic acid methylester [33]


Plumbagic acid [68,91]




Plumbagic acid Bu. Ester [51]




Isoshinznolone [78]








Isoshinanolone [46,51]



Benzyl 2,5-dihydroxy-6-methoxybenzoate [56]



1-acetoxy-4-hydroxy-2-methyl-5-methoxynaphthalene [74]


4-hydroxybenzaldehyde [77]

Androsta-1,4-diene-3,17-dione [78]

Trans-cinnamic acid [77]

Glucose [82]

Fructose [82]

3.2 Macro, micro and some essential elements detected in *P. zeylanica*:

Some elemental analysis has been done for leaves, stems and roots of *P. zeylanica* exists with abundant amounts of elements like four macro-elements (Na, K, Ca and Mg), five essential microelements (Zn, Fe, Mn, Cr and Co), and eight other elements (Mo, Sb, Bi, Cd, Sr, Pb, Cd and As) respectively were detected by inductively Coupled plasma atomic emission spectrometry (ICP-AES) [85].

4. Pharmacological activities of *P. zeylanica*.

The whole plant and its root have been used as a folk medicine in Taiwan for the treatment of rheumatic pain, menostasis, carbuncle and injury by bumping [86]. All parts of the plant are used, but the roots have tremendous pharmacological properties. The pulped roots or aerial parts are reported abortifacient, while powdered bark, root or leaves are used to treat gonorrhea, syphilis, tuberculosis, rheumatic pain, swellings, wound healing [87] dyspepsia, piles, diarrhoea, skin diseases, leprosy and also reported to possess antibacterial, antifungal [88], cantharides [83].

The pharmacological importance of this perennial shrub lies in its ability to produce a naphthoquinone, called plumbagin [89]. The main constituent in the root and

leaves is Plumbagin (2-methyl-5-hydroxy-1,4-naphthoquinone). Plumbagin is a yellow crystalline bioactive phytoconstituent [42] about 0.03% of dry weight of the roots. Plumbagin showing a broad range of pharmaceutical activities. Table 2 & 4

Pharmacological effects of plumbagin have been investigated on shortness of breath [90]. In Ayurvedic and Unani system of medicines, the plant has been described for significant anticancer [83, 91], antitumor [92], anti-inflammatory [22], antioxidant [93, 94], anti-mycobacterial [95] and antimicrobial activities [25, 39, 96, 97], rheumatic pain, sprains, scabies, skin diseases, and wounds. The roots of the plant and its constituents are credited with potential therapeutic properties including anti-atherogenic, cardioprotective, hepatoprotective, neuroprotective, and central nervous system stimulating properties [98], activity against canine distemper virus [99].

4.1 Pharmacological activities of different solvent extracts of *P. zeylanica*.

Acetone extract of *P. zeylanica* also effects on chromosomal aberrations induced by ethinylestradiol in cultured human lymphocytes [100]. hypolipidemic effect [101], anti-tumor [102-104], antimicrobial, anticancer, wound healing [105], anti-inflammatory and altered T-cell

proliferative activities ^[106, 107], and anti-fertility actions ^[108-112]. Plumbagin has also shown antibacterial activity against both gram-positive and gram-negative bacteria ^[128, 113-117], antihyperglycemic ^[118], insecticidal ^[119, 120], anti-allergic ^[121, 122] and antigonorrhoeal activity ^[123]. Besides, it has been also found active against certain yeasts fungi ^[124, 125], protozoa ^[19] and in tumor inhibitory activity ^[125]. It has also demonstrated significant

hyperglycemia, hypolipidemic, and antiatherosclerotic effects in rats ^[24, 65, 126-129]. The root of *P. zeylanica* has been reported to be a powerful poison when given orally or applied to ostiumuteri, causes abortion ^[130] cytotoxic and anti-insecticidal property ^[48, 131, 132]. Different kinds of solvent extracts of *P. zeylanica* have shown their broad spectrum pharmacological activities. Table-3

Table 2: Pharmacological activities of plumbagin (2-methyl-5-hydroxy-1,4-naphtho-quinone)


 <p>Plumbagin</p>	Antibacterial activity Antitumour activity Antyfertility activity Synergistic activity Anticoagulant activity Hypolipidaemic activity GST activity Antibacterial activity Blood coagulation activity AntiHelicobacter pylori activity	[28] [103,104] [104,111] [95] [127] [24] [93] [97] [129] [113]
--	--	---

Table 3: Pharmacological activities of some solvent extracts of *Plumbago Zeylanica* herb.

Type of plant extract	Dose ranges	Negative Control	Animal model / Microorganisms	Duration of the study	Results	Ref.
Methanolic extract of leaves	25, 50 & 100 mg	-	Indian earthworm (<i>Pheretima posthuma</i>)	2 days	Anthelmintic activity	133
Methanolic extract of root	50,100, &150 µg/ml	-	<i>Helicobacter pylori</i>	1 day	Anti- bacterial activity	134
Ethanol, acetone or ethyl acetate of Rhizome	30 µl	-	<i>Helicobacter pylori</i>	3 day	Anti- bacterial activity	113
Ethanolic extract of root	0.64-10.24 mg/ml	Ethanol	<i>E. coli</i> and <i>Shigella</i>	2 days	Anti- bacterial activity	135
Plumbago zeylanica root	100 mg/kg	-	Human study	14 days	Anti- hyper Cholesterolemic activity	34
Ethanolic extract of stems	500, 1000 mg/kg	48/80	Wistar Mice	2 days	Antiallergic activity	121
Methanol, chloroform and alcoholic extracts of leaves	50, 100 mg/ml	-	<i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i> , <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i>	1 day	Antibacterial activity	41
Methanol, Chloroform and	1 mg/ml	-	<i>E. coli</i> , <i>Salmonella typhi</i> , <i>Klebsiella</i>	2 days	Antibacterial activity	23

aqueous extract of root			<i>pneumoniae, Serratia marcescens, Proteus vulgaris, Pseudomonas aeruginosa, Staphylococcus aureus, Bacillus cereus</i>			
Petrol. ether, ethanol and aqueous extract of root	1200 µg/ml	-	<i>Staphylococcus aureus</i> and <i>Micrococcus luteus</i>	2 days	Antibacterial activity	39 172
Methanolic extract of leaves	50, 100 mg/ml	-	<i>Baci. subtilis, Staphylococcus aureus, Escherichia coli</i> and <i>Salmonella typhi</i>	2 days	Antibacterial activity	133 172
Ethanol extract of root	100, 200 mg/kg	Cancer cell line	Male Swiss albino mice	14 days	Anticancer activity	136
Ethanol extract of root	250 mg/kg bw	3-methyl-4- di methyl amino azo-benzene	Wistar albino rats	7 days	Anticarcinogenic activity	137
Hydroalcoholic extract of leaves	250, 500 mg/kg	Pentylene tetrazole	Wistar albino rats	1 hour	Anti-convulsion activity	138
Methanolic extract of root	4–10 mg/ml	Occluded dermal irritation	Albino rabbits, Swiss mice and Albino rats	1 day	Anti-dermatotoxicity	90
Ethanol extract of root	250 mg/kg bw	Alloxan	Wistar albino rats	21 day	Antidiabetic activity	118
Aqueous extract of root	100, 200 mg/kg	STZ	Wistar albino rats	42 day	Antidiabetic activity	139
Aqueous extract of leaves	100, 200 mg/kg	STZ	Wistar albino rats	28 day	Antidiabetic activity	140
Aqueous extract of leaves	100, 200 mg/kg	STZ	Wistar albino rats	28 day	Antidiabetic activity	141
Methanol, chloroform extract of whole plant	50 µl	-	<i>Rhizoctonia solani Kuhn, Bipolaris spp., Ustilago maydis</i> and <i>Alternaria alternata</i>	2 day	Antifungal activity	142 172
Ethanol extract of root	250, 500 mg /kg bw	Cyclophosphamide	Swiss albino mice	7 day	Antigenotoxic activity	94
Petroleum ether extract of root	300 mg/kg	Paracetamol	Wistar albino rats	7 days	Antihepatotoxic activity	143
Methanolic extract of aerial parts	35, 70 mg/kg	CCl ₄	Wistar albino rats	14 days	Antihepatotoxic activity	36
Aqueous extract of root	20,40, and 80 mg/kg	Diet-induced hyperlipidemic rats	Wistar albino rats	7 days	Antihyperlipidemic effect	17
Methanol, extract of root	300, 500 mg /kg	Carrageenin	Wistar albino rats	7 days	Anti-inflammatory activity	66
Petroleum ether, chloroform and acetone extract of root	0.1 ml	-	<i>Salmonella typhi, Staphylococcus aureus, Escherichia coli,</i>	2 days	Antimicrobial activity	144

			<i>Aspergillus niger</i> , <i>Pencillium</i> sp. And <i>Fusarium</i> <i>oxysporum</i>			
Methanolic extract of root	7.5 mg/ml	RPMI & Dimethyl Sulfoxide	male Sprague–Dawley rats	3 hours	Antimutagenic effect	145
Root powder	7.5 mg/kg bw	Phenyl-hydrazine	Male Wistar Albino rats	21 hours	Antioxidant activity	146
Ethanolic extract of root	1 mg/L	DPPH(1,1-diphenyl-2-picrylhydrazyl)ABTS (2,2-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt	<i>In-vitro</i> study	1 hour	Antioxidant activity	76
Ethanolic extract of root	100 mg/kg	-	<i>In-vitro</i>	2 hours	Antioxidant activity	147
Methanolic extract of leaves	50,100 mg/ml	Butylated hydroxy anisole	<i>In-vitro</i>	2 hours	Antioxidant activity	133
Methanolic extract of root	0.8–200 µg/ml	Guanidine hydrochloride, amantadine, and phosphonoformic acid	coxsackievirus B3 Nancy (CVB3), influenza A virus Hong Kong/1/68 (H3N2), and herpes simplex virus type 1 Kupka (HSV-1)	2 days	Antiviral activity	148
Aqueous extract of plant	0.5 µg/ml	-	Hepatitis B-virus	2 days	Antiviral activity	149
Ethanolic extract of root	250 mg/kg	Cholesterol	Rabbit	28 days	Hypolipidaemic activity	150
Ethanolic extract of root	250 mg/kg	Diet-induced hyperlipidemic rats	Rabbit	28 days	Hypolipidemic activity	151
Ethanolic extract of root	250 mg/kg bw	-	BALB/C mice	6 weeks	Immunomodulatory activity	152
Aqueous root extract	4 mg/ml	Turkey egg albumin	Balb/c mice	56 days	Immunosuppressive activity	153
Chloroform extract of root	100, 200 & 400 mg/kg	Scopolamine	Swiss albino mice	10 days	Memory Enhancing effect	154

Table 4: Plumbagin with putative anticancer and anti-proliferative tested in either *in vivo* or *in vitro* models

Cancer Cells	Results	Ref
Human Prostate cancer cell (PC-3, LNCaP, and C4-2)	Decrease in cell viability, apoptosis induction, Generation of ROS, depletion of intra cellular GSH	155

Human Melanoma A375.S2	Reduced amounts of cyclin B1, cyclin A, Cdc2, and Cdc25C and enhanced the levels of inactivated phosphorylated Cdc2 and Cdc25C, increased the activation of apoptosis signal-regulating kinase 1, JNK and extracellular signal-regulated kinase 1/2 (ERK1/2) and finally blocking ERK and JNK	156
Human non small cell lung cancer cells, A549	Activation of JNK and SP600125 (Aanthra [1,9-cd]pyrazol-6(2H)-one-1,9- pyrazoloanthrone), a specific inhibitor of JNK, decreased apoptosis by inhibiting the phosphorylation of p53 and subsequent increased in the interaction of p53 and MDM2. SP6000125 also inhibited the phosphorylation of Bcl-2 (Ser70	157
Human Peripheral blood lymphocytes	Effective cell growth inhibition, induces apoptosis, generates single-strand of DNA breaks and cytotoxic action	158
Human Prostate Cancer	Inhibition of both cultured Prostate Cancer cells and DU145 xenografts (a) the expression of protein kinase Cepsilon (PKCepsilon), phosphatidylinositol 3- kinase, phosphorylated AKT, phosphorylated Janus-activated kinase-2, and phosphorylated signal transducer and activator of transcription 3 (Stat3); (b) the DNA-binding activity of transcription factors activator protein-1, nuclear factor kappaB, and Stat3; and (c) Bcl-xL, cdc25A, and cyclooxygenase-2 expression	159
Human acute promyelocytic leukemia cells	Inhibition of proliferation of NB4 cells, chromosomes condensation and apoptotic body formation, cell proliferation and induce apoptosis of APL cell line NB4 cells	160
MCF7 and Bowes cancer cell lines	Inhibition of the proliferation of MCF7 and Bowes cells	83
Human hepatoma	Inhibition of the certain glycolytic enzymes and gluconeogenesis.	137
Human peripheral blood mononuclear cells	Involve the regulation of cell cycle progression, interleukin-2 and interferon production	161
MDA-MB-231 cells	Inhibitory effect on the protein levels of p- PI3K, p-Akt, p-JNK, p-ERK1/2, MMP-2, MMP-9, VEGF and HIF- 1 α	162
Human breast cancer cells	Inactivation of NF-kappaB and Bcl-2	163
Human breast cancer cells	Inhibit Akt activity and enhanced the activation of Chk2, resulting in increased inactive phosphorylation of Cdc25C and Cdc2.	164
Lung A549 cells	Increased the expression of p53 and phosphorylated p53 (Ser15 and Ser392) and regulates the levels of cell cycle related molecules in A549 and activates JNK	157
Human ovarian cancer cells	Bound to the active site of ER- α and inhibit classical ER- α signaling pathways	165
Cervical cancer cells	Lower dose of radiation in combination with plumbagin could induce apoptosis more effectively and activation of caspase 3 in C33A cells. Induction of apoptosis by irradiation and involves caspase-dependent pathways.	166
Human promyelocytic leukemia cells	Induced apoptotic cell death and inhibits tumor growth without obvious toxicity and triggering the mitochondria-dependent apoptosis of tumor cells by increasing ROS	91
Ovarian cancer cells	Induced loss of mitochondrial membrane potential, nuclear condensation, DNA fragmentation, and morphological changes	167
Human cervical cancer	Induced cell death is through the generation of ROS and subsequent induction of apoptosis caused loss of mitochondrial membrane potential and morphological changes characteristic of apoptosis, such as the translocation of phosphatidyl serine, nuclear condensation, and DNA fragmentation.	168
sarcoma-180	Ehrlich ascites model was evaluated and identified as less toxic, justified with the help of LD50 survival studies and stud	169

Azoxymethane induced intestinal carcinogenesis	Promising chemopreventive agents for human intestinal neoplasia	170
3T3-L1 cells	Activated PI3-kinase and/or PDK1 stimulate Akt activity with Ras–Raf–MEK1/2–ERK1/2 pathway	171

5. Conclusion

The study showed that the ethanol and petroleum ether and other solvent extract from the leaves, roots and stems of *Plumbago zeylanica* have anti microbial, antiviral, antioxidant, antifungal, anti-allergic and other wonderful medicinal properties. It is the most important medicinal plant extensively used in herbal formulations for centuries. The evidence presented in this review has shown that *Plumbago zeylanica* L. has tremendous potential to be integrated into conventional medical practice for the treatment and management of various metabolic syndromes, hepatotoxic, diabetes, inflammation, cancer and other disease complications. Some phenolic compounds have also been known as antioxidant agents, which act as free radical terminators and have shown medicinal activity as well as exhibiting physiological functions. It was reported that compounds such as flavonoids, which contain hydroxyls, are responsible for the radical scavenging effects of most plants. Development and research on *Plumbago zeylanica* through modern pharmaceutical technologies and analytical protocols is essential to assure its quality, safety and efficacy. It is anticipated that this review will provide some valuable information for ongoing explorations of this fascinating species its phytochemicals and pharmacological dynamics.

6. References

- Farombi EO. African indigenous plants with chemotherapeutic potentials and biotechnological approach to the production of bioactive prophylactic agents. *African J Biotech* 2003; 2:662-671.
- Baker JT *et al.* Natural products drug discovery and development. New perspective on international collaborations. *Nat prod* 1995; 58:1325-1357.
- Sandhy B, Thomas S, Isabel W and Shenbagavathai R. Ethnomedicinal plants used by the valaiyan community of Piranmalai hills (Reserved forest), Tamil Nadu, India. A pilot study. *African Journal of Traditional Complements and Alternative Medicines* 2006; 3:101-114.
- Ayyanar M, Iganacimuthu S. Traditional knowledge of kani tribals in Kouthalai of Tirunelveli hills. Tamil Nadu, India. *J of Enthopharmacology* 2005; 102:246-255.
- Rajan S, Sethuraman M, Mukherjee PK. Ethnobiology of the Nilgries Hills, India. *Phytotherapy Research* 2002; 16:98-116.
- Evans CE, Bansa A, Samuel OA. Efficacy of some nuppe medicinal plants against *Salmonella typhi*: an *in vitro* study. *J of Ethnopharmacology* 2002; 80:21-24.
- Bisignano G, Germano MP, Nostro A, Sanogo R. Drugs used in Africa as dyes: antimicrobial activities. *Phytotherapy Research* 1996; 9:346-350.
- Lis-balchin M, Deans SG. Antimicrobial effects of hydrophilic extracts of *Pelargonium* species (*Geraniaceae*). *Letters in Applied Microbiology* 1996; 23:205-207.
- Maoz M, Neeman I. Antimicrobial effects of aqueous plant extracts on the fungi *Microsporum canis* and *Trichophyton rubrum* and on three bacterial species. *Letters in Applied Microbiology* 1998; 26:61-63.
- Hammer KA, Carson CF, Riley TV. Antimicrobial activity of essential oils and other plant extracts. *J of Applied Microbiology* 1999; 86:985-990.
- Silver LL, Bostian KA. Discovery and development of new antibiotics: the problem of antibiotic resistance. *Antimicrobial Agents and Chemotherapy* 1993; 37:377-383.
- Recio MC. A review of some antimicrobial compounds isolated from medicinal plants reported in the literature 1978-1988. *Phytotherapy Research* 1989; 3:117-125.
- Cragg GM, Newman DJ, Snader KM. Natural products in drug discovery and development. *J of Nat Prod* 1997; 60:52-60.
- Shariff ZU. Modern herbal therapy for common ailments. *Nature Pharmacy Series, Spectrum Books Limited, Ibadan, Nigeria in Association with Safari Books (Export) Limited, United Kingdom*, 2001, 1:9-84.
- Moammir HA, Nancy E, Dreckschmidt, Ajit K. Plumbagin a medicinal plant –derived naphthaquinone, is a novel inhibitor of the Growth and Invasion of Hormone-Refractory Prostate cancer. *Cancer Res* 2008; 68(21):9024-9032.
- Dutta S, Vankatesh D, Souza R, Shenoy BD, Udupi RH and Udupa N. Niosomal Delivery of plumbagin ester for better antifertility activity. *Indian Drugs* 2002; 39(3):163-165.
- Pendurkar, Sudha R, Mengi, Sushma A. Antihyperlipidemic effect of aqueous extract of *Plumbago zeylanica* roots in diet induced hyperlipidemic rat. *Pharmaceutical Biology* 2009; 47(10):1004-1010.
- Jiangsu. New Medical College, "Zhongyao Dictionary (Encyclopedia of Chinese Materia Medica)". Scientific & Technological Press, Shanghai, 1979; 711-712.
- Simonsen HT *et al.* *In vitro* screening of Indian medicinal plants for antiparasitic activity. *Journal of Ethnopharmacology* 2001; 74:195-204.
- Ahmad I, Mehmood Z, Mohammad F, Ahmad S. Antimicrobial potency and synergistic activity of five traditionally used Indian medicinal plants. *Journal of Medicinal and Aromatic Plant Sciences* 2000; 23:173-176.
- Mehmood Z, Ahmad I, Mohammad F, Ahmad S. Indian medicinal plants: A potential source of anticandidal drugs. *Pharmaceutical Biology* 1999; 37:237-242.
- Oyedapo OO. Studies on the bioactivity of the extract of *Plumbago zeylanica*. *Phytotherapy Research* 1996; 13:346-348.
- Jeyachandran R, Mahesh A, Cindrella L, Sudhakar S,

- Pazhanichamy K. Antibacterial activity of plumbagin and root extracts of *plumbago zeylanica* L. *acta biologica cracoviensia Series Botanica* 2009; 51(1):17–22.
24. Sharma I, Gusain D, Dixit VP. Hypolipidaemic and antiatherosclerotic effects of plumbagin in rabbits. *Indian Journal of Physiology and Pharmacology* 1991; 35:10–14.
 25. Van-der VLM. Distribution of plumbagin in the *Plumbaginaceae*. *Phytochemistry* 1974; 11:3247–3248.
 26. Melo AM *et al.* First observations on the topical use of primin, plumbagin and mayteni in patients with skin cancer. *Revista do do Instituto de Antibioticos* 1974; 14:9–16.
 27. Brice HE. Antibacterial substances produced by flowering plants. *Australian Journal of Experimental Biology and Medicine Science* 1955; 33:547–554.
 28. Durga R, Sridhar P, Polasa H. Effect of plumbagin on antibiotic resistance in bacteria. *Indian Journal of Medical Research* 1990; 91:18–20.
 29. Gujar GT. Plumbagin, a naturally occurring naphthoquinone. Its pharmacological and pesticidal activity. *Fitoterapia* 1990; 59:387–393.
 30. Kostova I, Manolov I, Nicolova I, Danchev ND. New metal complexes of 4-methyl-7-hydroxycoumarin sodium salt and their pharmacological activity. *Il Farmaco* 2001; 56:707–713.
 31. Kofinas C, Chinou I, Loukis A, Harvala C, Roussakis C, Maillard M *et al.* Cytotoxic Coumarins from the aerial parts of *Tordylium apulum* and their effects on a non-small-cell bronchial carcinoma cell line. *Planta Medica* 1998; 64:174–176.
 32. Uchiyama T, Hara S, Makino M, Fujimoto Y. Seco-Adianane-type triterpenoids from *Dorstenia brasiliensi* (Moraceae). *Phytochemistry* 2002; 60:761–764.
 33. Lin LC, Yang LL, Chou CJ. Cytotoxic naphthoquinones and plumbagic acid glucosides from *Plumbago zeylanica*. *Phytochemistry* 2003; 62:619–622.
 34. Chetty KM *et al.* Pharmaceutical studies and therapeutic uses of *Plumbago Zeylanica* L. root *Ethnobotanical Leaflets* 2006; 10:294–304.
 35. Ming Y *et al.* Chemical constituents of *Plumbago zeylanica*. *Advanced Materials Research* 2011; 308–310:1662–1664.
 36. Rajesh K, Sushil K, Arjun P, Jayalakshmi S. hepatoprotective activity of aerial parts of *plumbago zeylanica* linn against carbon tetrachloride-induced hepatotoxicity in rats. *International J of pharmacy and pharmaceutical sciences* 2009; 1:suppl- 1.
 37. Lubaina AS *et al.* Shoot multiplication and direct organogenesis of an important medicinal plant *Plumbago zeylanica* L. (Plumbaginaceae). *Journal of Research in Biology* 2011; 6:424–428.
 38. Vishnukanta *et al.* Evaluation of anticonvulsant activity of *Plumbago zeylanica* Linn. leaf extract. *Asian Journal of Pharmaceutical and Clinical Research* 2010; 3(1):76–78.
 39. Ravikumar VR, Sudha T. Phytochemical and antimicrobial studies on *Plumbago zeylanica*. *IJRPC*, 2011; 1(2):185–188.
 40. Kakad S, Wabale AS, Kharde MN. Phytochemical Screening and Antimicrobial Studies on *Plumbago zeylanica* L. *Adv Biores* 2013; 4(3):115–117.
 41. Dhale DA, Markandeya SK. Antimicrobial and Phytochemical Screening of *Plumbago zeylanica* leaf. *J of Experimental Sciences* 2011; 2(3):04–06.
 42. Navneet K, Mishra BB, Tiwari VK, Tripathi V. Difuranonaphthoquinones from *Plumbago zeylanica* roots. *Phytochem Lett* 2010; 3:62–65.
 43. Gunaherath GM, Gunatilaka AA. Studies on medicinal and related plants of Sri Lanka. Part-18. Structure of a new naphthoquinone from *Plumbago zeylanica*. *J Chem Soc Perkin Trans* 1988; 1:407–410.
 44. Gupta A, Gupta A, Singh J. New naphthoquinones from *Plumbago zeylanica*. *Pharm Biol* 1999; 37:321–323.
 45. Dinda B, Saha S. A new binaphthoquinone from *Plumbago zeylanica* Linn. *Indian J Chem* 1989; 28:984–986.
 46. Gunaherath GM, Gunatilaka AA, Sultanbawa MU, Salasubramaniam S. 1,2(3)-Tetrahydro-3,3'-biplumbagin: A naphthalenone and other constituents from *Plumbago zeylanica*. *Phytochemistry* 1983; 22:1245–1247.
 47. Sankaram AV, Srinivasarao A, Sidhu GS. Chitranone- a new binaphthaquinone from *Plumbago zeylanica*. *Phytochemistry* 1976; 15:237–238.
 48. Yuan-Chuen W, Tung-Liang H. High-performance liquid chromatography for quantification of plumbagin, an antihelicobacter pylori compound of *plumbago zeylanica* L. *J Chromatogr* 2005; A 1094:99–104.
 49. Amatya S, Ghimire U, Tuladhar SM. Isolation of aromatic sulphone from *Plumbago zeylanica* Linn. *Pak J Sci Ind Res* 2007; 50:184–185.
 50. Rahman MS, Anwar MN. Fungitoxic and cytotoxic activity of a novel compound 1,2 benzenedicarboxylic acid, diisooctyl ester of *Plumbago zeylanica* Linn. *Asian J Microbiol Biotechnol Environ Sci* 2006; 8:461–464.
 51. Lin LC, Chou CJ. Meroterpenes and C-glucosylflavonoids from the aerial parts of *Plumbago zeylanica*. *Chinese Pharm J* 2003; 55:77–81.
 52. Gupta A, Rai R, Siddiqui IR, Singh J. Two new triterpenoids from *Plumbago zeylanica*. *Fitoterapia* 1998; 69:420–422.
 53. Dinda B, Saha S. Chemical constituents of *Plumbago zeylanica* aerial parts and *Thevetia neriifolia* roots. *J Indian Chem Soc* 1990; 67:88–89.
 54. Dinda B, Saha S. Free amino acids of *Plumbago zeylanica*. *J Indian Chem Soc* 1987; 64:261.
 55. Gupta A, Siddiqui IR, Singh J. A new anthraquinone glycoside from the roots of *Plumbago zeylanica*. *Indian J Chem Section B: Organic Chemistry Including Medicinal Chemistry* 2000; 39B:796–798.
 56. Gupta MM, Verma RK, Gupta AP. A chemical investigation of *Plumbago zeylanica*. *Curr Res Med Aromat Plants* 1995; 17:161–164.
 57. Chowdhury AK, Chakder SK, Khan AK. Isolation and characterization of chemical constituents of *Plumbago zeylanica* root. *J Bangladesh Acad Sci* 1981; 5:71–74.
 58. Dhar SK, Rao PG. Hormonal profile of plumbagin. *Fitoterapia* 1995; 66:442–446.
 59. Hsieh YJ, Lin LC, Tsai TH. Determination and identification of plumbagin from the roots of *Plumbago zeylanica* L. by liquid chromatography with tandem mass spectrometry. *J Chromatogr* 2005; A1083:141–145.
 60. Nguyen AT *et al.* Cytotoxicity of five plants used as anticancer remedies in Vietnamese traditional medicine. *Recent Prog Med Plants* 2006; 15:137–147.
 61. Chao L, Yuan L, Fang D, Qingyan M, Zewen G. Extraction, isolation and identification of plumbagin in *P. zeylanica* L. *Shizhen Guoyi Guoyao* 2006; 17:919.

62. Ariyanathan S, Saraswathy A, Rajamanickam GV. Quality control standards for the roots of three *Plumbago* species. *Indian J Pharm Sci* 2010; 72:86-91.
63. Kishore N, Mishra BB, Tiwari VK, Tripathi V. A novel naphthaquinone from *Plumbago zeylanica* L. roots. *Chem Nat Compd* 2010; 46:517-519.
64. Kishore N, Mishra BB, Tiwari VK, Tripathi V. Difuranonaphthoquinone from *Plumbago zeylanica* L. Roots *Phytochem Lett* 2010; 3:62-65.
65. Kumar R, Kumar S, Patra A, Jayalakshmi S. Hepatoprotective activity of aerial parts of *Plumbago zeylanica* Linn. Against carbon tetrachloride- induced hepatotoxicity in rats. *Int J Pharmacy Pharmaceut Sci* 2009; 1:171-175.
66. Arunachalam KD, Velmurugan P, Raja RB. Anti-inflammatory and cytotoxic effects of extract from *Plumbago zeylanica*. *Afri J Microbiol Res* 2010; 4:1239-1245.
67. Do DR, Nguyen XD. Chemical constituents of *Plumbago zeylanica* Linn. *Tap Chi Hoa Hoc* 1996; 34:67-70.
68. Qian XL, Zhou PZ. Study on the chemical constituents of *Plumbago zeylanica* Linn. I. Isolation and identification of the constituents of *Plumbago zeylanica* Linn. *Huaxue Xuebao* 1980; 38:405-408.
69. Padhye SB, Kulkarni BA. Root constituents of *Plumbago zeylanica*. *J Univ Poona Sci Technol* 1973; 44:27-29.
70. Israni SA, Kapadia NS, Lahiri SK, Yadav GK, Shah MB. An UV-visible spectrophotometric method for the estimation of plumbagin. *Int J Chem Tech Res* 2010; 2:856-859.
71. Sankaram AV, Siddhu GS. New biplumbagin and 3-chloroplumbagin from *P. zeylanica*. *Tetrahedron Lett* 1971; 26:2385-2388.
72. Dinda B, Saha S. A dihydronaphthaquinone from *Plumbago zeylanica*. *Chem Ind* 1986; 23:823.
73. Sankaram AV, Rao AS. Zeylanone and Isozeylanone, two novel quinones from *P. zeylanica*. *Tetrahedron* 1979; 35:1777-1782.
74. Sankaram AV, Sidhu GS. Synthesis of 3,3' Bi Plumbagin. *Indian J Chem* 1974; 12:519-520.
75. Gunaherath KG, Gunatilaka AA, Thomson RH. Structure of plumbazeylanone: a novel trimer of plumbagin from *P. zeylanica*. *Tetrahedron Lett* 1984; 25:4801-4804.
76. Nile SH, Khobragade CN. Antioxidant activity and flavonoid derivatives of *P. zeylanica*. *J Nat Prod* 2010; 3:130-133.
77. Zhang QR, Mei ZN, Yang GZ, Xiao YX. Chemical constituents from aerial parts of *P. zeylanica* L. *Zhong Yao Cai* 2007; 30:558-560.
78. Huang XY, Tan MX, Wu Q, Chen Y, Wang HS. Chemical constituents from the aerial parts of *P. zeylanica* L. *J Chinese Pharmaceut Sci* 2008; 17:144-147.
79. Rai MK, Pandey AK, Acharya D. Ethno-medicinal Plants Used by Gond Tribe of Bhanadehi, District Chhindwara, Madhya Pradesh. *J non-timber Forest Products* 2000; 7:237-241.
80. Yuan L, Zewen G, Yuxin H, Yan Q. *P. zeylanica* L. preparation for treating cancer and rheumatoid arthritis. China Patent: Faming Zhuanli Shenqing Gongkai Shuomingshu 2005, 3.
81. Qian XL, Liang XT, Cong PZ. Study on the chemical constituents of *P. zeylanica* L. II. Structural determination of plumbagic acid. *Huaxue Xuebao* 1980; 38:377-380.
82. Iyengar MA, Pendse GS. Pharmacognosy of the root of *P. zeylanica*. *Indian J Pharm* 1962; 24:290-291.
83. Nguyen AT, Malonne H, Duez P, Vanhaelen FR, Vanhaelen M, Fontaine J. Cytotoxic constituents from *Plumbago zeylanica*. *Int J Pharm* 2004; 75:500-504.
84. Tan MX, Chen ZF, Wang HS, Liu YC, Liang H. Analysis of macroelements and microelements in Chinese traditional medicine *Plumbago zeylanica* Linn by ICPAES. *Guang Pu Xue Yu Guang Pu Fen Xi* 2009; 29:1112-1114.
85. Harbone JB. "Phytochemical Method, A Guide to modern techniques of plant Analysis", Ed 3, springer (Indian) pvt. ltd., New Delhi, 2005, 22:5-16,
86. Okoli C, Akah O, Ezugworie U. Anti-inflammatory activity of extracts of root bark of *securidaca longipedunculata* fres (*polygalaceae*). *Afri J Trad Cam* 2005; 3:54-63.
87. Thakur RS, Puri HS, Husain A. Major medicinal plants of India. Lucknow, India: Central Institute of Medicinal and Aromatic Plants, 1989.
88. Uma DP, Soloman FE, Sharda AC. Indian medicinal plants and their roots. *Pharmaceut Biol* 1999; 37:231-236.
89. Ayo RG, Amupitip JO, Yimin Z. Cytotoxicity and antimicrobial studies of 1, 6, 8-trihydroxy-3-methyl-anyhraquinone (emodin) isolated from the leaves of *cassia nigricans*. *Vahl*, 2007; 6:1276-1279.
90. Teshome K, Gebre-Mariam T, Asres K, Perry F, Engidawork E. Toxicity studies on dermal application of plant extract of *P. zeylanica* used in Ethiopian traditional medicine. *J Ethnopharmacology* 2008; 117:236-248.
91. Xu KH, Lu DP. Plumbagin induces ROS-mediated apoptosis in human promyelocytic leukemia cells *in vivo*. *Leuk Res* 2010; 34:658-665.
92. Yang SJ, Chang SC, Wen HC, Chen CY, Liao JF, Chang CH. Plumbagin activates ERK1/2 and Akt via superoxide, Src and PI3-kinase in 3T3-L1 Cells. *Eur J Pharmacol* 2010; 638:21-28.
93. Sivakumar V, Prakash R, Murali MR, Devaraj H, Devaraj SN. *In vivo* micronucleus assay and GST activity in assessing genotoxicity of plumbagin in Swiss albino mice. *Drug Chem Toxicol* 2005; 28:499-507.
94. Sivakumar V, Devaraj SN. Protective Effect of *Plumbago zeylanica* Against Cyclophosphamide-Induced Genotoxicity and Oxidative Stress in Swiss Albino Mice. *Drug Chem Toxicol* 2006; 29:279-288.
95. Mossa JS, El-Ferally FS, Muhammad I. Antimycobacterial constituents from *Juniperus procera*, *Ferula communis* and *P. zeylanica* and their *in vitro* synergistic activity with isonicotinic acid hydrazide. *Phytother Res* 2004; 18:934-937.
96. Ahmad I, Mehmood Z, Mohammad F. Screening of some Indian medicinal plants for their antimicrobial properties. *J. Ethnopharmacol* 1998; 62:183-193.
97. Van-der VLM, Lotter AP. The constituents in the roots of *P. auriculata* Lam. and *P. zeylanica* L. responsible for antibacterial activity. *Planta Med* 1971; 20:8-13.
98. Bopaiah CP, Pradhan N. Central nervous system stimulatory action from the root extract of *Plumbago zeylanica* in rats. *Phytother Res* 2001; 15:153-156.
99. Bagla VP, McGaw LJ, Eloff JN. The antiviral activity of six South African plants traditionally used against infections in ethnoveterinary medicine. *Vet Microbiol* 2011.

100. Siddique YH, Ara G, Faisal M, Afzal M. Protective role of *P. zeylanica* extract against the toxic effects of ethinylestradiol in the third instar larvae of transgenic *Drosophila melanogaster* (hsp70-lacZ)Bg9 and cultured human peripheral blood lymphocytes. *Alternative Medicine Studies* 2011; 1:e726-9.
101. Ram A. Effect of *Plumbago zeylanica* in hyperlipidaemic rabbits and its modification by vitamin E. *Indian J Pharmacol* 1996; 28:161-166.
102. Gupta MM, Verma RK, Uniyal GC, Jain SP. Determination of plumbagin by normal phase high performance liquid chromatography. *J Chromat* 1993; 637:209-212.
103. Kavimani S, Ilango R, Madheswaran M, Jayakar B, Gupta M, Majumdar UK. Antitumor activity of plumbagin against Dalton's ascitic lymphoma. *Indian J Pharm Sci* 1996; 58:194-196.
104. Kini DP *et al.* Antitumor and antifertility activities of plumbagin controlled release formulations. *Indian J Exp Biol* 1997; 35:374-379.
105. Reddy JS, Rao PR, Reddy MS. Wound healing effects of *Heliotropium indicum*, *Plumbago zeylanicum* and *Acalypha indica* in rats. *J Ethnopharmacol* 2002; 79:249-251.
106. Aparanji P, Kumar BV, Kumar SP, Sreedevi K, Rao DN, Rao RA. Induction of anti-inflammatory and altered T-cell proliferative responses by the ethanolic extract of *P. zeylanica* in adjuvant-induced arthritic rats. *Pharm Biol* 2005; 43:784-789.
107. Aparanji P, Kumar BV, Rao TR, Rao DN, Rao RA. Alleviation of Collagen Induced Arthritis by *P. zeylanica* in Mice. *Pharm Biol* 2007; 45:54-59.
108. Edwin S, Joshi SB, Jain DC. Antifertility activity of leaves of *P. zeylanica* L. in female albino rats. *Eur J Contracept Reprod Health Care* 2009; 14:233-239.
109. Edwin S, Joshi SB, Jain DC. Bioassay-guided isolation of anti-inflammatory and antinociceptive compound from *P. zeylanica* leaf. *Pharm Biol* 2010; 48:381-387.
110. Chowdhury AK, Sushanta KC, Khan AK. Antifertility activity of *P. zeylanica* L. Root. *Indian J Med Res* 1982; 76:99-101.
111. Premakumari P, Rathinam K, Santhakumari G. Antifertility activity of Plumbagin. *Indian J Med Res* 1977; 65:829-838.
112. Bhargava SK. Effects of Plumbagin on reproductive function of male dog. *Indian J Exp Biol* 1984; 22:153-156.
113. Wang YC, Huang TL. Anti-*Helicobacter pylori* activity of *P. zeylanica* L. *Fems. Immunol Med Microbiol* 2005; 43:407-412.
114. Atkinson N. Antibacterial substances from flowering plants. 3. Antibacterial activity of dried Australian plants by a rapid direct plate test. *Aust J Exp Biol* 1956; 34:17-26.
115. Beg AZ, Ahmad I. Effect of *Plumbago zeylanica* extract and certain curing agents on multidrug resistant bacteria of clinical origin. *World Journal of Microbiology and Biotechnology* 2000; 16:841-844.
116. Lemma H *et al.* Anti-bacterial activity of *P. zeylanica* L. roots on some pneumonia causing pathogens. *Ethiop J Sci* 2002; 25:285-294.
117. Lakhmi VV, Padma S, Polasa H. Elimination of multidrug-resistant plasmid in bacteria by plumbagin, a compound derived from a plant. *Curr Microbiol* 1987; 16:159-161.
118. Olagunju JA, Jobi AA, Oyedapo OO. An investigation into the biochemical basis of the observed hyperglycemia in rats treated with ethanol root extract of *P. zeylanica*. *Phytother Res* 1999; 13:346-348.
119. Kubo I, Uchida M, Klocke JA. An insect ecdysis inhibitor from the Africa medicinal plant *Plumbago capensis*. *Agric Biol Chem* 1983; 47:911-913.
120. Kumar MS, Banerji A. Growth disruption in *Opisina arenosella* Walker by plumbagin, a natural occurring insect growth regulator. *Insect Sci Appl* 2002; 22:321-323.
121. Dai Y, Hou LF, Chan YP, Cheng L, Bur PP. Inhibition of immediate allergic reactions by ethanol extract from *P. zeylanica* stems. *Biol Pharm Bull* 2004; 27:429-432.
122. Karnick CR, Tiwari KC, Majumber R. Cultivation trials, pharmacognosy and ethnobotanical investigations of *P. zeylanica* L. (chitraka) of the Indian system of medicine. *Int J Crude Drug Res* 1982; 20:193-199.
123. Gundidza M, Manwa G. Activity of chloroform extract from *P. zeylanica* against *Neisseria gonorrhoeae*. *Fitoterapia*. 1990; 61:47-49.
124. Bamode RS, Shukla VN. Studies on fungitoxic properties of *P. zeylanica* L. extract. *Punjabrao Krishi Vidyapeeth Res J* 1974; 2:110-112.
125. Krishnaswamy M, Purushothaman KK. Plumbagin: A study of its anticancer, antibacterial and antifungal properties. *Indian J Exp Biol* 1980; 18:876-877.
126. Olagunju JA, Kazeem OW, Oyedapo OO. Further studies on the mechanism of carbohydrate intolerance induced in the rat by an ethanolic root extract of *P. zeylanica*. *Pharm Biol* 2000; 38:362-366.
127. Santhakumari G, Rathinam K. Anticoagulant activity of Plumbagin. *Indian J Exp Biol* 1978; 16:485-487.
128. Shen Z, Dong Z, Cheng P, Li L, Chen Z, Liu J. Effects of Plumbagin on platelet aggregation and platelet-neutrophil interactions. *Planta Med* 2003; 69:605-609.
129. Vijayakumar R, Senthilvelan M, Ravindran R, Devi RS. *P. zeylanica* action on blood coagulation profile with and without blood volume reduction. *Vascul Pharmacol* 2006; 45:86-90.
130. Azad C, Sushanta KC, Azadkhan AK. Antifertility activity of *Plumbago zeylanica* L. root. *Indian J Med Res* 1982; 76:99-101.
131. Yen-Ju H, Lei-Chwen L, Tung-Hu T. Measurement and pharmacokinetic study of plumbagin in a conscious freely moving rat using liquid chromatography/tandem mass spectrometry. *J Chromatography* 2006; 844:1-5.
132. Vanisree M, Hsin-Sheng T. Plant cell culture- an alternative and efficient source for the production of biologically important secondary metabolites. *Int J Appl Sci Eng* 2004; 2:29-48.
133. Katakai MS *et al.* Antibacterial activity, in vitro antioxidant activity and anthelmintic activity of methanolic extract of *P. zeylanica* L. Leaves. *J Pharmacy Res* 2010; 3:80-84.
134. Wang YC, Huang TL. Anti-*Helicobacter pylori* activity of *P. zeylanica* L. *Fems. Immunol Med Microbiol* 2005; 43:407-412.
135. Aqil F, Ahmad I. Antibacterial properties of traditionally used Indian medicinal plants. *Methods Find Exp Clin Pharmacol* 2007; 29:79-92.

136. Hiradeve S, Kishor D, Vijay K, Bibhilesh M. Evaluation of anticancer activity of *P. zeylanica* L. leaf extract. Int J Biomed Res 2010; 1:1-9.
137. Parimala R, Sachdanandam P. Effect of plumbagin on some glucose metabolizing enzymes studied in rats in experimental hepatoma. Mol Cell Biochem 1993; 125:59-63.
138. Vishnukanta S, Rana AC. Evaluation of anticonvulsant activity *P. zeylanica* L. leaf extract. Asian J Pharma Clin Res 2010; 3:76-78.
139. Zarmouh MM, Subramaniam K, Viswanathan S, Kumar PG. Cause and effect of *Plumbago zeylanica* root extract on blood glucose and hepatic enzymes in experimental diabetic rats. African J Microbiol Res 2010; 4:2674-2677.
140. Kumar G, Sharmila BG, Maheswaran R, Rema S, Rajasekara PM, Murugesan AG. Effect of *P. zeylanica* L. on blood glucose and plasma antioxidant status in STZ diabetic rats. J Nat Remedies 2007; 7:66-77.
141. Sharmila BG, Kumar G, Rajasekara PM. Hypoglycaemic effect of Kodiveli (*Plumbago zeylanica* L.) in STZ diabetic rats. J Theoret Exp Biol 2006; 3:1-5.
142. Koppula S, Ammani K, Varaprasad B, Bramhachari PV. Inhibition of plant pathogenic fungi by ethnobotanically selected plant extracts. J Pharmacy Res 2010; 3:2334-2336.
143. Kanchana N, Sadiq AM. Hepatoprotective effect of *P. zeylanica* on paracetamol induced liver Toxicity in rats. Int J Pharmacy Pharma Sci 2011; 3:151-154.
144. Mallikadevi T, Paulsamy SP. *zeylanica* L. - A potential plant for Antimicrobial activity. Plant Arch 2010; 10:547-550.
145. Demma J, Engidawork E, Hellman B. Potential genotoxicity of plant extracts used in Ethiopian traditional medicine. J Ethnopharmacol 2009; 122:136-142.
146. Natarajan KS, Narasimhan M, Shanmugasundaram KR, Shanmugasundaram ER. Antioxidant activity of a salt-spice-herbal mixture against free radical induction. J Ethnopharmacol 2006; 21(105):76-83.
147. Zahin M, Aqil F, Ahmad I. The *in vitro* antioxidant activity and total phenolic content of four Indian medicinal plants. Int J Pharmacy Pharma Sci 2009; 1:88-95.
148. Gebre-Mariam T, Neubert R, Schmidt PC, Wutzler P, Schmidtke M. Antiviral activities of some Ethiopian medicinal plants used for the treatment of dermatological disorders. J Ethnopharmacol 2006; 104:182-187.
149. Chen W, Yu Z, Li S. Effects of the water-soluble extracts from the single herb of gandunging against hepatitis-B virus *in vitro*. Zhong Yao Cai 1999; 22:463-465.
150. Ram A, Lauria P, Gupta R, Sharma VN. Hypolipidaemic effect of *Myristica fragrans* fruit extract in rabbits. J Ethnopharmacol 1996; 55:49-53.
151. Dwivedi S. Effect of *P. zeylanica* in hyperlipidemia rabbits and its modification by vitamin E. Indian J Pharmacol 1997; 29:138-139.
152. Abdul KM, Ramchender RP. Modulatory effect of plumbagin (5-hydroxy-2-methyl- 1,4-naphthoquinone) on macrophage functions in BALB/c mice. I. Potentiation of macrophage bactericidal activity. Immunopharmacol 1995; 30:231-236.
153. Poosarla A, Athota RR. Immunosuppressive properties of aqueous extract of *P. zeylanica* in Balb/c mice. J Med Plants Res 2010; 4:2138-2143.
154. Mittal V, Sharma S, Pawan JK, Anil HMJ. *P. zeylanica* roots: A potential source for improvement of learning and memory. Int J Pharma Bio Sci 2010; 6:1-6.
155. Powolny AA, Singh SV. Plumbagin- induced apoptosis in human prostate cancer cells is associated with modulation of cellular redox status and generation of reactive oxygen species. Pharma Res 2008; 25:2171-180.
156. Wang CC, Chiang YM, Sung SC, Hsu YL, Chang JK, Kuo PL. Plumbagin induces cell cycle arrest and apoptosis through reactive oxygen species/c-Jun Nterminal kinase pathways in human melanoma A375.S2 cells. Cancer Lett 2008; 259:82-98.
157. Hsu YL, Cho CY, Kuo PL, Huang YT, Lin CC. Plumbagin (5-hydroxy-2- methyl-1, 4-naphthoquinone) induces apoptosis and cell cycle arrest in A549 cells through p53 accumulation via c-Jun NH2-terminal kinase-mediated phosphorylation at serine 15 *in vitro* and *in vivo*. J Pharmacol Exp Ther 2006; 318:484-494.
158. Nazeem S *et al.* Plumbagin induces cell death through a copper-redox cycle mechanism in human cancer cells. Mutagenesis 2009; 24:413-418.
159. Aziz MH, Dreckschmidt NE, Verma AK. Plumbagin, a medicinal plant-derived naphthoquinone, is a novel inhibitor of the growth and invasion of hormone-refractory prostate cancer. Cancer Res 2008; 68:9024-9032.
160. Zhao YL, Lu DP. Effects of plumbagin on the human acute promyelocytic leukemia cells *in vitro*. Zhongguo Shi Yan Xue Ye Xue Za Zhi 2006; 14:208-211.
161. Tsai WJ *et al.* Seselin from *Plumbago zeylanica* inhibits phytohemagglutinin (PHA)-stimulated cell proliferation in human peripheral blood mononuclear cells. J Ethnopharmacol 2008; 119:67-73.
162. Sathya S *et al.* 3 β -hydroxylup- 20(29)-ene-27,28-dioic acid dimethyl ester, a novel natural product from *P. zeylanica* inhibits the proliferation and migration of MDA-MB-231 cells. Chemico-Biol Interact 2010; 188:412-420.
163. Ahmad A, Banerjee S, Wang Z, Kong D, Sarkar FH. Plumbagin-induced apoptosis of human breast cancer cells is mediated by inactivation of NF-kappaB and Bcl-2. J Cell Biochem 2008; 105:1461-1471.
164. Kuo PL, Hsu YL, Cho CY. Plumbagin induces G2-M arrest and autophagy by inhibiting the AKT/mammalian target of rapamycin pathway in breast cancer cells. Mol Cancer Ther 2006; 6:3209-3221.
165. Thasni KA, Rakesh S, Rojini G, Ratheeshkumar T, Srinivas G, Priya S. Estrogen-dependent cell signaling and apoptosis in BRCA1-blocked BG1 ovarian cancer cells in response to plumbagin and other chemotherapeutic agents. Annl Oncol 2008; 19:696-705.
166. Nair S, Nair RR, Srinivas P, Pillai MR. Radiosensitizing effects of plumbagin in cervical cancer cells is through modulation of apoptotic pathway. Mol Carcinog 2008; 47:22-33.
167. Srinivas G, Annab LA, Gopinath G, Benerji A, Srinivas P. Antisense blocking of BRCA1 enhances sensitivity to plumbagin but not tamoxifen in BG-1 ovarian cancer cells. Mol Carcinog 2004; 39:15-25.
168. Srinivas P, Gopinath G, Banerji A, Dinakar A, Srinivas G. Plumbagin induces reactive oxygen species, which mediate apoptosis in human cervical cancer cells. Mol Carcinog 2004; 40:201-211.

169. Naresh RA, Udupa N, Devi PU. Niosomal plumbagin with reduced toxicity and improved anticancer activity in BALB/C mice. *J Pharmacy Pharmacol* 1996; 48:1128–1132.
170. Sugie S *et al.* Inhibitory effects of plumbagin and juglone on azoxymethane-induced intestinal carcinogenesis in rats. *Cancer Lett* 1998; 127:177–183.
171. Yang SJ *et al.* Plumbagin activates ERK1/2 and Akt via superoxide, Src and PI3-kinase in 3T3-L1 cells. *Eur J Pharmacol* 2010; 638:21-28.
172. Paras J, Sharma HP, Fauziya B, Binit B, Soni K, Chanchala P. Pharmacological Profiles of Ethno-Medicinal Plant. *P. zeylanica*. *Int J Pharm Sci Rev Res* 2014; 24(1) n° 29:157-163.