PART 2: Random Variables and Integration

Mathematical reading notes — revised edition, 2026

Topics. Measurable functions (§3.2), random variables (§3.5), their laws (§3.6), nonnegative Lebesgue integrals and monotone convergence (§4.2), signed integrals and dominated convergence (§4.3), and expectation (§4.4).

1. Measurable functions (§3.2)

Let $(S_1, \mathcal{S}_1)$ and $(S_2, \mathcal{S}_2)$ be measurable spaces. A function $f : S_1 \to S_2$ is measurable when

$$f^{-1}(B) \in \mathcal{S}_1 \qquad \text{for every } B \in \mathcal{S}_2.$$

The condition concerns *preimages*, not images: taking the preimage of every measurable target set must give a measurable subset of the domain. This is also called $\mathcal{S}_1/\mathcal{S}_2$ measurability.

Example: indicator functions

For $A \in \mathcal{S}$, the indicator $\mathbf{1}_A : S \to \mathbb{R}$ is defined by $\mathbf{1}_A(x) = 1$ when $x \in A$ and 0 otherwise. For any Borel set $B \subset \mathbb{R}$, its preimage is one of the measurable sets

$$\mathbf{1}_A^{-1}(B) = \begin{cases} \varnothing, & 0 \notin B,\ 1 \notin B, \\ A^c, & 0 \in B,\ 1 \notin B, \\ A, & 0 \notin B,\ 1 \in B, \\ S, & 0 \in B,\ 1 \in B. \end{cases}$$

Thus indicator functions of measurable sets are measurable.

Generators of sigma-algebras

For a family $\mathcal{C} \subseteq \mathcal{P}(S)$, write

$$\sigma(\mathcal{C}) = \bigcap \{\mathcal{A} : \mathcal{A} \text{ is a sigma-algebra on } S,\ \mathcal{C} \subseteq \mathcal{A}\}.$$

It is the smallest sigma-algebra containing $\mathcal{C}$. For example, if $S = \{1, 2, 3\}$ and $\mathcal{C} = \{\{1\}\}$, then

$$\sigma(\mathcal{C}) = \{\varnothing, \{1\}, \{2, 3\}, S\}.$$

On the real line, the Borel sigma-algebra is generated by open sets, or already by the half-lines $(-\infty, a)$:

$$\mathcal{B}(\mathbb{R}) = \sigma\big(\{(-\infty, a) : a \in \mathbb{R}\}\big).$$

Proposition 1 (Testing measurability on a generator)**.** *If $\mathcal{C}$ generates $\mathcal{S}_2$, a function $f : S_1 \to S_2$ is measurable if and only if $f^{-1}(C) \in \mathcal{S}_1$ for every $C \in \mathcal{C}$.*

Proof. One implication follows from the definition. For the converse define $\mathcal{G} = \{B \subseteq S_2 : f^{-1}(B) \in \mathcal{S}_1\}$. Preimages preserve complements and countable unions, so $\mathcal{G}$ is a sigma-algebra. As $\mathcal{C} \subseteq \mathcal{G}$, it contains $\sigma(\mathcal{C}) = \mathcal{S}_2$. □

A continuous map between metric spaces, each equipped with its Borel sigma-algebra, is measurable: preimages of open sets are open, and the open sets generate the Borel sigma-algebra.

2. Random variables and their laws (§§3.5–3.6)

A probability space $(\Omega, \mathcal{A}, P)$ consists of a sample space, a sigma-algebra, and a probability measure. A random variable taking values in $(S, \mathcal{S})$ is a measurable map $X : \Omega \to S$. For a measurable set $B \in \mathcal{S}$,

$$P(X \in B) = P\big(X^{-1}(B)\big) = P\big(\{\omega : X(\omega) \in B\}\big).$$

When $S = \mathbb{R}^n$ with its Borel sigma-algebra, such a measurable map is called a random vector. In the real-valued case, $P(X = x)$ is shorthand for the probability of $X^{-1}(\{x\})$.

The *law*, or distribution, of X is the pushforward measure

$$\mu_X(B) = P(X \in B) = P\big(X^{-1}(B)\big), \qquad B \in \mathcal{S}.$$

This transfers probabilities on an abstract outcome space Ω to probabilities about the values taken by X.

Proposition 2. *The law μ_X is a probability measure on $(S, \mathcal{S})$.*

Proof. Because $X^{-1}(\varnothing) = \varnothing$ and $X^{-1}(S) = \Omega$, we have $\mu_X(\varnothing) = 0$ and $\mu_X(S) = 1$. If the measurable sets B_i are disjoint, their preimages are disjoint and

$$\mu_X\Big(\bigcup_i B_i\Big) = P\Big(\bigcup_i X^{-1}(B_i)\Big) = \sum_i P\big(X^{-1}(B_i)\big) = \sum_i \mu_X(B_i).$$

□

Proposition 3 (Realizing any probability law)**.** *For any probability measure μ on $(S, \mathcal{S})$, there is a random variable whose law is μ.*

Proof. Take $(\Omega, \mathcal{A}, P) = (S, \mathcal{S}, \mu)$ and the identity random variable $X(\omega) = \omega$. Then $X^{-1}(B) = B$ and $\mu_X(B) = \mu(B)$. Thus μ is the law of X. □

3. The nonnegative Lebesgue integral (§4.2)

Let $f : S \to [0, \infty]$ be measurable on a measure space $(S, \mathcal{S}, \mu)$. A *nonnegative simple function* is a finite sum $s = \sum_{i=1}^m a_i \mathbf{1}_{A_i}$ with $a_i \geq 0$ and measurable A_i. Define its integral as $\int s\, d\mu = \sum_{i=1}^m a_i \mu(A_i)$ for a disjoint simple representation. The integral of a nonnegative measurable function is

$$\int_S f\, d\mu = \sup\Big\{\int_S s\, d\mu : 0 \leq s \leq f,\ s \text{ nonnegative and simple}\Big\}.$$

Equivalently one can use lower sums over finite measurable partitions. This is analogous to approximating a graph from below using rectangles, except that measurable sets replace intervals.




Illustration: approximation by simple functions from below

Proposition 4 (Monotone convergence theorem)**.** *If $0 \leq f_1 \leq f_2 \leq \cdots$ are measurable and $f_n \uparrow f$ pointwise, then*

$$\int_S f_n\, d\mu \uparrow \int_S f\, d\mu.$$

We use this theorem without reproducing its full proof. It allows limits of increasing nonnegative functions to pass through the integral.

4. Signed integrals and dominated convergence (§4.3)

For a real-valued measurable function f, let

$$f_+ = \max\{f, 0\}, \qquad f_- = \max\{-f, 0\}, \qquad f = f_+ - f_-.$$

The integral is defined by $\int f\,d\mu = \int f_+\,d\mu - \int f_-\,d\mu$ whenever the difference is defined. It is *integrable* when

$$\int_S |f|\,d\mu < \infty.$$

For integrable f, g and real scalars a, b, the integral is linear: $\int (af + bg)\,d\mu = a\int f\,d\mu + b\int g\,d\mu$. Moreover, if $f \le g$ almost everywhere, then $\int f\,d\mu \le \int g\,d\mu$ (provided the integrals exist), and

$$\left|\int_S f\,d\mu\right| \le \int_S |f|\,d\mu.$$

Proposition 5 (Dominated convergence theorem)**.** *If the measurable functions f_n converge to f almost everywhere and there is a* single integrable *function $g \ge 0$ satisfying $|f_n| \le g$ almost everywhere for every n, then f is integrable and*

$$\lim_{n\to\infty} \int_S f_n\,d\mu = \int_S f\,d\mu.$$

The one-sided inequality $f_n < g$ alone is *not* sufficient. The same integrable majorant must control the absolute values of the entire sequence.

Example: Lebesgue but not Riemann integrable

On $[0,1]$, consider $f(x) = \mathbf{1}_{[0,1]\setminus\mathbb{Q}}(x)$, equal to one at irrational numbers and zero at rational numbers. Both rationals and irrationals are dense in every subinterval; hence every nontrivial Riemann partition has lower sum zero and upper sum one. The function is therefore not Riemann integrable. As the rationals form a countable set of Lebesgue measure zero,

$$\int_{[0,1]} f(x)\,dx = 1.$$

5. Expectation (§4.4)

For an integrable real random variable X on $(\Omega, \mathcal{A}, P)$, its expectation is

$$\mathbb{E}X = \int_\Omega X\,dP.$$

The properties of the Lebesgue integral immediately give linearity $\mathbb{E}(aX + bY) = a\mathbb{E}X + b\mathbb{E}Y$, monotonicity ($X \le Y \implies \mathbb{E}X \le \mathbb{E}Y$ for integrable variables), and $|\mathbb{E}X| \le \mathbb{E}|X|$.

Convergence theorems for expectation

- **Dominated convergence.** If $X_n \to X$ almost surely and $|X_n| \le Y$ almost surely for all n, where $Y \ge 0$ and $\mathbb{E}Y < \infty$, then $\mathbb{E}X_n \to \mathbb{E}X$.
- **Bounded convergence on a probability space.** If $X_n \to X$ almost surely and $|X_n| \le C$ for one finite constant C and all n, then $\mathbb{E}X_n \to \mathbb{E}X$.
- **Monotone convergence.** If $0 \le X_n \uparrow X$ almost surely, then $\mathbb{E}X_n \uparrow \mathbb{E}X$, with the possibility of $+\infty$.

Individual integrability of the X_n is not enough by itself to exchange expectation with the limit.

Reading note

The distinction between pointwise or almost-sure convergence and convergence of integrals becomes important later, when probability laws and second moments are passed to limits in the Wasserstein argument.