



Emergent climate and CO₂ sensitivities of net primary productivity in ecosystem models do not agree with empirical data in temperate forests of eastern North America

CHRISTINE R. ROLLINSON^{1,2} , YAO LIU³, ANN RAIHO⁴, DAVID J. P. MOORE³, JASON MCLACHLAN⁴, DANIEL A. BISHOP⁵, ALEX DYE⁶, JACLYN H. MATTHES⁷, AMY HESSL⁶, THOMAS HICKLER^{8,9}, NEIL PEDERSON⁵, BENJAMIN POULTER^{10,11}, TRISTAN QUAIFFE¹², KEVIN SCHAEFER¹³, JÖRG STEINKAMP⁸  and MICHAEL C. DIETZE¹

¹Department of Earth & Environment, Boston University, 685 Commonwealth Ave, Boston, MA 02215, USA, ²Morton Arboretum, 4100 Illinois Route 53, Lisle, IL 60532, USA, ³School of Natural Resources, University of Arizona, 1064 E. Lowell St., Tucson, AZ 85721, USA, ⁴Department of Biological Sciences, University of Notre Dame, 176 Galvin Life Science Center, Notre Dame, IN 46556, USA, ⁵Havard Forest, 324 N. Main St, Petersham, MA 10366, USA, ⁶Department of Geology and Geography, West Virginia University, P.O. Box 6300, Morgantown, WV 26506, USA, ⁷Department of Biological Sciences, Wellesley College, 106 Central Street, Wellesley, MA 02481, USA, ⁸Senckenberg Biodiversity and Climate Research Centre (BiK-F), Senckenberganlage 25, Frankfurt am Main D-60325, Germany, ⁹Department of Physical Geography and Geosciences, Goethe University, Altenhöferallee 1, Frankfurt am Main 60438, Germany, ¹⁰Biospheric Science Laboratory, NASA Goodard Space Flight Center, Greenbelt, MD 22071, USA, ¹¹Institute on Ecosystem and Department of Ecology, Montana State University, Bozeman, MT 59717, USA, ¹²Department of Meteorology, University of Reading, Earley Gate, PO Box 243, Reading, RG6 6BB, UK, ¹³National Snow and Ice Data Center, Cooperative Institute for Research in Environmental Sciences, University of Colorado, 449 UCB, Boulder, CO 80309, USA

Abstract

Ecosystem models show divergent responses of the terrestrial carbon cycle to global change over the next century. Individual model evaluation and multimodel comparisons with data have largely focused on individual processes at subannual to decadal scales. Thus far, data-based evaluations of emergent ecosystem responses to climate and CO₂ at multidecadal and centennial timescales have been rare. We compared the sensitivity of net primary productivity (NPP) to temperature, precipitation, and CO₂ in ten ecosystem models with the sensitivities found in tree-ring reconstructions of NPP and raw ring-width series at six temperate forest sites. These model-data comparisons were evaluated at three temporal extents to determine whether the rapid, directional changes in temperature and CO₂ in the recent past skew our observed responses to multiple drivers of change. All models tested here were more sensitive to low growing season precipitation than tree-ring NPP and ring widths in the past 30 years, although some model precipitation responses were more consistent with tree rings when evaluated over a full century. Similarly, all models had negative or no response to warm-growing season temperatures, while tree-ring data showed consistently positive effects of temperature. Although precipitation responses were least consistent among models, differences among models to CO₂ drive divergence and ensemble uncertainty in relative change in NPP over the past century. Changes in forest composition within models had no effect on climate or CO₂ sensitivity. Fire in model simulations reduced model sensitivity to climate and CO₂, but only over the course of multiple centuries. Formal evaluation of emergent model behavior at multidecadal and multicentennial timescales is essential to reconciling model projections with observed ecosystem responses to past climate change. Future evaluation should focus on improved representation of disturbance and biomass change as well as the feedbacks with moisture balance and CO₂ in individual models.

Keywords: climate change, climate sensitivity, ecosystem model, emergent response, model-data comparison, net primary productivity, paleoecology, tree rings

Received 13 May 2016; revised version received 10 December 2016 and accepted 14 December 2016

Introduction

Changes in temperature and precipitation regimes over the past millennium have been associated with shifts in ecosystem composition and structure in the

Correspondence: Christine R. Rollinson, tel. +1 630 719 2422, fax +1 630 725 2049, e-mail: crollinson@mortonarb.org

paleoecological record (Prentice *et al.*, 1991; Davis & Shaw, 2001; Shuman *et al.*, 2002; Clifford & Booth, 2015). Modern empirical data from experiment- and observation-based studies provide evidence that rapidly increasing temperatures, altered precipitation regimes, and rising atmospheric CO₂ concentrations are causing changes in ecosystem dynamics today (Boisvenue & Running, 2006; Morin *et al.*, 2009; Fisichelli *et al.*, 2013; Peñuelas *et al.*, 2013). However, the effects of climate change in long-lived ecosystems such as the temperate forests of the eastern and Midwestern United States remain contested because changes in forest composition and structure take decades to centuries to occur (Renwick & Rocca, 2014). These observation-based studies capture emergent, long-term ecosystem responses to climate and CO₂ variability that are the product of feedbacks and interactions among physiological and biogeochemical processes.

Terrestrial ecosystem models are used to make mechanistic, process-based projections of ecosystem response to changing climate and CO₂ in the past, present, and future. However, the mechanistic, bottom-up approach used to build and evaluate ecosystem models is mismatched in spatial and temporal scale from observations of ecosystem response to climate change. Differences in model parameterization or structural representation of physiological process can cause major divergences in the resulting ecosystem dynamics through time (De Kauwe *et al.*, 2013; Walker *et al.*, 2015). Model intercomparisons of ecosystem response to climate change over the next century typically show increased divergence of ecosystem dynamics at the end of the century (e.g., Friedlingstein *et al.*, 2014). Although most policy and management decisions occur at subcentennial temporal scales, life spans of common temperate tree species in the Northeastern United States can range from 120 years for *Betula papyrifera* to over 800 years for *Tsuga canadensis* (Burns & Honkala, 1990). This means that in the absence of widespread disturbance or management, even the model simulations of ecosystem response to climate change are far shorter than the multigenerational scales at which changes in forest composition or structure in response to shifts in climate occur. These slow, gradual shifts in forest composition or structure may mediate ecosystem sensitivity to climate and CO₂ in both models and reality. For example, mortality and recruitment are inherent processes underlying the types of ecosystem-scale responses to climate change observed in the past and occur at temporal and spatial scales beyond that which we can observe with current available ecological data. This can then cause the observed responses of ecosystems to past climate change over multiple

centuries to be different from that observed in shorter studies focused on a few decades.

Tree rings provide annually resolved records of individual and forest response to environmental variation over the temporal scales of decades and centuries across temperate forests. Tree rings have been used to understand forest responses to climate variability at individual sites and entire continents (Williams *et al.*, 2013; Charney *et al.*, 2016; D'Orangeville *et al.*, 2016), but these approaches have been largely disconnected from assessments of how climate change impacts ecosystems in models. Tree rings are increasingly used to quantify and constrain components of the terrestrial carbon cycle in forests as well as individual- and forest-level responses to climate (Graumlich *et al.*, 1989; Davis *et al.*, 2009; Babst *et al.*, 2013; Dye *et al.*, 2016). These efforts are critical for understanding long-term forest responses to climate variability as multiple field experiments have indicated that long-term individual and ecosystem responses to warming and elevated CO₂ diverge from initial responses found in the first few years of manipulation (Melillo *et al.*, 2002; Hollister *et al.*, 2005; Matesanz *et al.*, 2009; Norby *et al.*, 2010). This long-term evaluation of climate impacts on forest ecosystems is particularly important for disentangling the simultaneous directional shifts in temperature, precipitation, CO₂, and disturbance that complicate much of modern ecological research (Foster *et al.*, 1998; Turner *et al.*, 2003; Gómez-Aparicio *et al.*, 2011).

This article compares emergent sensitivity of net primary productivity (NPP) to climate and CO₂ in ecosystem models to those found in tree-ring data at multiple temporal scales. The goals of this article are as follows: (i) determine whether annual ecosystem NPP sensitivity to climate and CO₂ in ecosystem models matches those found in individual tree- and forest-level tree-ring data; (ii) compare the climate and CO₂ sensitivities in models and data from short temporal extents where patterns are dominated by interannual climate variability with those from centennial-scale records that contain low-frequency climatic shifts; and (iii) quantify the contribution of differences in model responses to climate and CO₂ to model ensemble uncertainty in NPP dynamics through time. To achieve these goals, we analyze the temporal trends of NPP in ten ecosystem models that have been run from 850 to 2010 A.D. at six temperate forest sites in the Upper Midwestern and Northeastern United States. The influences of growing season temperature, precipitation, and CO₂ on change in model NPP are then compared to those found in plot-level NPP reconstructions from tree rings at two sites as well as raw ring widths from plot-based sampling and the International Tree Ring Databank (ITRDB) at four sites. Climate and CO₂ responses are

analyzed for three temporal extents: (i) the scale of past paleoecological responses to climate change (850–2010 A.D.); (ii) the period of historical temperature and precipitation records (1901–2010 A.D.); and (iii) the scale of robust modern ecological data (1980–2010 A.D.). We use these results to explain model-data discrepancies in regional drivers of NPP through time as well as quantify the drivers of uncertainty within the model ensemble over the past millennium.

Materials and methods

Model experimental overview

We modeled ecosystem carbon flux and composition change in response to climate variation from 850 to 2010 A.D. at six sites in the Northeastern and Upper Midwestern United States. These sites are located on an east–west gradient that represent present-day differences in temperature, precipitation, and forest types (Table 1 and Appendix S1). Models were presented with common, continuous 6-hourly meteorological climate and monthly CO₂ forcing data. CO₂ was taken from the Law Dome time series (850–2000) and the NOAA Mauna Loa record (2001–2010) (Keeling *et al.* 2009; Wei *et al.*, 2014). Climate drivers included air temperature, water-equivalent precipitation rate, total incoming shortwave radiation, incoming longwave radiation, surface air pressure, specific humidity, and wind speed. The continuous, 6-hourly 850–2010 meteorological driver set was developed using an artificial neural network to spatially and temporally down-scale output from CCSM4 output from the Paleoclimate Modeling Intercomparison Project, Phase III (PMIP3) past millennium simulations (850–1849, Crucifix *et al.*, 2012) and Coupled Model Intercomparison Project, Phase 5 (CMIP5) simulations (1850–1900, Taylor *et al.*, 2012) using 6-hourly, 0.5-degree CRUNCEP data (1901–2010, Wei *et al.*, 2014) according to Kumar *et al.* (2012). Due to mismatches at transitional periods between CCSM4 and CRUNCEP products, all variables except wind were bias-corrected to avoid sharp jumps in climatology in the time series. Temperature was corrected as an additive bias, whereas short- and longwave radiation, and precipitation were corrected using a ratio bias to conserve the hydrological sensitivity among meteorological variables (Hempel *et al.*, 2013).

Ten models representing variants of five independent ecosystem models completed simulations of ecosystem dynamics at all six sites from 850 to 2010 A.D. and provided monthly or annual-resolution output on composition and the carbon cycle (Table 2). Full model protocol can be found in Appendix S2. Dynamic vegetation models allow plant communities to self-assemble and change through time, while static vegetation models prescribe plant functional types (PFTs) by fractional area based on potential vegetation distribution from Ramankutty & Foley (1999). All models were spun up to steady state at 850 A.D. by cycling the first 20 years of forcing data (850–869 A.D.) with a constant CO₂ concentration of 277 ppm. Not all models used all meteorological variables as drivers. LINKAGES was the only model to not include CO₂ as a driver and only includes temperature and precipitation drivers, whereas all other models included at a minimum, temperature, precipitation, shortwave radiation, and CO₂. Even though LINKAGES does not include CO₂ as a driver of ecosystem dynamics, it was included in our analyses to help isolate potential confounding effects of increasing CO₂ with other drivers of ecosystem change.

Our analyses focus on two model characteristics (dynamic vegetation and fire) and the dynamics of two ecosystem properties (composition and aboveground biomass) that are linked to slow processes associated with ecosystem responses to climate and CO₂ change (Table 2). However, because PFTs in all models except for SiBCASA, a biome-based model, respond to climate independently, they may still experience shifts in the relative community composition if defined by fluctuations in PFT biomass. Composition variability through time was quantified as the standard deviation of percent evergreen biomass through time within each site for each model. Although fire was possible in ED, CLM, and LPJ model variants, the conditions for fire were not triggered in ED2 and ED2-LU, so the effects of fire were only present in four models (CLM-BGC, CLM-CN, LPJ-WSL, and LPJ-GUESS). Fire in ecosystem models primarily affects ecosystems by reducing biomass, but mortality or changes in growth can also cause similar biomass fluctuations in all models. To calculate biomass variability, biomass was first normalized to the site mean so that biomass through time was expressed as a percent of mean. Biomass variability was then calculated as the standard deviation of this normalized biomass through time at each site for each model. All models except JULES-STATIC used aboveground biomass (AGB) as the measure of biomass variability. Leaf area index was used as

Table 1 Location, modern forest type, and mean model driver growing season (May through September), temperature and precipitation for the six modeling locations, and data types that were available at each site for sensitivity analysis

Name	Longitude	Latitude	Forest type	Temp (°C)	Precip (mm)	Data types
Demming Lake	−95.17	47.17	Mixed	15.6 ± 1.0	375 ± 74	M, RW
Billy's Lake	−94.58	46.28	Mixed	16.4 ± 1.0	398 ± 81	M
UNDERC	−89.53	46.22	Mixed	14.3 ± 0.9	411 ± 74	M, RW
Minden Bog	−82.83	43.61	Evergreen	16.4 ± 0.8	375 ± 65	M
Harvard Forest	−72.18	42.54	Deciduous	15.6 ± 0.7	520 ± 86	M, RW, NPP _{TR}
Howland Forest	−68.73	45.25	Mixed	13.9 ± 0.8	492 ± 80	M, RW, NPP _{TR}

Temperature and precipitation values presented are mean ± standard deviation from 850 to 2010 A.D. Data type codes are as follows: M, ecosystem models; RW, raw tree-ring widths; NPP_{TR}, tree-ring NPP reconstruction.

Table 2 List of models and key model characteristics used in analyses

Model	Vers.	Vegetation scheme	Composition variability (%)	Fire occurrence	Biomass variability (%)	Citation
CLM-BGC	4.5	Static	32	Yes	22	Oleson <i>et al.</i> (2010)
CLM-CN	4.5	Static	29	Yes	17	Oleson <i>et al.</i> (2010)
ED2	2.1	Dynamic	20	No	20	Medvigy <i>et al.</i> (2009)
ED2-LU	2.1	Dynamic	21	No	19	Medvigy <i>et al.</i> (2009)
JULES-STATIC	4.1	Static	25	No	3	Best <i>et al.</i> (2011), Clark <i>et al.</i> (2011)
JULES-TRIFFID	4.1	Dynamic	02	No	6	Best <i>et al.</i> (2011), Clark <i>et al.</i> (2011)
LINKAGES	1.0	Dynamic	22	No	19	Post & Pastor (1996)
LPJ-GUESS	3.1	Dynamic	35	Yes	24	Sitch <i>et al.</i> (2003), Gerten <i>et al.</i> (2004) and Smith <i>et al.</i> (2014)
LPJ-WSL	1.0	Dynamic	28	Yes	12	Sitch <i>et al.</i> (2003), Gerten <i>et al.</i> (2004)
SiBCASA		Static	00	No	4	Schaefer <i>et al.</i> (2008)

Static vegetation models were prescribed site composition based on modern forest-type information (Table 1). However, because composition was assessed as fraction biomass or leaf area (LPJ & JULES models), relative composition could shift through time. Composition variability and biomass variability were quantified as the standard deviation of the relative fraction evergreen or biomass through time over the full modeling temporal extent. For fire occurrence, 'No' indicates no fire occurred in the model simulations at any locations. Version numbers are not tracked in SiBCASA.

a proxy for biomass in JULES-STATIC because there is no biomass or vegetation dynamics beyond leaf area simulated in JULES without coupling to the TRIFFID model. In other models, LAI had a mean correlation with AGB of 0.73 ± 0.18 and ranged from 0.55 in CLM-BGC to 0.99 in LINKAGES. In JULES-TRIFFID, the correlation between LAI and AGB was 0.93.

Empirical data

We used two types of tree-ring data to provide an empirical estimate of climate sensitivity for trees and forests from our study region. First, stand-level aboveground NPP was reconstructed from sampling mapped fixed-area plots according to Dye *et al.* (2016), which enable aboveground NPP to be calculated on a per area basis that is comparable to ecosystem model output (Table 1). NPP was reconstructed from five plots in two separate stands at Harvard and three plots in a single stand at Howland. A nested sampling scheme was used at both locations where two to three increment cores were taken from all trees greater than 10 cm diameter at breast height (DBH) in a 13-m-radius plot and trees greater than 20 cm DBH were cored in a 20-m-radius plot. Growth measurements from all cores for each were averaged and used to reconstruct DBH through time that was then converted to aboveground biomass and annual aboveground biomass increment (proportional to NPP) for each tree and plot using species-specific equations that can be found in the appendix of Dye *et al.* (2016). Due to the potential decline in sample replication, and the potential for unmeasured mortality, analyses of tree-ring NPP sensitivity to climate were restricted to 1980–2010 (Clark *et al.*, 2001; Foster *et al.*, 2014).

In addition to tree-ring estimates of NPP, we also analyzed the sensitivity of raw ring width (RW) to climate from the trees used to generate the NPP estimates as well as from nine datasets from the International Tree Ring Databank (ITRDB)

(Grissino-Mayer & Fritts, 1997). Records from the ITRDB were found in close proximity to the Demming Lake, Howland, and UNDERC sites, allowing more robust comparisons of climate sensitivity across the modeled study sites (Appendix S3). In cases where multiple cores existed for a tree, the mean of ring width for each year was used. Although we include supplemental sensitivity analyses using *a priori* detrended ring-width index (RWI, Appendix S4), we chose raw ring widths as our response variable rather than RWI or basal area increment (BAI) for two reasons. First, we chose not to detrend ring widths *a priori* for our primary analyses to preserve potential low-frequency effects of changing climate or CO₂ in the modern era that might be confounded with increasing stem size. Second, BAI was not used because measurements of actual diameters were not available for ITRDB trees. The CRUNCEP climate records used to drive the models were matched to the tree-ring data for each site.

Sensitivity analysis

Model and tree-ring sensitivities to climate were analyzed using generalized additive models (GAMs) where NPP or RW is the sum of nonlinear temperature, precipitation, and CO₂ effects. In these analyses, we used site-level annual NPP as the response variable for ecosystem models, plot-level annual aboveground woody increment for tree-ring NPP, and mean raw ring width for each individual in the RW analysis. Temperature, precipitation, and CO₂ effects were estimated with three-knot thin plate regression splines, while an additional size effect was fit with three-knot cubic smoothing spline using the *gam* function of the *mgcv* package in R 3.2.3 (Wood, 2012). Even though LINKAGES lacks CO₂ as a model driver, this effect was included in our statistical analyses as a test of the statistical model's ability to attribute variation in NPP to climate and CO₂. The size term was included to account for effects of biomass (models, tree-ring NPP) or individual size

(raw ring width) that could alter the maximum potential NPP or ring width independent of climate. Total site or plot above-ground biomass was used for the size effect in models and tree-ring NPP GAMs, while reconstructed DBH was used for size in the RW GAMs. DBH was reconstructed by subtracting ring widths from the DBH at the time of sampling at Harvard and Howland, but was reconstructed by summing ring widths from the ITRDB records, where no DBH information was available. Because ring-width sums may not accurately reflect the true DBH of ITRDB trees, in RW models the size effect was fit separately for each ITRDB core. This effectively individually detrends ITRDB cores by removing effects of increasing size through time while simultaneously assessing the climate and CO₂ sensitivity of both ITRDB and plot-based samples. We also performed supplemental analyses with *a priori* detrended RWI using a two-thirds spline that removes long-term trends in tree-ring series (Appendix S5). Because *a priori* detrending removes age- and size-based trends in the data, the size factor was removed from our climate and CO₂ GAM in RWI analyses. We also include an analysis with all datasets restricted to Harvard and Howland sites where tree-ring NPP records are available (Appendix S6).

We normalized the sensitivities of model NPP and tree-ring width to climate and CO₂ for all analyses to facilitate comparison of model agreement of change in NPP through time and relative influences of temperature, precipitation, and CO₂ as drivers of that change. In all cases, the response variable of NPP or RW was converted to a percent deviation from the model or tree-ring dataset mean. We assessed model agreement of both change in NPP and drivers of change as the standard deviation around the ensemble mean before and after 1901, where CO₂ begins to sharply increase. To prescribe the effects of individual drivers on change in NPP or RW through time, we quantified the relative effects of temperature, precipitation, and CO₂ on NPP or RW in each year. The CO₂ effect was offset from the GAM-estimated effect so that the beginning of simulation CO₂ concentration in 850–855 A.D. received a weight of zero.

Models were analyzed at three temporal scales to determine whether observations from short periods in the modern era were able to capture the relationships between climate and CO₂ change and ecosystem responses over periods with low-frequency, directional shifts in climate. These three temporal extents are as follows: (i) scale of robust, modern empirical records (1980–2010), (ii) the scale of observational climate data (1901–2010), and (iii) the full model simulation extent (850–2010). This model-centric analysis was complemented with comparisons to the shifts in sensitivity seen in the RW models at the two shorter scales. Because GAMs center the spline-based effects on the means of the given data (i.e., the mean of observed temperature, precipitation, and CO₂), all three temporal scales were recentered on their respective 1980–2010 means to facilitate cross-scale comparisons. Differences in climate and CO₂ sensitivities within and across temporal scales both within individual models and at the ensemble level were assessed by comparing the mean slopes and 95% confidence intervals around each effect. We used an ANOVA to compare variability in the model ensemble across temporal scales. In

this ANOVA, we compared the absolute deviation of mean climate and CO₂ sensitivity for each model to that of the ensemble mean at each scale.

We also quantified the effects of slow ecosystem processes among models on NPP sensitivity to climate and CO₂ using linear regression. Here, we considered two categorical characteristics of models as well as two related continuous variables to identify trends in model dynamics and sensitivity to climate and CO₂. Categorical model characteristics included vegetation scheme (static or dynamic) and the presence of fire (yes or no), while composition (fraction evergreen) and biomass variability described similar model dynamics as continuous effects. Each characteristic of slow ecosystem processes was correlated with the mean slopes of model sensitivities to temperature, precipitation, and CO₂. We performed this correlation analysis at all three temporal scales to determine whether the effects of these slow processes were significant at particular, characteristic scales.

Results

Spatial and temporal patterns of NPP in ecosystem models

Terrestrial ecosystem models disagreed about both the spatial patterns of NPP in the Northeastern and Upper Midwestern United States as well as the change in NPP through time (Fig. 1). Midwestern sites tended toward lower NPP than those further east with ensemble-mean NPP ranging from $3.60 \pm 2.14 \text{ MgC m}^{-2} \text{ yr}^{-1}$ (mean \pm SD) at Billy's Lake to $6.0 \pm 2.5 \text{ MgC m}^{-2} \text{ yr}^{-1}$ at Harvard Forest. ED-LU had the highest mean NPP of the ensemble ($7.12 \pm 2.80 \text{ MgC m}^{-2} \text{ yr}^{-1}$), and JULES-STATIC had the lowest ($1.51 \pm 1.17 \text{ MgC m}^{-2} \text{ yr}^{-1}$), but this pattern varied across individual sites and times. Tree-ring estimates of NPP (NPP_{TR}) at both Harvard and Howland were much lower than predicted by models (Fig. 1). From 1980 to 2010, NPP_{TR} was $1.73 \pm 0.27 \text{ MgC ha}^{-1} \text{ yr}^{-1}$ at Harvard and $1.17 \pm 0.13 \text{ MgC ha}^{-1} \text{ yr}^{-1}$ at Howland. The mean NPP of the model ensemble from 1980 to 2010 was $7.47 \pm 3.73 \text{ MgC ha}^{-1} \text{ yr}^{-1}$ at Harvard and $7.07 \pm 3.93 \text{ MgC ha}^{-1} \text{ yr}^{-1}$ at Howland. NPP_{TR} was lower than the lowest model NPP at both sites ($2.90 \pm 0.38 \text{ MgC ha}^{-1} \text{ yr}^{-1}$ at Harvard and $3.48 \pm 0.30 \text{ MgC ha}^{-1} \text{ yr}^{-1}$ at Howland).

To facilitate comparisons of the impacts of climate and CO₂ on NPP, through time, we analyzed percent change in NPP relative to each model's mean. Once relativized, models and tree rings displayed similar levels of variability in NPP and ring width: Model NPP variability was $16 \pm 8\%$, NPP_{TR} was 12% , and RW was 20% . Even though ED2-LU had the highest NPP variability in absolute terms, with a standard deviation through time of $1.83 \text{ MgC m}^{-2} \text{ yr}^{-1}$,




Fig. 1 Net primary production (NPP) across all sites for the full modeling temporal extent (850–2010) for all ten ecosystem models. Sites are as follows: (a) Demming Lake, (b) Billy's Lake, (c) UNDERC, (d) Minden Bog, (e) Harvard Forest, and (f) Howland Forest. Lines indicate the 10-year running means. Black dots at Harvard and Howland indicate tree-ring estimates of NPP.

JULES-STATIC displayed higher temporal NPP variability relative to its mean (29%). In contrast, SiBCASA showed the least absolute and relative NPP variability through time: $0.28 \text{ MgC m}^{-2} \text{ yr}^{-1}$, 5% of its mean. The higher relative NPP variability of JULES-STATIC was accompanied by the most stable biomass (temporal standard deviations of 3%), while JULES-TRIFFID showed the least temporal variability of composition with only a mean 2% variability through time (Table 2). LPJ-GUESS, whose mean NPP and NPP variability were close to the ensemble mean, had the highest variability of both composition and biomass (35% and 24%, respectively).

NPP and RW sensitivity to climate and CO₂

Comparisons between models and tree-ring data at subcentennial temporal scales reveal large disparities between the effects of climate and CO₂ on NPP and tree growth (Fig. 2). At the 1980–2010 temporal extent, the ensemble of models and tree-ring data agreed on a positive relationship between NPP or growth and precipitation, but mean model sensitivity ($0.09 \pm 0.05\% \text{ mm}^{-1} \text{ yr}^{-1}$) was nine times higher than both tree-ring NPP and ring widths (both $0.01\% \text{ mm}^{-1} \text{ yr}^{-1}$). The overall signs of sensitivity were opposite between models and tree-ring data for temperature (model ensemble = $-0.99 \pm 4.03\% \text{ }^{\circ}\text{C}^{-1}$, RW = $11.0\% \text{ }^{\circ}\text{C}^{-1}$, NPP_{TR} = $14.2\% \text{ }^{\circ}\text{C}^{-1}$) and CO₂ (model ensemble = $0.20 \pm 0.16\% \text{ ppm}^{-1}$, RW = $-0.7\% \text{ ppm}^{-1}$, NPP_{TR} = $0.01\% \text{ ppm}^{-1}$). However, the 95% confidence interval for NPP_{TR} does encompass 0 and is consistent with relatively CO₂-

insensitive models such as LPJ-WSL. *A priori* detrending of ring width to ring-width index dramatically reduced temperature, precipitation, and CO₂ sensitivity in the individual-based tree-ring analysis, causing temperature and CO₂ sensitivities to be more consistent with model sensitivities, but less consistent with NPP_{TR} (Appendix S5). Trends within and among the sensitivities of models and tree-ring data were similar between analyses including all sites and when analyses were restricted to just Harvard and Howland (Appendix S6). Increasing temporal scale had relatively minor effects on climate sensitivity in model NPP and increased sensitivity in RW (Fig. 2 and Appendix S7). This resulted in greater consistency among precipitation responses in RW and models, but continued model-data disagreement in temperature and CO₂ at the 1901–2010 scale. Temperature sensitivity of NPP in LINKAGES was a notable exception to this trend and showed similar positive effects of lower growing season temperatures as the tree-ring datasets, but then diverged and displayed negative effects of warm temperatures.

Spatial and temporal variability in NPP among models corresponded to differences among models in emergent ecosystem sensitivities to temperature, precipitation, and CO₂. At the full temporal extent of our model simulations (850–2010), models showed consistent positive effects of precipitation and CO₂ on NPP, although with varying magnitudes of the effect (Fig. 2). For precipitation, the ensemble of models had an average $0.13 \pm 0.11\%$ increase in NPP per millimeter precipitation, but individual models showed




Fig. 2 Normalized sensitivity of net primary production (NPP) and tree-ring width to growing season temperature, growing season precipitation, and CO₂ across three temporal scales. Climate and CO₂ effects have been normalized to the mean NPP or ring width for each model to facilitate comparison across models and data of different scales. Colored lines indicate the mean estimated response, and color shaded area indicate a 95% confidence interval around that response. Gray shaded regions indicate the ranges of climate or CO₂ not observed in the models at that temporal scale and are presented only for aiding visualization and as such differences among models in these regions should not be interpreted.

sensitivities ranging from $0.02 \pm 0.03\% \text{ mm}^{-1} \text{ yr}^{-1}$ in SiBCASA to $0.32 \pm 0.16\% \text{ mm}^{-1} \text{ yr}^{-1}$ in JULES-STATIC. Ensemble NPP sensitivity to CO₂ was $0.36 \pm 0.34\% \text{ ppm}^{-1}$ with individual effects in models with CO₂ ranging from $0.06 \pm 0.04\% \text{ ppm}^{-1}$ in CLM-BGC to $1.03 \pm 0.20\% \text{ ppm}^{-1}$ in JULES-STATIC. LINKAGES, which lacks CO₂ effects in its model structure, supports the ability of the GAM approach to correctly attribute impacts of climate and CO₂ on NPP by displaying a small NPP response of $-0.01 \pm 0.00\% \text{ ppm}^{-1}$. Temperature was the only effect included in models to show differences in the direction of NPP sensitivity. Most models had a generally negative temperature effect, and the ensemble-mean effect was a decrease of $2.37 \pm 9.14\%$ in NPP per °C. JULES-STATIC showed the strongest effect with a $23.48 \pm 0.89\%$ decrease per °C. Overall, LINKAGES had a positive temperature effect ($14.77 \pm 19.43\% \text{ }^{\circ}\text{C}$), although this effect became negative at higher temperatures. Over the full course of model simulations, the presence of fire was correlated with decreased sensitivity to temperature ($t = -2.3$, $P = 0.03$) and CO₂ ($t = -2.8$, $P < 0.01$). Decreased model NPP sensitivity to temperature was also associated with increased biomass variability through time ($t = -2.7$, $P = 0.01$).

Models with more variable composition over multiple centuries also tended to be less sensitive to CO₂, but this trend was not significant ($t = -2.0$, $P = 0.06$).

An analysis of variance indicated there was greater agreement among model sensitivities to precipitation and CO₂ at the two shorter temporal scales than at the multicentennial (850–2010) extent (Fig. 2; precipitation: $t = 4.6$, $P < 0.01$, CO₂: $t = 4.0$, $P < 0.01$), but temperature showed similar ensemble variability across temporal scales ($P > 0.05$). The presence of fire in ecosystem models correlated with reduced sensitivity to temperature, precipitation, and CO₂, but only at the multicentennial temporal scale (Table 3). Slow ecosystem processes had weakened effects on model NPP sensitivity to climate and CO₂ at the shorter temporal scales of empirical data and were not significantly associated with patterns in model sensitivities at these multicentennial scales (Table 3). Similarly, models with high temporal variability of biomass were also less sensitive to temperature, but only over the course of multiple centuries. Factors associated with changes in composition including composition stability and whether a model had dynamic or static vegetation had no correlation with climate sensitivity in our model ensemble.

Table 3 ANOVA effect sizes of changes in climate and CO₂ sensitivity for four key characteristics of ecosystem models and ecosystem dynamics

Effect	Character	1980–2010	1901–2010	850–2010
Temperature	Static vegetation	3.8 ± 3.68	4.13 ± 3.68	4.17 ± 3.68
	Composition Var.	12.67 ± 14.92	7.67 ± 17.3	−8.16 ± 20.47
	Fire occurs	0.99 ± 3.31	−1.4 ± 3.31	−8.18 ± 3.31*
	Biomass Var.	−0.67 ± 7.20	−9.89 ± 8.82	−20.03 ± 9.15*
Precipitation	Static vegetation	−0.01 ± 0.20	0.15 ± 0.20	0.06 ± 0.20
	Composition Var.	0.03 ± 0.20	−0.06 ± 0.23	−0.25 ± 0.27
	Fire occurs	−0.01 ± 0.04	−0.02 ± 0.04	−0.11 ± 0.04*
	Biomass Var.	0.06 ± 0.10	0.01 ± 0.12	−0.2 ± 0.13
CO ₂	Static vegetation	−0.01 ± 0.20	0.15 ± 0.20	0.06 ± 0.20
	Composition Var.	−0.34 ± 0.70	−0.33 ± 0.81	−1.81 ± 0.96
	Fire occurs	−0.17 ± 0.15	−0.15 ± 0.15	−0.45 ± 0.15*
	Biomass Var.	0.09 ± 0.39	−0.13 ± 0.48	−0.61 ± 0.50

Effects are expressed as mean change in normalized NPP (%) per unit climate effect ± standard error. Static vegetation and fire effects are relative to dynamic vegetation scheme and absence of fire, respectively.

*Bold indicate significance at $P < 0.05$.

Drivers of model ensemble patterns agreement through time

Differences in model sensitivity to increasing CO₂ explained increasing model ensemble uncertainty in change in NPP since 1900 (Fig. 3). Model ensemble variability of normalized NPP prior to 1900 was $11.0 \pm 5.3\%$ (Fig. 3b). After 1900, this variability more than doubled to $20.6 \pm 7.4\%$. This shift in model agreement occurred as NPP in most models shifted from being primarily influenced by precipitation to showing varying degrees of CO₂ enhancement (Fig. 3b, c). In the 1980–2010 extent, the effect of CO₂ on NPP in models that include dynamic CO₂ ranged from +2% to +110% with model variability in CO₂ response proportionate to that of intermodel NPP deviation during this time period (CO₂ variability = $29 \pm 3\%$; NPP variability = $28 \pm 5\%$, Fig. 3c). LINKAGES was the only model to not include a CO₂ driver, and our statistical method misattributed a slight 0.6% decrease in NPP to increasing CO₂.

Discussion

Analysis of emergent responses to temperature, precipitation, and CO₂ at multidecadal and multicentennial timescales reveals many inconsistencies among models and data. The ten terrestrial ecosystem models included in our study showed variability in both mean NPP across space and relative changes in NPP through time (Fig. 1). Similar discrepancies have been widely observed in other multimodel comparisons, which have attributed differences among models to numerous causes including ecosystem feedbacks and uncertainties in model processes and parameterization

(Piao *et al.*, 2013; Friedlingstein *et al.*, 2014; Walker *et al.*, 2014). We found widely varying patterns of climate and CO₂ effects among models and data across multiple temporal scales (Fig. 2). However, the ensemble of models displayed less variation in climate and CO₂ responses at short temporal scales. Small differences in model structure and parameterization compound over time and drive model divergence, skewing model projections at ecology- and policy-relevant timescales. Fire and changes in biomass are only associated with differences in precipitation and CO₂ sensitivity among models at the multicentennial scale, indicating that feedbacks involving disturbance and biomass only have discernible impacts on climate and CO₂ over multiple forest generations (Table 3). Although model responses to precipitation are the most variable, responses to CO₂ are the greatest source of ensemble divergence of the past 100 years.

Models and empirical datasets have widely varying estimates of baseline NPP, so standardization is important to compare the impacts of climate and CO₂ through time across datasets. To be consistent across all models in the ensemble, we analyzed total ecosystem NPP, which includes the total carbon allocated to both above- and belowground tissues. Most models in our ensemble simulated NPP values within the range of what has been observed for forests common in the Northeastern and Upper Midwestern United states. Representative field-based estimates of NPP for our study region range from $1.3 \text{ MgC ha}^{-1} \text{ yr}^{-1}$ in mature boreal forests to as high as $8.7 \text{ MgC ha}^{-1} \text{ yr}^{-1}$ in the oak–hickory forests that dominate further south when a 50% carbon content of biomass is assumed (Jenkins *et al.*, 2001; Luyssaert *et al.*, 2007; Goulden *et al.*, 2011).




Fig. 3 (a) 10-year running mean net primary productivity for ten terrestrial ecosystem models (lines) with 95% confidence intervals from among-site variability (shaded areas). (b) 10-year running mean of the ensemble mean of normalized model net primary production (NPP) (colored line) with shaded 95% confidence interval (from spread among models) showing relative change in NPP through time in the model ensemble. Color of each line indicates the relative control of temperature (red), precipitation (blue), and CO₂ (green) on change in relative NPP, and dashed line indicates the normalized mean of 100%. (c) Mean effect of temperature (red), precipitation (blue), and CO₂ (green) on normalized NPP in the model ensemble. Lines indicate the ensemble mean, and the shaded area indicates the 95% confidence intervals.

Even if the models were perfectly accurate, their values would still be greater than the aboveground-only NPP quantified through tree rings in our study, which is estimated to be between 45% and 65% of the total carbon. Furthermore, our tree-ring productivity estimates only include the aboveground biomass increment from 1 year to the next, which will not capture any biomass that is produced and turns over within a year. A recent comparison of tree-ring estimates of NPP at Harvard Forest indicated good agreement with repeat forest censuses (Eisen & Plotkin, 2015; Dye *et al.*, 2016), but is substantially lower than values estimated through remote sensing for coniferous and deciduous forests at the same site (5.5 and 6.8 $\text{MgC ha}^{-1} \text{ yr}^{-1}$, respectively; Turner *et al.*, 2005). To reconcile total and aboveground NPP estimates, better quantification of amounts and controls of allocation and turnover among tissues is essential (Jenkins *et al.*, 2001; Litton *et al.*, 2007; Luysaert *et al.*, 2007). Some empirical studies suggest that the fraction of NPP allocated to aboveground wood could vary from year to year due to tree maturation,

increasing CO₂, or interannual climate variability (DeLucia *et al.*, 2005; Norby *et al.*, 2005; Doughty *et al.*, 2014). Although some ecosystem models allocate carbon in response to changing resource limitations (De Kauwe *et al.*, 2014), most do not and the models used in this analysis are based on simple allometric rules. Despite these inconsistencies in NPP values among models and empirical datasets, it is informative to compare the emergent sensitivities of productivity to climate and CO₂ across datasets.

Emergent model NPP sensitivities to temperature and precipitation are not supported by the responses observed in plot or individual tree-ring records. Specifically, most models were oversensitive to precipitation relative to tree rings and had predominantly negative or no response to temperature, whereas both tree-ring NPP and raw ring widths showed consistently positive responses (Fig. 2). The strongly negative effect of temperature coupled with a strongly positive effect of precipitation in the models suggests that the ecosystem models in our ensemble may be oversensitive to both temperature- and precipitation-driven droughts despite rapid postdrought recovery seen in many ecosystem models (Anderegg *et al.*, 2015). Although the tree rings in our study show positive correlations with temperature in both raw ring width and *a priori* detrended ring-width index, other studies provide support for negative effects of high temperatures on tree growth in the Eastern United States (e.g., Rollinson *et al.*, 2016). Nonetheless, models appear to be lacking moderating feedbacks that can cause positive temperature effects at low temperatures or reduce precipitation sensitivity. Indeed, the model in our ensemble that most closely matches the empirical relationship between temperature and NPP is LINKAGES, which calculates NPP directly from growing degree days rather than from separate GPP and autotrophic respiration functions (Post & Pastor, 1996).

Models were also oversensitive to precipitation relative to tree rings. Traditional tree-ring analyses have focused on growth responses to drought indices such as the Palmer Drought Severity Index that combines temperature, precipitation, and soil moisture into a single drought metric (e.g., Speer *et al.*, 2009; McEwan *et al.*, 2011). However, process-based ecosystem models drive productivity from independent forcing of temperature and precipitation, and temperature- vs. precipitation-based droughts may impact ecosystems differently (Anderegg *et al.*, 2013). Furthermore, models can vary in their representation of soil and hydrology so that the same temperature and precipitation drivers may result in different droughts both among models and compared to empirical system (Cook *et al.*, 2015). Tree-ring studies that have used mixed-

modeling frameworks to look at the simultaneous influences of temperature and precipitation on growth have revealed stronger influences of temperature on growth than precipitation (Rollinson *et al.*, 2016). To improve the ecosystem-scale model responses to both temperature and precipitation, further empirical research is needed to separate temperature and precipitation effects on tree growth.

Model responses to precipitation were more variable than temperature or CO₂, and consequently, there was little consensus in our model ensemble about when and where ecosystem productivity was driven by temperature vs. precipitation (Figs 2 and 3). Despite sensitivity disparities with tree-ring data, the strong control of precipitation in many models at the multicentennial scale supports hypotheses that drought may have been instrumental in driving ecosystem dynamics and species shifts in New England 500–600 years before present (Clifford & Booth, 2015). However, the relative insensitivity of productivity to temperature is difficult to reconcile with apparent temperature-driven millennial-scale compositional shifts in the paleoecological record (Prentice *et al.*, 1991; Blois *et al.*, 2013). In the model ensemble, fire and changes in biomass variability reduce model sensitivity to temperature rather than changes in relative composition, which suggests this lack of temperature sensitivity is not due to stabilizing shifts in plant functional types. Challenges of recreating spatial and temporal patterns of observed composition are a pervasive problem in ecosystem modeling and are known to have cascading impacts that bias other aspects of ecosystem dynamics including NPP and transpiration (Matthes *et al.*, 2016). Improved plant functional type parameterization within and across models through careful data collection will undoubtedly help address some of the discrepancies in magnitude and relative importance of temperature and precipitation seen between models and data in our study.

Differences in model responses to increasing CO₂ drive model ensemble uncertainty in NPP since 1900 (Fig. 3). The discrepancy of CO₂ enhancement effects in the model ensemble reflects the long-running debate in empirical studies about whether or not increased CO₂ has sustained effects on tree growth and ecosystem productivity. One synthesis from free-air CO₂ enrichment (FACE) studies, where conditions were raised to ~170 ppm above modern, increased productivity by 23%, or 13% ppm⁻¹ (Norby *et al.*, 2005). This is slightly less than half of the 43% increase in NPP attributed to CO₂ in our model ensemble over a similar increase in CO₂ over the past 100 years (Fig. 3). However, a 30-year study that found 12% increased stem width from increased CO₂ attributed most of this difference to early growth increases that diminished as the trees matured

(Hättenschwiler *et al.*, 1997). The effect of increasing competition for resources such as nitrogen and light during forest maturation has been used to explain a lack of CO₂ enhancement on tree growth in older forests quantified through tree rings (van der Sleen *et al.*, 2014; Fernández-de-Uña *et al.*, 2016). In tree rings, rising CO₂ has been associated with increased water use efficiency, but there has been little evidence that this translates into increased stem growth (Gedalof & Berg, 2010; Andreu-Hayles *et al.*, 2011; Peñuelas *et al.*, 2011; van der Sleen *et al.*, 2014; Frank *et al.*, 2015).

Many of the explanations for a lack of CO₂-induced growth increases in tree-ring widths center around the challenges of separating out the effects of increasing CO₂ from increasing tree size and stand-level forest dynamics (Jacoby & D'Arrigo, 1997; Andreu-Hayles *et al.*, 2011). In our results, stand-level NPP from tree rings showed no CO₂ effect, but individual ring widths displayed a negative effect that arises from the aforementioned challenges even though we explicitly accounted for increasing individual size in our analyses (Fig. 2 and Appendix S5). This model- and data-based uncertainty in the long-term effect of increasing CO₂ on ecosystems is the greatest source of uncertainty in current and future forest carbon cycle modeling (Sitch *et al.*, 2008; IPCC, 2013). Both fire and changes in biomass are correlated with CO₂ sensitivity in our ensemble of models across temporal scales (Table 3), indicating that disturbance and recovery processes may have strong influence on how CO₂ impacts ecosystem NPP. Consequently, accurate representation of causes of forest disturbances and process of recovery at local scales is necessary to reconcile models with empirical data and to make improved predictions of future forest dynamics under continued global change (Seidl *et al.*, 2011; Dietze *et al.*, 2014; Steinkamp & Hickler, 2015).

Models and data display conflicting responses of NPP to climate and CO₂, and these differences are more pronounced at the multicentennial timescales of past climate-driven ecosystem change. Models are generally oversensitive to growing season precipitation relative to tree-ring datasets and display negative responses to temperature, while tree rings show consistent positive effects. Further research on independent temperature and precipitation effects is necessary to diagnose systematic weaknesses in ecosystem models. Similarly, differences in model responses to CO₂ drive divergence of model ensemble NPP over the past century as the effects compound through time, while CO₂ concentration consistently increases. Despite empirical evidence for positive CO₂ effects on individual processes included in ecosystem models, the net effects on tree growth and forest-level NPP remain unclear and poorly constrained. Differences in disturbance as measured

through the presence of fire in model simulations only impact the sensitivity of productivity to climate and CO₂ at multicentennial timescales that capture multiple tree generations. The paths toward model improvement and reconciling discrepancies with data will be model-specific as each model has unique structural and parameterization requirements. However, accurate representation of fire and other disturbance processes and ecosystem recovery in models will require improved synthesis of short-term ecophysiological processes with gradual shifts in forest composition and structure seen in historical and paleoecological records. We advocate that this process of refinement includes consideration and formal evaluation of emergent, ecosystem-level behavior at the multidecadal and multicentennial temporal scales at which changes in forest composition and biomass occur. This approach would better align model evaluation with the organizational and temporal scales of forest dynamics and policy decisions.

Acknowledgements

This work was supported by NSF Macrosystems grant DEB #1241891 (Dietze, Matthes, Rollinson), 1241851 (Moore, Liu, Quaife, Schaefer, Hickler, Steinkamp), 1241930 (Pederson, Hessler, Dye, Poulter), and 1241874 (McLachlan, Raiho). Model and data contributions are as follows (in model alphabetical order, * indicates contribution sufficient to warrant authorship): CLM – Liu*, Moore*, F. Montane, A. Arellano, D. Ricciuto; ED, ED-LU – Rollinson*, Matthes*, and Dietze*; JULES, JULES-TRIFFID – Quaife*; LINKAGES – Raiho*, Dietze*; LPJ-GUESS – Steinkamp*, Hickler*, M. Forrest, C. Werner; LPJ-WSL – Poulter*; SiBCASA – Schaefer*, E. Jafarov; and Tree-Ring Data – Dye*, Bishop*, Pederson*, Hessler*. Analyses and writing were performed by Rollinson, Rollinson, Dietze, Liu, McLachlan, Moore, and Raiho contributed to the conceptual framework of the manuscript. All authors contributed to manuscript preparation. All codes used for analyses and figures in this manuscript are publically available on Github: <https://github.com/PaleON-Project/Temporal-Scaling-MS>.

References

- Anderegg LDL, Anderegg WRL, Berry JA (2013) Not all droughts are created equal: translating meteorological drought into woody plant mortality. *Tree Physiology*, **33**, 701–712.
- Anderegg WRL, Schwalm C, Biondi F *et al.* (2015) Pervasive drought legacies in forest ecosystems and their implications for carbon cycle models. *Science*, **349**, 528–532.
- Andreu-Hayles L, Planells O, Gutiérrez E, Muntan E, Helle G, Anchukaitis KJ, Schleser GH (2011) Long tree-ring chronologies reveal 20th century increases in water-use efficiency but no enhancement of tree growth at five Iberian pine forests. *Global Change Biology*, **17**, 2095–2112.
- Babst F, Poulter B, Trouet V *et al.* (2013) Site- and species-specific responses of forest growth to climate across the European continent. *Global Ecology and Biogeography*, **22**, 706–717.
- Best MJ, Pryor M, Clark DB *et al.* (2011) The Joint UK Land Environment Simulator (JULES), model description. Part 1: energy and water fluxes. *Geoscientific Model Development*, **4**, 677–699.
- Blois JL, Williams JW, Fitzpatrick MC *et al.* (2013) Modeling the climatic drivers of spatial patterns in vegetation composition since the Last Glacial Maximum. *Ecography*, **36**, 460–473.
- Boisvenue C, Running SW (2006) Impacts of climate change on natural forest productivity – evidence since the middle of the 20th century. *Global Change Biology*, **12**, 862–882.
- Burns RM, Honkala BH (1990) *Silvics of North America: 1. Conifers; 2. Hardwoods*. Agriculture Handbook 654. Vol 2. U.S. Department of Agriculture, Forest Service, Washington, DC.
- Charney ND, Babst F, Poulter B *et al.* (2016) Observed forest sensitivity to climate implies large changes in 21st century North American forest growth. *Ecology Letters*, **19**, 1119–1128.
- Clark DA, Brown S, Kicklighter DW, Chambers JQ, Thomlinson JR, Ni J (2001) Measuring net primary production in forest concepts and field methods. *Ecological Applications*, **11**, 356–370.
- Clark DB, Mercado LM, Sitch S *et al.* (2011) The Joint UK Land Environment Simulator (JULES), model description. Part 2: carbon fluxes and vegetation dynamics. *Geoscientific Model Development*, **4**, 701–722.
- Clifford MJ, Booth RK (2015) Late-Holocene drought and fire drove a widespread change in forest community composition in eastern North America. *The Holocene*, **25**, 1102–1110.
- Cook BI, Ault TR, Smerdon JE (2015) Unprecedented 21st century drought risk in the American Southwest and Central Plains. *Science Advances*, **1**, 1–7.
- Crucifix M, Harrison S, Brierley C (2012) Recent and deep pasts in paleoclimate model intercomparison project. *Eos, Transactions American Geophysical Union*, **93**, 539–539.
- Davis MB, Shaw RG (2001) Range shifts and adaptive responses to Quaternary climate change. *Science (New York, N.Y.)*, **292**, 673–679.
- Davis SC, Hessler AE, Scott CJ, Adams MB, Thomas RB (2009) Forest carbon sequestration changes in response to timber harvest. *Forest Ecology and Management*, **258**, 2101–2109.
- De Kauwe MG, Medlyn BE, Zaehle S *et al.* (2013) Forest water use and water use efficiency at elevated CO₂: a model-data intercomparison at two contrasting temperate forest FACE sites. *Global Change Biology*, **19**, 1759–1779.
- De Kauwe MG, Medlyn BE, Zaehle S *et al.* (2014) Where does the carbon go? A model-data intercomparison of vegetation carbon allocation and turnover processes at two temperate forest free-air CO₂ enrichment sites. *New Phytologist*, **203**, 883–899.
- DeLucia EH, Moore DJ, Norby RJ (2005) Contrasting responses of forest ecosystems to rising atmospheric CO₂: implications for the global C cycle. *Global Biogeochemical Cycles*, **19**, 1–9.
- Dietze M, Serbin S, Davidson C *et al.* (2014) A quantitative assessment of a terrestrial biosphere model's data needs across North American biomes. *Journal of Geophysical Research: Biogeosciences*, **119**, 286–300.
- D'Orangeville L, Duchesne L, Houle D, Kneeshaw D, Côté B, Pederson N (2016) Northeastern North America as a potential refugium for boreal forests in a warming climate. *Science*, **352**, 1452–1455.
- Doughty CE, Malhi Y, Araujo-Murakami A *et al.* (2014) Allocation trade-offs dominate the response of tropical forest growth to seasonal and interannual drought. *Ecology*, **95**, 2192–2201.
- Dye A, Barker-Plotkin A, Bishop D, Pederson N, Poulter B, Hessler A (2016) Comparing tree-ring and permanent plot estimates of aboveground net primary production in three Eastern U.S. forests. *Ecosphere*, **7**, e01454.
- Eisen K, Plotkin AB (2015) Forty years of forest measurements support steadily increasing aboveground biomass in a maturing, Quercus-dominant northeastern forest. *Journal of the Torrey Botanical Society*, **142**, 97–112.
- Fernández-de-Uña L, McDowell NG, Cañellas I, Gea-Izquierdo G (2016) Disentangling the effect of competition, CO₂ and climate on intrinsic water-use efficiency and tree growth. *Journal of Ecology*, **104**, 678–690.
- Fischelli NA, Frelich LE, Reich PB (2013) Temperate tree expansion into adjacent boreal forest patches facilitated by warmer temperatures. *Ecography*, **36**, 001–010.
- Foster DR, Motzkin G, Slater B (1998) Land-use history as long-term broad-scale disturbance: regional forest dynamics in central New England. *Ecosystems*, **1**, 96–119.
- Foster JR, D'Amato AW, Bradford JB (2014) Looking for age-related growth decline in natural forests: unexpected biomass patterns from tree rings and simulated mortality. *Oecologia*, **175**, 363–374.
- Frank DC, Poulter B, Saurer M *et al.* (2015) Water-use efficiency and transpiration across European forests during the Anthropocene. *Nature Climate Change*, **5**, 579–583.
- Friedlingstein P, Meinshausen M, Arora VK, Jones CD, Anav A, Liddicoat SK, Knutti R (2014) Uncertainties in CMIP5 climate projections due to carbon cycle feedbacks. *Journal of Climate*, **27**, 511–526.
- Gedalof Z, Berg AA (2010) Tree ring evidence for limited direct CO₂ fertilization of forests over the 20th century. *Global Biogeochemical Cycles*, **24**, 2–7.

- Gerten D, Schaphoff S, Haberlandt U, Lucht W, Sitch S (2004) Terrestrial vegetation and water balance – hydrological evaluation of a dynamic global vegetation model. *Journal of Hydrology*, **286**, 249–270.
- Gómez-Aparicio L, García-Valdés R, Ruiz-Benito P, Zavala MA (2011) Disentangling the relative importance of climate, size and competition on tree growth in Iberian forests: implications for forest management under global change. *Global Change Biology*, **17**, 2400–2414.
- Goulden ML, Mcmillan AMS, Winston GC, Rocha AV, Manies KL, Harden JW, Bond-Lamberty BP (2011) Patterns of NPP, GPP, respiration, and NEP during boreal forest succession. *Global Change Biology*, **17**, 855–871.
- Graumlich LJ, Brubaker LB, Grier CC (1989) Long-term trends in forest net primary productivity: Cascade Mountains, Washington. *Ecology*, **70**, 405–410.
- Grissino-Mayer HD, Fritts HC (1997) The International Tree-Ring Data Bank: an enhanced global database serving the global scientific community. *The Holocene*, **7**, 235–238.
- Hättenschwiler S, Miglietta F, Raschi A, Körner C (1997) Thirty years of in situ tree growth under elevated CO₂: a model for future forest responses? *Global Change Biology*, **3**, 463–471.
- Hempel S, Frieler K, Warszawski L, Schewe J, Piontek F (2013) A trend-preserving bias correction – the ISI-MIP approach. *Earth System Dynamics*, **4**, 219–236.
- Hollister RD, Webber PJ, Tweedie CE (2005) The response of Alaskan arctic tundra to experimental warming: differences between short- and long-term responses. *Global Change Biology*, **11**, 525–536.
- IPCC (2013) Summary for policy makers. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM), pp. 3–29. Cambridge University Press, Cambridge, UK and New York, NY, USA.
- Jacoby GC, D'Arrigo RD (1997) Tree rings, carbon dioxide, and climatic change. *Proceedings of the National Academy of Sciences of the United States of America*, **94**, 8350–8353.
- Jenkins JC, Birdsey RA, Pan Y (2001) Biomass and NPP estimation for the mid-Atlantic region (USA) using plot-level forest inventory data. *Ecological Applications*, **11**, 1174–1193.
- Keeling RF, Piper SC, Bollenbacher AF, Walker JS (2009) Atmospheric CO₂ records from sites in the SIO air sampling network. In: *Trends: A Compendium of Data on Global Change*. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, TN, USA.
- Kumar J, Brooks B-GJ, Thornton PE, Dietze MC (2012) Sub-daily statistical downscaling of meteorological variables using neural networks. *Procedia Computer Science*, **9**, 887–896.
- Litton CM, Raich JW, Ryan MG (2007) Carbon allocation in forest ecosystems. *Global Change Biology*, **13**, 2089–2109.
- Luyssaert S, Inglima I, Jung M *et al.* (2007) CO₂ balance of boreal, temperate, and tropical forests derived from a global database. *Global Change Biology*, **13**, 2509–2537.
- Matesanz S, Brooker RW, Valladares F, Klotz S (2009) Temporal dynamics of marginal steppe vegetation over a 26-year period of substantial environmental change. *Journal of Vegetation Science*, **20**, 299–310.
- Matthes JH, Goring S, Williams JW, Dietze MC (2016) Benchmarking historical CMIP5 land-climate feedbacks across the Upper Midwest and Northeastern United States. *Journal of Geophysical Research: Biogeosciences*, **121**, 523–535.
- McEwan RW, Dyer JM, Pederson N (2011) Multiple interacting ecosystem drivers: toward an encompassing hypothesis of oak forest dynamics across eastern North America. *Ecography*, **34**, 244–256.
- Medvigy D, Wofsy SC, Munger JW, Hollinger DY, Moorcroft PR (2009) Mechanistic scaling of ecosystem function and dynamics in space and time: Ecosystem Demography model version 2. *Journal of Geophysical Research*, **114**, 1–21.
- Melillo JM, Steudler PA, Aber JD *et al.* (2002) Soil warming feedbacks the carbon-cycle climate. *Science*, **298**, 2173–2176.
- Morin X, Lechowicz MJ, Augspurger C, O'Keefe J, Viner D, Chuine I (2009) Leaf phenology in 22 North American tree species during the 21st century. *Global Change Biology*, **15**, 961–975.
- Norby RJ, Delucia EH, Gielen B *et al.* (2005) Forest response to elevated CO₂ is conserved across a broad range of productivity. *Proceedings of the National Academy of Sciences of the United States of America*, **102**, 18052–18056.
- Norby RJ, Warren JM, Iversen CM, Medlyn BE, McMurtrie RE (2010) CO₂ enhancement of forest productivity constrained by limited nitrogen availability. *Proceedings of the National Academy of Sciences of the United States of America*, **107**, 19368–19373.
- Oleson KW, Lawrence DM, Gordon B *et al.* (2010) Technical description of version 4.0 of the Community Land Model (CLM). NCAR/TN-478 + STR NCAR Technical Note, 266.
- Peñuelas J, Canadell JG, Ogaya R (2011) Increased water-use efficiency during the 20th century did not translate into enhanced tree growth. *Global Ecology and Biogeography*, **20**, 597–608.
- Peñuelas J, Sardans J, Estiarte M *et al.* (2013) Evidence of current impact of climate change on life: a walk from genes to the biosphere. *Global Change Biology*, **19**, 2303–2338.
- Piao S, Sitch S, Ciais P *et al.* (2013) Evaluation of terrestrial carbon cycle models for their response to climate variability and to CO₂ trends. *Global Change Biology*, **19**, 2117–2132.
- Post WM, Pastor J (1996) Linkages – an individual-based forest ecosystem model. *Climatic Change*, **34**, 253–261.
- Prentice IC, Bartlein PJ, Webb T (1991) Vegetation and climate change in eastern North America since the last glacial maximum. *Ecology*, **72**, 2038–2056.
- Ramankutty N, Foley JA (1999) Estimating historical changes in global land cover: croplands from 1700 to 1992. *Global Biogeochemical Cycles*, **13**, 997–1027.
- Renwick KM, Rocca ME (2014) Temporal context affects the observed rate of climate-driven range shifts in tree species. *Global Ecology and Biogeography*, **24**, 44–51.
- Rollinson CR, Kaye MW, Canham CD (2016) Interspecific variation in growth responses to climate and competition of five eastern tree species. *Ecology*, **97**, 1003–1011.
- Schaefer K, Collatz GJ, Tans P *et al.* (2008) Combined simple biosphere/Carnegie-Ames-Stanford approach terrestrial carbon cycle model. *Journal of Geophysical Research*, **113**, G03034.
- Seidl R, Fernandes PM, Fonseca TF *et al.* (2011) Modelling natural disturbances in forest ecosystems: a review. *Ecological Modelling*, **222**, 903–924.
- Shuman B, Webb T, Bartlein P, Williams JW (2002) The anatomy of a climatic oscillation: vegetation change in eastern North America during the Younger Dryas chronozone. *Quaternary Science Reviews*, **21**, 1777–1791.
- Sitch S, Smith B, Prentice IC *et al.* (2003) Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model. *Global Change Biology*, **9**, 161–185.
- Sitch S, Huntingford C, Gedney N *et al.* (2008) Evaluation of the terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using five Dynamic Global Vegetation Models (DGVMs). *Global Change Biology*, **14**, 2015–2039.
- van der Sleen P, Groenendijk P, Vlam M *et al.* (2014) No growth stimulation of tropical trees by 150 years of CO₂ fertilization but water-use efficiency increased. *Nature Geoscience*, **8**, 24–28.
- Smith B, Warland D, Arneith A, Hickler T, Leadley P, Siltberg J, Zaehle S (2014) Implications of incorporating N cycling and N limitations on primary production in an individual-based dynamic vegetation model. *Biogeosciences*, **11**, 2027–2054.
- Speer JH, Grissino-Mayer HD, Orvis KH, Greenberg CH (2009) Climate response of five oak species in the eastern deciduous forest of the southern Appalachian Mountains, USA. *Canadian Journal of Forest Research*, **39**, 507–518.
- Steinkamp J, Hickler T (2015) Is drought-induced forest dieback globally increasing? *Journal of Ecology*, **103**, 31–43.
- Taylor KE, Stouffer RJ, Meehl GA (2012) An overview of CMIP5 and the experiment design. *Bulletin of the American Meteorological Society*, **93**, 485–498.
- Turner MG, Collins SL, Lugo AL, Magnuson JJ, Rupp TS, Swanson FJ (2003) Disturbance dynamics and ecological response: the contribution of long-term ecological research. *BioScience*, **53**, 46–56.
- Turner DP, Ritts WD, Cohen WB *et al.* (2005) Site-level evaluation of satellite-based global terrestrial gross primary production and net primary production monitoring. *Global Change Biology*, **11**, 666–684.
- Walker AP, Hanson PJ, De Kauwe MG *et al.* (2014) Comprehensive ecosystem model-data synthesis using multiple data sets at two temperate forest free-air CO₂ enrichment experiments: model performance at ambient CO₂ concentration. *Journal of Geophysical Research: Biogeosciences*, **119**, 937–964.
- Walker AP, Zaehle S, Medlyn BE *et al.* (2015) Predicting long-term carbon sequestration in response to CO₂ enrichment: how and why do current ecosystem models differ? *Global Biogeochemical Cycles*, **29**, 476–495.
- Wei Y, Liu S, Huntzinger DN *et al.* (2014) NACP MsTMIP: Global and North American Driver Data for Multi-Model Intercomparison. ORNL DAAC, Oak Ridge, TN, USA. Available at: https://daac.ornl.gov/cgi-bin/dsviewer.pl?ds_id=1220 (accessed 23 September 2013).
- Williams AP, Allen CD, Macalady AK *et al.* (2013) Temperature as a potent driver of regional forest drought stress and tree mortality. *Nature Climate Change*, **3**, 292–297.
- Wood S (2012) mgcv: GAMs with GCV/AIC/REML smoothness estimation and GAMMs by PQL Version 1.8-16. Available at: <https://cran.r-project.org/web/packages/mgcv/> (accessed 8 November 2016).

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Appendix S1 Growing season temperature and precipitation time series for 850–2010 A.D. for all six sites that were used as model drivers and in for sensitivity analysis.

Appendix S2 Model simulation protocol.

Appendix S3 Location and record information for tree-ring width information from the International Tree Ring Databank (Grisino-Mayer & Fritts, 1997).

Appendix S4 Mean and 95% confidence interval of spine-detrended ring width index (RWI) for all tree ring records used in analyses.

Appendix S5 Sensitivity of relativized NPP and tree-ring width index (RWI) to growing season temperature, growing season precipitation, and CO₂ across three temporal scales.

Appendix S6 Sensitivity of relativized NPP and tree-ring width to growing season temperature, growing season precipitation, and CO₂ across three temporal scales restricted to Harvard and Howland sites.

Appendix S7 Relativized sensitivity of model NPP or ring width to growing season (May–September) temperature, growing season precipitation, and CO₂ by temporal extent.

Appendix S8 Model simulation protocol PaleON Model-Data Inter-comparison project Phase 1a: site-level comparison.