

A KERNEL-CERTIFIED CORE FOR THE STUDY OF AGRAWAL'S CONJECTURE AT $r = 5$

THE UNICO/NOUS PIPELINE

ABSTRACT. This note collects the part of our structural study of Agrawal's conjecture at $r = 5$ that is mechanically verified. Every statement called a theorem below is proved in Lean 4 over Mathlib, with no `sorry`, no `admit` and no added axioms; the corresponding declaration name is given with each statement. Every computational claim carries a replayable certificate with its detection criterion embedded and SHA-256 manifests. We then pose, without any claim of proof, the two open problems that our working material identifies as central. Nothing in this note asserts the conjecture.

1. SETTING

Agrawal's conjecture (arising from the AKS primality test [1]; see also [2, 3, 4]) states: if r is prime, $r \nmid n$, and $(X - 1)^n \equiv X^n - 1 \pmod{n, X^r - 1}$, then n is prime or $n^2 \equiv 1 \pmod{r}$. Our study concerns $r = 5$. This note isolates its mechanically certified core; the statements are elementary, and their interest lies in the role they play in the wider study and in the fact that a proof assistant has checked every one of them.

Throughout, p is an odd prime. Let

$$R_p = (\mathbf{Z}/p)[X] / (X^2 - X - 1), \quad \varepsilon = \text{the class of } X,$$

so that $\varepsilon^2 = \varepsilon + 1$, and set $\sqrt{5} := 2\varepsilon - 1$, which satisfies $(\sqrt{5})^2 = 5$ in R_p (`GoldenRing`, `eps_sq`, `sqr5_sq`). Let F_n, L_n denote the Fibonacci and Lucas numbers, with $L_n = F_{n+1} + F_{n-1}$ (`lucas`). Define

$$A_n = \begin{cases} L_n, & n \text{ even,} \\ F_n, & n \text{ odd,} \end{cases} \quad H_n = \gcd(A_n, 5^{n-1} + 1),$$

and, dually, $B_n = F_n$ (n even), L_n (n odd), with $J_n = \gcd(B_n, 5^{n-1} - 1)$.

2. THE GOLDEN FROBENIUS AND THE INERTIA OF J_n

Theorem 2.1 (`golden_frobenius`). *Assume 5 is not a square in $\mathbf{Z}/p$ and $p \neq 2$. Then $\varepsilon^p = 1 - \varepsilon$ in R_p .*

An autonomous multi-model pipeline (Claude Fable 5, Claude Opus 4.8, GPT-5.6-Sol, with an independent external review pass), orchestrated by Daniele Cappello, who carried the messages, audited the artifacts and made no mathematical contribution. Repository with the Lean sources and all computational certificates: github.com/Solarys431/agrawal-r5.

Certified proof sketch. By Euler's criterion $5^{(p-1)/2} = -1$ in $\mathbf{Z}/p$ (`euler_nonsquare`), hence $(\sqrt{5})^p = (5)^{(p-1)/2}\sqrt{5} = -\sqrt{5}$ (`sqrt5_pow_p`). Expanding $(2\varepsilon - 1)^p$ by the Frobenius endomorphism and cancelling the unit 2 gives the claim. $\square$

Corollary 2.2 (`golden_pow_p_succ`). *Under the same hypotheses, $\varepsilon^{p+1} = -1$ in R_p .*

Theorem 2.3 (`inertia_J`). *Assume 5 is not a square in $\mathbf{Z}/p$ and $p \neq 2$. Then for no $n \geq 1$ do $\varepsilon^{2n} = 1$ in R_p and $5^{n-1} = 1$ in $\mathbf{Z}/p$ hold simultaneously.*

Certified proof sketch. If n is odd, raising $\varepsilon^{2n} = 1$ to the power $(p+1)/2$ and comparing with $(\varepsilon^{p+1})^n = (-1)^n = -1$ gives $1 = -1$, absurd for $p \neq 2$. If n is even, $5 = 5^n = (5^{n/2})^2$ exhibits 5 as a square, against the hypothesis. $\square$

Via the identity $\varepsilon^{n+1} = F_{n+1}\varepsilon + F_n$ in R_p (`golden_pow_fib`, `golden_pow_pred`) one obtains the divisibility forms, in which the hypothesis is purely arithmetic by quadratic reciprocity (`not_isSquare_five`):

Theorem 2.4 (`inertia_J_fib_mod5`, `inertia_J_lucas_mod5`). *Let $p \equiv \pm 2 \pmod{5}$, $p \neq 2$. For even $n \geq 2$, p does not divide $\gcd(F_n, 5^{n-1} - 1)$; for odd $n \geq 1$, p does not divide $\gcd(L_n, 5^{n-1} - 1)$. Equivalently, no prime inert in $\mathbf{Q}(\sqrt{5})$ divides any J_n .*

3. THE CANONICAL SUPPORT WITNESS

Theorem 3.1 (`support_witness`). *Let $p \equiv 2 \pmod{5}$, $p \neq 2$, and put $n_0 = (p+1)/2$. Then $n_0 \equiv 4 \pmod{5}$, $p \mid 5^{n_0-1} + 1$, and p divides L_{n_0} if n_0 is even, F_{n_0} if n_0 is odd. In particular $p \mid H_{n_0}$, with an index in the class 4 mod 5.*

4. THE GOLDEN-CYCLOTOMIC ENTANGLEMENT

Theorem 4.1 (`entanglement`). *In any commutative ring, if $\zeta^4 + \zeta^3 + \zeta^2 + \zeta + 1 = 0$, then*

$$\zeta^3(1 + \zeta + \zeta^4)^2(\zeta - 1)^4 = 5.$$

In particular this holds in $\mathbf{Z}[\zeta_5]$ with $\varepsilon = 1 + \zeta + \zeta^4$ the image of the golden unit.

The proof is a single `linear_combination` with an explicit cofactor; $\zeta^5 = 1$ follows from the same hypothesis (`zeta_pow_five`).

5. THE FERMAT-SHADOW GLUE

Lemma 5.1 (`mul_dvd_gcd_mul`). *If $x \mid M$ and $y \mid M$, then $xy \mid \gcd(x, y) M$.*

Theorem 5.2 (`fermat_shadow`). *Let n be squarefree with $5 \nmid n$, and suppose that for every prime $p \mid n$*

$$(*) \quad 5^{n/p-1} \equiv 1 \pmod{p}.$$

Then $n \mid 5^{n-1} - 1$; that is, n is a Fermat pseudoprime to base 5.

Remark 5.3. Hypothesis (*) is exactly the local norm constraint that our working study derives for squarefree counterexamples to Agrawal’s congruence at $r = 5$. That derivation is *not* part of this certified core: here (*) is an explicit hypothesis, and the theorem certifies the arithmetic glue $n - 1 = p(n/p - 1) + (p - 1)$ together with the squarefree gluing.

6. COMPUTATIONAL CERTIFICATES

Independently of the kernel-checked statements above, the repository carries replayable certificates, each embedding its own detection criterion and reconstruction data, with SHA-256 manifests:

- (1) seven *certified-empty fibres* of a three-factor sieve associated with the study (pairs (13, 37), (17, 113), (13, 277), (23, 167), (7, 823), (83, 347), (463, 1327)): for each pair, the finitely many levels required by the certificate are integrally factored, all prime factors are proven prime, and no prime in the certificate’s detector classes survives;
- (2) a self-certifying census of H_n for all $n \equiv 4 \pmod{5}$, $n \leq 10^5$: 9,725 nontrivial values, all factorizations proven, and *no* prime factor $\equiv \pm 1 \pmod{5}$ occurs;
- (3) SHA-256 sums for a prime-by-prime corpus up to 10^9 used by the working study.

These are certificates of finite computations; they prove exactly what they state and nothing beyond.

7. TWO OPEN PROBLEMS

We pose the two statements that our working material identifies as the central open problems of the $r = 5$ study. We claim no proof and no partial proof here.

Conjecture 7.1 (golden inertia). *For every $n \equiv 4 \pmod{5}$, every odd prime factor of H_n distinct from 5 is $\equiv \pm 2 \pmod{5}$.*

Theorem 3.1 shows that every prime $p \equiv 2 \pmod{5}$ does occur in some H_n with $n \equiv 4 \pmod{5}$; the census in item (2) above verifies Conjecture 7.1 for all $n \leq 10^5$.

Conjecture 7.2 (fibre emptiness). *The pattern of item (1) above is general: for every admissible pair, the corresponding fibre is empty.*

8. FORMALIZATION DATA

Eight modules, 646 source lines, 45 named declarations; no `sorry`, no `admit`, no axioms beyond Mathlib’s. Mathlib is pinned to a fixed upstream commit; a clean clone builds with `lake exe cache get && lake build`. The table mapping each statement of this note to its declaration and file is in the repository `README`.

REFERENCES

- [1] M. Agrawal, N. Kayal, and N. Saxena, *PRIMES is in P*, Ann. of Math. 160 (2004), 781–793.
- [2] H. W. Lenstra, Jr. and C. Pomerance, *Remarks on Agrawal’s conjecture*, AIM note, 2003.
- [3] H. W. Lenstra, Jr. and C. Pomerance, *Primality testing with Gaussian periods*, J. Eur. Math. Soc. 21 (2019), 1229–1269.
- [4] T. Váša, *Agrawal’s conjecture and Carmichael numbers*, Student Scientific Conference, Comenius University Bratislava, 2009.