# M21 Hyoscine Butyl bromide

| | |
|---|---|
| **Scope of Practice** | P |
| **Classification** | • Anticholinergic (antimuscarinic) |
| **Usual Presentation** | • 20mg/1ml ampule |
| **Mechanism of Action** | • Hyoscine butyl bromide blocks muscarinic acetylcholine receptors (primarily M3) on smooth muscle, reducing parasympathetic-mediated contractions |
| **Indications** | • Lower abdominal cramping associated with gastroenteritis, rental colic, irritable bowel disease and dysmenorrhea. |
| **Contraindications** | • Tachyarrhythmias<br>• Narrow angle glaucoma<br>• Stenosis of the GI or unitary tract<br>• Myasthenia gravis<br>• Hypertrophy of the prostate<br>• Renal Impairment |
| **Precautions** | • Nil |
| **Side Effects** | • Tachycardia<br>• Bradycardia<br>• Hypotension<br>• Urinary urgency<br>• Dry mouth<br>• Constipation<br>• Drowsiness |
| **Adult Dose & Administration** | • 20mg IV/IM/SC |
| **Pediatric Dose & Administration** | • Not for pediatric administration |
| **Use in Pregnancy and Lactation** | • Pregnancy: Safety has not been established. Administration in pregnancy is not advised.<br>• Lactation: Safety has not been established. Administration in lactation is not advised. |
| **Additional Information** | • Nil |