M27 Midazolam

Scope of Practice	EP
Classification	• Benzodiazepine
Usual Presentation	• 5mg/1ml ampule
Mechanism of Action	• Potentiates the action of the GABA neurotransmitter producing sedation and seizure inhibition
Indications	• Seizures • Sedation of agitated or combative patients
Contraindications	• Known allergy
Precautions	• Respiratory distress • Shock and/or hypotension
Side Effects	• Reduced level of consciousness • Respiratory Depression • Hypotension
Adult Dose & Administration	**Seizures** • 5mg IV/IO/IM (repeat once after 10 minutes if required) **Sedation** • 1-5mg IV/IO/IM **Dilution for IV/IO administration** • In a 5ml syringe, draw up 4mls of normal saline • Add 1ml (5mg) of Midazolam. • Concentration is now 1mg/1ml
Pediatric Dose & Administration	**Seizures** • 0.1mg/kg **IV** (max 5mg – may repeat once after 10 minutes if required) • 0.2mg/kg **IM** (max 5mg – may repeat once after 10 minutes if required) **Dilution for IV administration** • In a 5 or 10mL syringe, draw up 4mls of normal saline • Add 1mL (5mg) of Midazolam. • Concentration is now 1mg/1ml
Use in Pregnancy and Lactation	• Pregnancy: Fetal harm has been demonstrated however, midazolam should be administered if clinically indicated. • Lactation: Safety has not been established however, midazolam should be administered if clinically indicated.
Additional Information	• Benzodiazepines may cause airway, breathing and circulation compromise. • If administered, all possible monitoring should be in place, including nasal end-tidal CO2 monitoring. • An IV (if not available in place) should be established as soon as possible after administration.