M33 Pethidine

Scope of Practice	EP
Classification	• Opiate Analgesic
Usual Presentation	• 50mg/1ml ampule
Mechanism of Action	• Binds with Mu opioid pain receptors creating analgesia
Indications	• Moderate to Severe Pain
Contraindications	• Known allergy • Respiratory depression • SBP <90mmHg • Renal Impairment • Concurrent use of Mono Amine Oxidase Inhibitors • Suspected bowel obstruction
Precautions	• Respiratory distress or difficulty • Elderly • Pediatrics age <1 year (increased risk or respiratory depression) • Morbid obesity • Moderate Hypotension / Shock
Side Effects	• Respiratory depression • Hypotension • Drowsiness / reduced level of consciousness • Arrhythmias
Adult Dose & Administration	• IM: 25-100mg (maximum 400mg/24 hours) • IV: 25-50mg. Administer slowly over 2 minutes. (maximum 200mg/24 hours) **Dilution for IV administration** • In a 10ml syringe, draw up 9mls of normal saline • Add 1ml (50mg) of Pethidine. • Concentration is 5mg/1ml (50mg in 10ml)
Pediatric Dose & Administration	• IM: 0.5-2mg/kg (repeat after 4 hours)
Use in Pregnancy and Lactation	• Pregnancy: May cause fetal harm. Pethidine should not be administered during pregnancy. • Lactation: Safety has not been established. Pethidine should only be administered if the benefit outweighs the risk. If administered, the mother should be advised to stop breast feeding until advised by their physician.
Additional Information	• Nil