M17 Glucose 40% Oral Paste

Scope of Practice	EMR
Classification	• Carbohydrate
Usual Presentation	• Plastic tube containing 15g glucose
Mechanism of Action	• Provides glucose for cellular metabolism, raising blood glucose levels rapidly in hypoglycemia.
Indications	• Hypoglycemia (RBS<70mg/dl)
Contraindications	• Known allergy • Unresponsive patients or patients unable to swallow
Precautions	• Airway occlusion in semi-conscious patients (particularly children <2years old)
Side Effects	• May cause vomiting
Adult Dose & Administration	• 1 tube PO – adjusting amount of paste inserted according to space available. • Repeat after 5-10 minutes (if required)
Pediatric Dose & Administration	• ½ tube PO – adjusting amount of paste inserted according to space available. • Repeat after 5-10 minutes (if required)
Use in Pregnancy and Lactation	• Pregnancy: Safety has not been established. However, glucose should be administered if clinically indicated. • Lactation: Safety has not been established. However, glucose should be administered if clinically indicated.
Additional Information	• Glucose paste is not absorbed via the buccal mucosa and must be swallowed to be effective. • In very young children/neonates, if the patient is able to suck, apply a small amount of glucose paste to the end of a gloved finger and allow the child to suck the paste.