Combivent

Scope of Practice	EMT
Classification	• ß2 Agonist and anticholinergic

Usual Presentation	• 2.5ml nebule containing:
	  • 2.5mg Salbutamol
	  • 0.5mg Ipratropium Bromide

Mechanism of Action	• Salbutamol agonizes ß2 adrenergic receptors causing bronchodilation • Ipratropium Bromide blocks the action of acetylcholine at muscarinic receptors, resulting in bronchodilation.
Indications	• Asthma • COPD
Contraindications	• Known allergy
Precautions	• Nil
Side Effects	• Tachycardia • Palpitations • Headache • Dry mouth
Adult Dose & Administration	**Asthma & COPD** • 2.5ml Nebulized (2.5mg Salbutamol, 0.5mg Ipratropium Bromide) • Dilute dose with normal saline or sterile water to a maximum of 5ml.
Pediatric Dose & Administration	• Not recommended for pediatrics
Use in Pregnancy and Lactation	• Pregnancy: Safety has not been established. Combivent should be administered if clinically indicated. • Lactation: may be administered in lactation however, the mother should be advised to stop breast feeding until advised by their physician.
Additional Information	• Combivent does not have a significant role in the management of bronchospasm associated with smoke inhalation or chest infections. However, if significant bronchospasm is present, Combivent may be administered.