


4

Carbon and the Molecular Diversity of Life

KEY CONCEPTS

- 4.1 Organic chemistry is the study of carbon compounds
- 4.2 Carbon atoms can form diverse molecules by bonding to four other atoms
- 4.3 A few chemical groups are key to molecular function



▲ Carbon can bond to four other atoms or groups of atoms, making a large variety of molecules possible.

▲ **Figure 4.1** What properties make carbon the basis of all life?

Carbon: The Backbone of Life

Living organisms, such as the plants and the Qinling golden snub-nosed monkeys shown in **Figure 4.1**, are made up of chemicals based mostly on the element carbon. Carbon enters the biosphere through the action of plants and other photosynthetic organisms. Plants use solar energy to transform atmospheric CO_2 into the molecules of life, which are then taken in by plant-eating animals.

Of all the chemical elements, carbon is unparalleled in its ability to form molecules that are large, complex, and varied, making possible the diversity of organisms that have evolved on Earth. Proteins, DNA, carbohydrates, and other molecules that distinguish living matter from inanimate material are all composed of carbon atoms bonded to one another and to atoms of other elements. Hydrogen (H), oxygen (O), nitrogen (N), sulfur (S), and phosphorus (P) are other common ingredients of these compounds, but it is the element carbon (C) that accounts for the enormous variety of biological molecules.

Large biological molecules, such as proteins, are the main focus of Chapter 5. In this chapter, we investigate the properties of smaller molecules. We will use these small molecules to illustrate concepts of molecular architecture that will help explain why carbon is so important to life, at the same time highlighting the theme that emergent properties arise from the organization of matter in living organisms.

Organic chemistry is the study of carbon compounds

For historical reasons, compounds containing carbon are said to be organic, and their study is called **organic chemistry**. By the early 1800s, chemists had learned to make simple compounds in the laboratory by combining elements under the right conditions. Artificial synthesis of the complex molecules extracted from living matter seemed impossible, however. Organic compounds were thought to arise only in living organisms, which were believed to contain a life force beyond the jurisdiction of physical and chemical laws.

Chemists began to chip away at this notion when they learned to synthesize organic compounds in the laboratory. In 1828, Friedrich Wöhler, a German chemist, tried to make an “inorganic” salt, ammonium cyanate, by mixing solutions of ammonium ions (NH_4^+) and cyanate ions (CNO^-). Wöhler was astonished to find that instead he had made urea, an organic compound present in the urine of animals.

The next few decades saw laboratory synthesis of increasingly complex organic compounds, supporting the view that physical and chemical laws govern the processes of life. Organic chemistry was redefined as the study of carbon compounds, regardless of origin. Organic compounds range from simple molecules, such as methane (CH_4), to colossal ones, such as proteins, with thousands of atoms.


Organic Molecules and the Origin of Life on Earth

EVOLUTION In 1953, Stanley Miller, a graduate student of Harold Urey’s at the University of Chicago, helped bring the abiotic (nonliving) synthesis of organic compounds into the context of evolution. Study **Figure 4.2** to learn about his classic experiment. From his results, Miller concluded that complex organic molecules could arise spontaneously under conditions thought at that time to have existed on the early Earth. You can work with the data from a related experiment in the **Scientific Skills Exercise**. These experiments support the idea that abiotic synthesis of organic compounds, perhaps near volcanoes, could have been an early stage in the origin of life (see Chapter 25).

The overall percentages of the major elements of life—C, H, O, N, S, and P—are quite uniform from one organism to another, reflecting the common evolutionary origin of all life. Because of carbon’s ability to form four bonds, however, this limited assortment of atomic building blocks can be used to build an inexhaustible variety of organic molecules. Different species of organisms, and different individuals within a species, are distinguished by variations in the types

Can organic molecules form under conditions estimated to simulate those on the early Earth?

Experiment In 1953, Stanley Miller set up a closed system to mimic conditions thought at that time to have existed on the early Earth. A flask of water simulated the primeval sea. The water was heated so that some vaporized and moved into a second, higher flask containing the “atmosphere”—a mixture of gases. Sparks were discharged in the synthetic atmosphere to mimic lightning.



Results Miller identified a variety of organic molecules that are common in organisms. These included simple compounds, such as formaldehyde (CH_2O) and hydrogen cyanide (HCN), and more complex molecules, such as amino acids and long chains of carbon and hydrogen known as hydrocarbons.

Conclusion Organic molecules, a first step in the origin of life, may have been synthesized abiotically on the early Earth. Although new evidence indicates that the early Earth’s atmosphere was different from the “atmosphere” used by Miller in this experiment, recent experiments using the revised list of chemicals also produced organic molecules. (We will explore this hypothesis in more detail in Chapter 25.)

Source: S. L. Miller, A production of amino acids under possible primitive Earth conditions, *Science* 117:528–529 (1953).

WHAT IF? If Miller had increased the concentration of NH_3 in his experiment, how might the relative amounts of the products HCN and CH_2O have differed?

SCIENTIFIC SKILLS EXERCISE

Working with Moles and Molar Ratios

Could the First Biological Molecules Have Formed Near Volcanoes on Early Earth? In 2007, Jeffrey Bada, a former graduate student of Stanley Miller's, discovered some vials of samples that had never been analyzed from an experiment performed by Miller in 1958. In this experiment, Miller used hydrogen sulfide gas (H_2S) as one of the gases in the reactant mixture. Since H_2S is released by volcanoes, the H_2S experiment was designed to mimic conditions near volcanoes on early Earth. In 2011, Bada and colleagues published the results of their analysis of these "lost" samples. In this exercise, you will make calculations using the molar ratios of reactants and products from the H_2S experiment.


How the Experiment Was Done According to his laboratory notebook, Miller used the same apparatus as in his original experiment (see Figure 4.2), but the mixture of gaseous reactants included methane (CH_4), carbon dioxide (CO_2), hydrogen sulfide (H_2S), and ammonia (NH_3). After three days of simulated volcanic activity, he collected samples of the liquid, partially purified the chemicals, and sealed the samples in sterile vials. In 2011, Bada's research team used modern analytical methods to analyze the products in the vials for the presence of amino acids, the building blocks of proteins.

Data from the Experiment The table below shows 4 of the 23 amino acids detected in the samples from Miller's 1958 H_2S experiment.

Product Compound	Molecular Formula	Molar Ratio (Relative to Glycine)
Glycine	$\text{C}_2\text{H}_5\text{NO}_2$	1.0
Serine	$\text{C}_3\text{H}_7\text{NO}_3$	3.0×10^{-2}
Methionine	$\text{C}_5\text{H}_{11}\text{NO}_2\text{S}$	1.8×10^{-3}
Alanine	$\text{C}_3\text{H}_7\text{NO}_2$	1.1

Interpret the Data

1. A *mole* is the number of grams of a substance that equals its molecular (or atomic) mass in daltons. There are 6.02×10^{23} molecules (or atoms) in 1.0 mole (Avogadro's number; see Concept 3.2). The data table shows the "molar ratios" of some of the products from the Miller H_2S experiment. In a molar ratio, each unitless value is expressed relative to a standard for that experiment. Here, the standard is the number of moles of the amino acid glycine, which is set to a value of 1.0. For instance, serine has a molar ratio of 3.0×10^{-2} , meaning that for every mole of glycine, there is 3.0×10^{-2} mole of serine. (a) Give the molar ratio of methionine to glycine and explain what it means. (b) How many molecules of glycine are present in 1.0 mole? (c) For every 1.0 mole of glycine in the sample, how many molecules of methionine are present? (Recall that to multiply two



▲ Some of Stanley Miller's notes from his 1958 hydrogen sulfide (H_2S) experiment along with his original vials.

- numbers with exponents, you add their exponents; to divide them, you subtract the exponent in the denominator from that in the numerator.)
2. (a) Which amino acid is present in higher amounts than glycine? (b) How many more molecules of that amino acid are present than the number of molecules in 1.0 mole of glycine?
 3. The synthesis of products is limited by the amount of reactants. (a) If one mole each of CH_4 , NH_3 , H_2S , and CO_2 is added to 1 liter of water (≈ 55.5 moles of H_2O) in a flask, how many moles of hydrogen, carbon, oxygen, nitrogen, and sulfur are in the flask? (b) Looking at the molecular formula in the table, how many moles of each element would be needed to make 1.0 mole of glycine? (c) What is the maximum number of moles of glycine that could be made in that flask, with the specified ingredients, if no other molecules were made? Explain. (d) If serine or methionine were made individually, which element(s) would be used up first for each? How much of each product could be made?
 4. The earlier published experiment carried out by Miller did not include H_2S in the reactants (see Figure 4.2). Which of the compounds shown in the data table can be made in the H_2S experiment but could not be made in the earlier experiment?

MB A version of this Scientific Skills Exercise can be assigned in MasteringBiology.

Data from E. T. Parker et al., Primordial synthesis of amines and amino acids in a 1958 Miller H_2S -rich spark discharge experiment, *Proceedings of the National Academy of Sciences USA* 108:5526-5531 (2011). www.pnas.org/cgi/doi/10.1073/pnas.1019191108.

of organic molecules they make. In a sense, the great diversity of living organisms we see on the planet (and in fossil remains) is made possible by the unique chemical versatility of the element carbon.

CONCEPT CHECK 4.1

1. Why was Wöhler astonished to find he had made urea?
2. **WHAT IF?** Miller carried out a control experiment without the electrical discharge and found no organic compounds. What might explain this result?

For suggested answers, see Appendix A.

CONCEPT 4.2

Carbon atoms can form diverse molecules by bonding to four other atoms

The key to an atom's chemical characteristics is its electron configuration. This configuration determines the kinds and number of bonds an atom will form with other atoms. Recall that it is the valence electrons, those in the outermost shell, that are available to form bonds with other atoms.


Molecule and Molecular Shape	Molecular Formula	Structural Formula	Ball-and-Stick Model (molecular shape in pink)	Space-Filling Model
(a) Methane. When a carbon atom has four single bonds to other atoms, the molecule is tetrahedral.	CH ₄	$ \begin{array}{c} \text{H} \\ \\ \text{H} - \text{C} - \text{H} \\ \\ \text{H} \end{array} $		
(b) Ethane. A molecule may have more than one tetrahedral group of single-bonded atoms. (Ethane consists of two such groups.)	C ₂ H ₆	$ \begin{array}{c} \text{H} \quad \text{H} \\ \quad \\ \text{H} - \text{C} - \text{C} - \text{H} \\ \quad \\ \text{H} \quad \text{H} \end{array} $		
(c) Ethene (ethylene). When two carbon atoms are joined by a double bond, all atoms attached to those carbons are in the same plane, and the molecule is flat.	C ₂ H ₄	$ \begin{array}{c} \text{H} \quad \text{H} \\ \backslash \quad / \\ \text{C} = \text{C} \\ / \quad \backslash \\ \text{H} \quad \text{H} \end{array} $		

▲ **Figure 4.3** The shapes of three simple organic molecules.

The Formation of Bonds with Carbon

Carbon has 6 electrons, with 2 in the first electron shell and 4 in the second shell; thus, it has 4 valence electrons in a shell that can hold up to 8 electrons. A carbon atom usually completes its valence shell by sharing its 4 electrons with other atoms so that 8 electrons are present. Each pair of shared electrons constitutes a covalent bond (see Figure 2.10d). In organic molecules, carbon usually forms single or double covalent bonds. Each carbon atom acts as an intersection point from which a molecule can branch off in as many as four directions. This enables carbon to form large, complex molecules.

When a carbon atom forms four single covalent bonds, the arrangement of its four hybrid orbitals causes the bonds to angle toward the corners of an imaginary tetrahedron. The bond angles in methane (CH₄) are 109.5° (**Figure 4.3a**), and they are roughly the same in any group of atoms where carbon has four single bonds. For example, ethane (C₂H₆) is shaped like two overlapping tetrahedrons (**Figure 4.3b**). In molecules with more carbons, every grouping of a carbon bonded to four other atoms has a tetrahedral shape. But when two carbon atoms are joined by a double bond, as in ethene (C₂H₄), the bonds from both carbons are all in the same plane, so the atoms joined to those carbons are in the same plane as well (**Figure 4.3c**). We find it convenient to write molecules as structural formulas, as if the molecules being represented are two-dimensional, but keep in mind that molecules are three-dimensional and that the shape of a molecule is central to its function.




▲ **Figure 4.4** Valences of the major elements of organic molecules. Valence is the number of covalent bonds an atom can form. It is generally equal to the number of electrons required to complete the valence (outermost) shell (see Figure 2.7). All the electrons are shown for each atom in the electron distribution diagrams (top). Only the valence shell electrons are shown in the Lewis dot structures (bottom). Note that carbon can form four bonds.

MAKE CONNECTIONS Draw the Lewis dot structures for sodium, phosphorus, sulfur, and chlorine. (Refer to Figure 2.7.)

The electron configuration of carbon gives it covalent compatibility with many different elements. **Figure 4.4** shows the valences of carbon and its most frequent bonding partners—hydrogen, oxygen, and nitrogen. These are the four major atomic components of organic molecules. These valences are the basis for the rules of covalent bonding in organic chemistry—the building code for the architecture of organic molecules.


How do the rules of covalent bonding apply to carbon atoms with partners other than hydrogen? We'll look at two examples, the simple molecules carbon dioxide and urea.

In the carbon dioxide molecule (CO_2), a single carbon atom is joined to two atoms of oxygen by double covalent bonds. The structural formula for CO_2 is shown here:



Each line in a structural formula represents a pair of shared electrons. Thus, the two double bonds in CO_2 have the same number of shared electrons as four single bonds. The arrangement completes the valence shells of all atoms in the molecule. Because CO_2 is a very simple molecule and lacks hydrogen, it is often considered inorganic, even though it contains carbon. Whether we call CO_2 organic or inorganic, however, it is clearly important to the living world as the source of carbon for all organic molecules in organisms.

Urea, $\text{CO}(\text{NH}_2)_2$, is the organic compound found in urine that Wöhler synthesized in the early 1800s. Again, each atom has the required number of covalent bonds. In this case, one carbon atom participates in both single and double bonds.



Urea and carbon dioxide are molecules with only one carbon atom. But as Figure 4.3 shows, a carbon atom can also use one or more valence electrons to form covalent bonds to other carbon atoms, each of which can also form four bonds. Thus, the atoms can be linked into chains of seemingly infinite variety.

Molecular Diversity Arising from Variation in Carbon Skeletons

Carbon chains form the skeletons of most organic molecules. The skeletons vary in length and may be straight, branched, or arranged in closed rings (**Figure 4.5**). Some carbon skeletons have double bonds, which vary in number and location. Such variation in carbon skeletons is one important source of the molecular complexity and diversity that characterize living matter. In addition, atoms of other elements can be bonded to the skeletons at available sites.



Hydrocarbons

All of the molecules that are shown in Figures 4.3 and 4.5 are **hydrocarbons**, organic molecules consisting of only carbon and hydrogen. Atoms of hydrogen are attached to the carbon skeleton wherever electrons are available for covalent bonding. Hydrocarbons are the major components of petroleum, which is called a fossil fuel because it consists of the partially decomposed remains of organisms that lived millions of years ago.

Although hydrocarbons are not prevalent in most living organisms, many of a cell's organic molecules have regions consisting of only carbon and hydrogen. For example, the



▼ **Figure 4.5** Four ways that carbon skeletons can vary.

(a) Length





Carbon skeletons vary in length.

(b) Branching





Skeletons may be unbranched or branched.

(c) Double bond position




The skeleton may have double bonds, which can vary in location.

(d) Presence of rings



Some carbon skeletons are arranged in rings. In the abbreviated structural formula for each compound (at the right), each corner represents a carbon and its attached hydrogens.

molecules known as fats have long hydrocarbon tails attached to a nonhydrocarbon component (**Figure 4.6**). Neither petroleum nor fat dissolves in water; both are hydrophobic compounds because the great majority of their bonds are relatively nonpolar carbon-to-hydrogen linkages. Another characteristic of hydrocarbons is that they can undergo reactions that release a relatively large amount of energy. The gasoline that fuels a car consists of hydrocarbons, and the hydrocarbon tails of fats serve as stored fuel for plant embryos (seeds) and animals.



▲ Figure 4.6 The role of hydrocarbons in fats. (a) Mammalian adipose cells stockpile fat molecules as a fuel reserve. This colorized micrograph shows part of a human adipose cell with many fat droplets, each containing a large number of fat molecules. (b) A fat molecule consists of a small, nonhydrocarbon component joined to three hydrocarbon tails that account for the hydrophobic behavior of fats. The tails can be broken down to provide energy. (Black = carbon; gray = hydrogen; red = oxygen.)

MAKE CONNECTIONS How do the tails account for the hydrophobic nature of fats? (See Concept 3.2.)

Isomers


Variation in the architecture of organic molecules can be seen in **isomers**, compounds that have the same numbers of atoms of the same elements but different structures and hence different properties. We will examine three types of isomers: structural isomers, *cis-trans* isomers, and enantiomers.

Structural isomers differ in the covalent arrangements of their atoms. Compare, for example, the two five-carbon compounds in **Figure 4.7a**. Both have the molecular formula C_5H_{12} , but they differ in the covalent arrangement of their carbon skeletons. The skeleton is straight in one compound but branched in the other. The number of possible isomers increases tremendously as carbon skeletons increase in size. There are only three forms of C_5H_{12} (two of which are shown in Figure 4.7a), but there are 18 variations of C_8H_{18} and 366,319 possible structural isomers of $C_{20}H_{42}$. Structural isomers may also differ in the location of double bonds.

In *cis-trans* isomers (formerly called *geometric isomers*), carbons have covalent bonds to the same atoms, but these atoms differ in their spatial arrangements due to the inflexibility of double bonds. Single bonds allow the atoms they join to rotate freely about the bond axis without changing the compound. In contrast, double bonds do not permit such rotation. If a double bond joins two carbon atoms, and each C also has two different atoms (or groups of atoms) attached to it, then two distinct *cis-trans* isomers are possible. Consider a simple molecule with two double-bonded carbons, each of which has an H and an X attached to it (**Figure 4.7b**). The arrangement with both Xs on the same side of the double bond is called a *cis isomer*, and that with the Xs on opposite sides is

▼ Figure 4.7 Three types of isomers, compounds with the same molecular formula but different structures.

(a) Structural isomers



Structural isomers differ in covalent partners, as shown in this example of two isomers of C_5H_{12} .

(b) *Cis-trans* isomers




cis isomer: The two Xs are on the same side.

trans isomer: The two Xs are on opposite sides.

Cis-trans isomers differ in arrangement about a double bond. In these diagrams, X represents an atom or group of atoms attached to a double-bonded carbon.

(c) Enantiomers




Enantiomers differ in spatial arrangement around an asymmetric carbon, resulting in molecules that are mirror images, like left and right hands. The two isomers here are designated the L and D isomers from the Latin for "left" and "right" (*levo* and *dextro*). Enantiomers cannot be superimposed on each other.

© Pearson Education, Inc.

DRAW IT There are three structural isomers of C_5H_{12} ; draw the one not shown in (a).

called a *trans isomer*. The subtle difference in shape between such isomers can dramatically affect the biological activities of organic molecules. For example, the biochemistry of vision involves a light-induced change of retinal, a chemical compound in the eye, from the *cis* isomer to the *trans* isomer (see Figure 50.17). Another example involves *trans* fats, which are discussed in Chapter 5.

Enantiomers are isomers that are mirror images of each other and that differ in shape due to the presence of an *asymmetric carbon*, one that is attached to four different atoms or groups of atoms. (See the middle carbon in

Drug	Effects	Effective Enantiomer	Ineffective Enantiomer
Ibuprofen	Reduces inflammation and pain	 S-Ibuprofen	 R-Ibuprofen
Albuterol	Relaxes bronchial (airway) muscles, improving airflow in asthma patients	 R-Albuterol	 S-Albuterol

▲ **Figure 4.8** The pharmacological importance of enantiomers. Ibuprofen and albuterol are drugs whose enantiomers have different effects. (*S* and *R* are used here to distinguish between enantiomers.) Ibuprofen is commonly sold as a mixture of the two enantiomers; the *S* enantiomer is 100 times more effective than the *R* form. Albuterol is synthesized and sold only as the *R* form of the drug; the *S* form counteracts the active *R* form.

the ball-and-stick models shown in **Figure 4.7c**.) The four groups can be arranged in space around the asymmetric carbon in two different ways that are mirror images. Enantiomers are, in a way, left-handed and right-handed versions of the molecule. Just as your right hand won't fit into a left-handed glove, a "right-handed" molecule won't fit into the same space as the "left-handed" version. Usually, only one isomer is biologically active because only that form can bind to specific molecules in an organism.

The concept of enantiomers is important in the pharmaceutical industry because the two enantiomers of a drug may not be equally effective, as is the case for both ibuprofen and the asthma medication albuterol (**Figure 4.8**). Methamphetamine also occurs in two enantiomers that have very different effects. One enantiomer is the highly addictive stimulant drug known as "crank," sold illegally in the street drug trade. The other has a much weaker effect and is the active ingredient in an over-the-counter vapor inhaler for treatment of nasal congestion. The differing effects of enantiomers in the body demonstrate that organisms are sensitive to even the most subtle variations in molecular architecture. Once again, we see that molecules have emergent properties that depend on the specific arrangement of their atoms.

CONCEPT CHECK 4.2

- DRAW IT** (a) Draw a structural formula for C_2H_4 . (b) Draw the *trans* isomer of $C_2H_2Cl_2$.
- Which molecules in Figure 4.5 are isomers? For each pair, identify the type of isomer.
- How are gasoline and fat chemically similar?
- Can propane (C_3H_8) form isomers? Explain.

For suggested answers, see Appendix A.


CONCEPT 4.3

A few chemical groups are key to molecular function

The properties of an organic molecule depend not only on the arrangement of its carbon skeleton but also on the chemical groups attached to that skeleton. We can think of hydrocarbons, the simplest organic molecules, as the underlying framework for more complex organic molecules. A number of chemical groups can replace one or more hydrogens of the hydrocarbon. These groups may participate in chemical reactions or may contribute to function indirectly by their effects on molecular shape; they help give each molecule its unique properties.

The Chemical Groups Most Important in the Processes of Life

Consider the differences between estradiol (a type of estrogen) and testosterone. These compounds are female and male sex hormones, respectively, in humans and other vertebrates. Both are steroids, organic molecules with a common carbon skeleton in the form of four fused rings. They differ only in the chemical groups attached to the rings (shown here in abbreviated form); the distinctions in molecular architecture are shaded in blue:




















The different actions of these two molecules on many targets throughout the body are the basis of gender, producing the contrasting features of male and female vertebrates. In this case, the chemical groups are important because they affect molecular shape, contributing to function.

In other cases, chemical groups are directly involved in chemical reactions; such groups are known as **functional groups**. Each has certain properties, such as shape and charge, that cause it to participate in chemical reactions in a characteristic way.


The seven chemical groups most important in biological processes are the hydroxyl, carbonyl, carboxyl, amino, sulfhydryl, phosphate, and methyl groups. The first six groups can be chemically reactive; of these, all except the sulfhydryl group are also hydrophilic and thus increase the solubility of organic compounds in water. The methyl group is not reactive, but instead often serves as a recognizable tag on biological molecules. Study **Figure 4.9** to become familiar with these biologically important chemical groups.

▼ **Figure 4.9** Some biologically important chemical groups.


Chemical Group	Group Properties and Compound Name	Examples
Hydroxyl group ($-\text{OH}$)  (may be written $\text{HO}-$)	Is polar due to electronegative oxygen. Forms hydrogen bonds with water, helping dissolve compounds such as sugars. Compound name: Alcohol (specific name usually ends in <i>-ol</i>)	 Ethanol , the alcohol present in alcoholic beverages
Carbonyl group ($>\text{C}=\text{O}$) 	Sugars with ketone groups are called ketoses; those with aldehydes are called aldoses. Compound name: Ketone (carbonyl group is within a carbon skeleton) or aldehyde (carbonyl group is at the end of a carbon skeleton)	 Acetone , the simplest ketone  Propanal , an aldehyde
Carboxyl group ($-\text{COOH}$) 	Acts as an acid (can donate H^+) because the covalent bond between oxygen and hydrogen is so polar. Compound name: Carboxylic acid, or organic acid	 $\rightleftharpoons$  + H^+ Acetic acid , which gives vinegar its sour taste Ionized form of $-\text{COOH}$ (carboxylate ion), found in cells
Amino group ($-\text{NH}_2$) 	Acts as a base; can pick up an H^+ from the surrounding solution (water, in living organisms). Compound name: Amine	 + H^+ $\rightleftharpoons$  Glycine , an amino acid (note its carboxyl group) Ionized form of $-\text{NH}_2$, found in cells
Sulfhydryl group ($-\text{SH}$)  (may be written $\text{HS}-$)	Two $-\text{SH}$ groups can react, forming a "cross-link" that helps stabilize protein structure. Hair protein cross-links maintain the straightness or curliness of hair; in hair salons, permanent treatments break cross-links, then re-form them while the hair is in the desired shape. Compound name: Thiol	 Cysteine , a sulfur-containing amino acid
Phosphate group ($-\text{OPO}_3^{2-}$) 	Contributes negative charge (1- when positioned inside a chain of phosphates; 2- when at the end). When attached, confers on a molecule the ability to react with water, releasing energy. Compound name: Organic phosphate	 Glycerol phosphate , which takes part in many important chemical reactions in cells
Methyl group ($-\text{CH}_3$) 	Affects the expression of genes when on DNA or on proteins bound to DNA. Affects the shape and function of male and female sex hormones. Compound name: Methylated compound	 5-Methyl cytosine , a component of DNA that has been modified by addition of a methyl group

ATP: An Important Source of Energy for Cellular Processes

The “Phosphate group” row in Figure 4.9 shows a simple example of an organic phosphate molecule. A more complicated organic phosphate, **adenosine triphosphate**, or **ATP**, is worth mentioning here because its function in the cell is so important. ATP consists of an organic molecule called adenosine attached to a string of three phosphate groups:



Where three phosphates are present in series, as in ATP, one phosphate may be split off as a result of a reaction with water. This inorganic phosphate ion, HOPO_3^{2-} , is often abbreviated P_i in this book, and a phosphate group in an organic molecule is often written as P . Having lost one phosphate, ATP becomes adenosine *diphosphate*, or ADP. Although ATP is sometimes said to store energy, it is more accurate to think of it as storing the potential to react with water. This reaction releases energy that can be used by the cell. You will learn about this in more detail in Chapter 8.



CONCEPT CHECK 4.3

1. What does the term *amino acid* signify about the structure of such a molecule?
2. What chemical change occurs to ATP when it reacts with water and releases energy?
3. **WHAT IF?** Suppose you had an organic molecule such as cysteine (see Figure 4.9, sulfhydryl group example), and you chemically removed the —NH_2 group and replaced it with —COOH . Draw the structural formula for this molecule and speculate about its chemical properties. Is the central carbon asymmetric before the change? After?

For suggested answers, see Appendix A.

The Chemical Elements of Life: A Review

Living matter, as you have learned, consists mainly of carbon, oxygen, hydrogen, and nitrogen, with smaller amounts of sulfur and phosphorus. These elements all form strong covalent bonds, an essential characteristic in the architecture of complex organic molecules. Of all these elements, carbon is the virtuoso of the covalent bond. The versatility of carbon makes possible the great diversity of organic molecules, each with particular properties that emerge from the unique arrangement of its carbon skeleton and the chemical groups appended to that skeleton. This variation at the molecular level provides the foundation for the rich biological diversity found on our planet.

4 Chapter Review

SUMMARY OF KEY CONCEPTS

CONCEPT 4.1

Organic chemistry is the study of carbon compounds (pp. 57–58)

- Organic compounds, once thought to arise only within living organisms, were finally synthesized in the laboratory.
- Living matter is made mostly of carbon, oxygen, hydrogen, and nitrogen. Biological diversity results from carbon's ability to form a huge number of molecules with particular shapes and properties.

? How did Stanley Miller's experiments support the idea that, even at life's origins, physical and chemical laws govern the processes of life?

CONCEPT 4.2

Carbon atoms can form diverse molecules by bonding to four other atoms (pp. 58–62)

- Carbon, with a valence of 4, can bond to various other atoms, including O, H, and N. Carbon can also bond to other carbon

atoms, forming the carbon skeletons of organic compounds.

These skeletons vary in length and shape and have bonding sites for atoms of other elements.

- **Hydrocarbons** consist of carbon and hydrogen.
- **Isomers** are compounds that have the same molecular formula but different structures and therefore different properties. Three types of isomers are **structural isomers**, **cis-trans isomers**, and **enantiomers**.


? Refer back to Figure 4.9. What type of isomers are acetone and propanal? How many asymmetric carbons are present in acetic acid, glycine, and glycerol phosphate? Can these three molecules exist as forms that are enantiomers?

CONCEPT 4.3

A few chemical groups are key to molecular function (pp. 62–64)

- Chemical groups attached to the carbon skeletons of organic molecules participate in chemical reactions (**functional groups**) or contribute to function by affecting molecular shape (see Figure 4.9).
- **ATP (adenosine triphosphate)** consists of adenosine attached to three phosphate groups. ATP can react with water, forming

inorganic phosphate and ADP (adenosine diphosphate). This reaction releases energy that can be used by the cell.



? In what ways does a methyl group differ chemically from the other six important chemical groups shown in Figure 4.9?


TEST YOUR UNDERSTANDING

LEVEL 1: KNOWLEDGE/COMPREHENSION

- Organic chemistry is currently defined as
 - the study of compounds made only by living cells.
 - the study of carbon compounds.
 - the study of natural (as opposed to synthetic) compounds.
 - the study of hydrocarbons.

- Which functional group is *not* present in this molecule?

- carboxyl
- sulfhydryl
- hydroxyl
- amino



- MAKE CONNECTIONS** Which chemical group is most likely to be responsible for an organic molecule behaving as a base (see Concept 3.3)?

- hydroxyl
- carbonyl
- amino
- phosphate


LEVEL 2: APPLICATION/ANALYSIS

- Which of the following hydrocarbons has a double bond in its carbon skeleton?


- C_3H_8
- C_2H_6
- C_2H_4
- C_2H_2

- Choose the term that correctly describes the relationship between these two sugar molecules:

- structural isomers
- cis-trans* isomers
- enantiomers
- isotopes



- Identify the asymmetric carbon in this molecule:



- Which action could produce a carbonyl group?
 - the replacement of the —OH of a carboxyl group with hydrogen
 - the addition of a thiol to a hydroxyl
 - the addition of a hydroxyl to a phosphate
 - the replacement of the nitrogen of an amine with oxygen
- Which of the molecules shown in question 5 has an asymmetric carbon? Which carbon is asymmetric?

LEVEL 3: SYNTHESIS/EVALUATION

9. EVOLUTION CONNECTION

DRAW IT Some scientists think that life elsewhere in the universe might be based on the element silicon, rather than on carbon, as on Earth. Look at the electron distribution diagram for silicon in Figure 2.7 and draw the Lewis dot structure for silicon. What properties does silicon share with carbon that would make silicon-based life more likely than, say, neon-based life or aluminum-based life?

10. SCIENTIFIC INQUIRY

50 years ago, pregnant women who were prescribed thalidomide for morning sickness gave birth to children with birth defects. Thalidomide is a mixture of two enantiomers; one reduces morning sickness, but the other causes severe birth defects. Today, the FDA has approved this drug for non-pregnant individuals with Hansen's disease (leprosy) or newly diagnosed multiple myeloma, a blood and bone marrow cancer. The beneficial enantiomer can be synthesized and given to patients, but over time, both the beneficial *and* the harmful enantiomer can be detected in the body. Propose a possible explanation for the presence of the harmful enantiomer.

11. WRITE ABOUT A THEME: ORGANIZATION

In 1918, an epidemic of sleeping sickness caused an unusual rigid paralysis in some survivors, similar to symptoms of advanced Parkinson's disease. Years later, L-dopa (below, left), a chemical used to treat Parkinson's disease, was given to some of these patients. L-dopa was remarkably effective at eliminating the paralysis, at least temporarily. However, its enantiomer, D-dopa (right), was subsequently shown to have no effect at all, as is the case for Parkinson's disease. In a short essay (100–150 words), discuss how the effectiveness of one enantiomer and not the other illustrates the theme of structure and function.



L-dopa

D-dopa

12. SYNTHESIZE YOUR KNOWLEDGE



Explain how the chemical structure of the carbon atom accounts for the differences between the male and female lions seen in the photo.

For selected answers, see Appendix A.

MasteringBiology®

Students Go to **MasteringBiology** for assignments, the eText, and the Study Area with practice tests, animations, and activities.

Instructors Go to **MasteringBiology** for automatically graded tutorials and questions that you can assign to your students, plus Instructor Resources.