

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that  $A[j] - A[i] = k$ ,  $i \neq j$ .

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as  $5 - 1 = 4$

So Return 1.

**For example:**

Input	Result
3	1
1 3 5	
4	

**Answer:** (penalty regime: 0 %)

```
1 #include <stdio.h>
2
3 int main() {
4     int n;
5     scanf("%d", &n);
6     int arr[n];
7     for (int i = 0; i < n; i++)
8         scanf("%d", &arr[i]);
9
10    int k;
11    scanf("%d", &k);
12
13    int i = 0, j = 1;
14    while (i < n && j < n) {
15        if (i != j && arr[j] - arr[i] == k) {
16            printf("1");
17            return 0;
18        } else if (arr[j] - arr[i] < k) {
19            j++;
20        } else {
21            i++;
22        }
23    }
24    printf("0");
25    return 0;
26 }
27
```

	Input	Expected	Got	
✓	3 1 3 5 4	1	1	✓
✓	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✓
✓	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✓
✓	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✓

Passed all tests! ✓

**Correct**

Marks for this submission: 1.00/1.00.