

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Definition Suppose ξ and $\{\xi_n, n \geq 1\}$, are defined on the same probability space (Ω, \mathcal{F}, P) . If for any $\varepsilon > 0$

$$\lim_{n \rightarrow \infty} P(|\xi_n - \xi| \geq \varepsilon) = 0,$$

or equivalently $\lim_{n \rightarrow \infty} P(|\xi_n - \xi| < \varepsilon) = 1$, then we say that ξ_n converges to ξ in probability, written $\xi_n \xrightarrow{P} \xi$.

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Throwing a dot in $[0, 1]$ randomly, the dot is located any point in $[0, 1]$ with the same possibility. Let ω denote the location of dot and define

$$\xi(\omega) = \begin{cases} 1, & \omega \in [0, 0.5], \\ 0, & \omega \in (0.5, 1], \end{cases} \quad \eta(\omega) = \begin{cases} 0, & \omega \in [0, 0.5], \\ 1, & \omega \in (0.5, 1]. \end{cases}$$

Then ξ and η have the same distribution function

$$F(x) = \begin{cases} 0, & x < 0, \\ \frac{1}{2}, & 0 \leq x < 1, \\ 1, & x \geq 1. \end{cases}$$

If we define $\xi_n = \xi$, for $n \geq 1$, then $\xi_n \xrightarrow{d} \eta$, but $|\xi_n - \eta| \equiv 1$.

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

- ① Suppose ξ and $\{\xi_n, n \geq 1\}$ are random variables defined on the probability space (Ω, \mathcal{F}, P) .
- (1) If $\xi_n \xrightarrow{P} \xi$, then $\xi_n \xrightarrow{d} \xi$.
- (2) If $\xi_n \xrightarrow{d} c$, where c is a **constant**, then $\xi_n \xrightarrow{P} c$.

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

① Suppose ξ and $\{\xi_n, n \geq 1\}$ are random variables defined on the probability space (Ω, \mathcal{F}, P) .

(1) If $\xi_n \xrightarrow{P} \xi$, then $\xi_n \xrightarrow{d} \xi$.

(2) If $\xi_n \xrightarrow{d} c$, where c is a **constant**, then $\xi_n \xrightarrow{P} c$.

Proof. (1) Let F and F_n be the cdfs of ξ and ξ_n respectively, and let x be a continuity point of F .

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

For any $\varepsilon > 0$,

$$\begin{aligned}(\xi_n \leq x) &= (\xi_n \leq x, |\xi_n - \xi| < \varepsilon) \\ &\quad + (\xi_n \leq x, |\xi_n - \xi| \geq \varepsilon) \\ &\subset (\xi \leq x + \varepsilon) \cup (|\xi_n - \xi| \geq \varepsilon).\end{aligned}$$

For any $\varepsilon > 0$,

$$\begin{aligned}(\xi_n \leq x) &= (\xi_n \leq x, |\xi_n - \xi| < \varepsilon) \\ &\quad + (\xi_n \leq x, |\xi_n - \xi| \geq \varepsilon) \\ &\subset (\xi \leq x + \varepsilon) \cup (|\xi_n - \xi| \geq \varepsilon).\end{aligned}$$

Thus

$$F_n(x) \leq F(x + \varepsilon) + P(|\xi_n - \xi| \geq \varepsilon).$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Since $\xi_n \xrightarrow{P} \xi$ as $n \rightarrow \infty$, we obtain

$$P(|\xi_n - \xi| \geq \varepsilon) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Since $\xi_n \xrightarrow{P} \xi$ as $n \rightarrow \infty$, we obtain

$$P(|\xi_n - \xi| \geq \varepsilon) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Thus

$$\limsup_{n \rightarrow \infty} F_n(x) \leq F(x + \varepsilon).$$

Similarly

$$(\xi \leq x) \subset (\xi_n \leq x + \varepsilon) \cup (|\xi - \xi_n| \geq \varepsilon)$$

and thus

$$F(x) \leq F_n(x + \varepsilon) + P(|\xi_n - \xi| \geq \varepsilon).$$

Similarly

$$(\xi \leq x) \subset (\xi_n \leq x + \varepsilon) \cup (|\xi - \xi_n| \geq \varepsilon)$$

and thus

$$F(x) \leq F_n(x + \varepsilon) + P(|\xi_n - \xi| \geq \varepsilon).$$

So

$$F(x - \varepsilon) \leq F_n(x) + P(|\xi_n - \xi| \geq \varepsilon).$$

Similarly

$$(\xi \leq x) \subset (\xi_n \leq x + \varepsilon) \cup (|\xi - \xi_n| \geq \varepsilon)$$

and thus

$$F(x) \leq F_n(x + \varepsilon) + P(|\xi_n - \xi| \geq \varepsilon).$$

So

$$F(x - \varepsilon) \leq F_n(x) + P(|\xi_n - \xi| \geq \varepsilon).$$

Thus

$$F(x - \varepsilon) \leq \liminf_{n \rightarrow \infty} F_n(x).$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

We conclude that

$$F(x-\varepsilon) \leq \liminf_{n \rightarrow \infty} F_n(x) \leq \limsup_{n \rightarrow \infty} F_n(x) \leq F(x+\varepsilon).$$

We conclude that

$$F(x-\varepsilon) \leq \liminf_{n \rightarrow \infty} F_n(x) \leq \limsup_{n \rightarrow \infty} F_n(x) \leq F(x+\varepsilon).$$

Letting $\varepsilon \rightarrow 0$ yields

$$\lim_{n \rightarrow \infty} F_n(x) = F(x).$$

That is

$$\xi_n \xrightarrow{d} \xi.$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

(2) If $\xi_n \xrightarrow{d} c$, then

$$\lim_{n \rightarrow \infty} F_n(x) = \begin{cases} 0, & x < c, \\ 1, & x > c. \end{cases}$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

(2) If $\xi_n \xrightarrow{d} c$, then

$$\lim_{n \rightarrow \infty} F_n(x) = \begin{cases} 0, & x < c, \\ 1, & x > c. \end{cases}$$

Hence for any $\varepsilon > 0$,

$$\begin{aligned} & P(|\xi_n - c| \geq \varepsilon) \\ &= P(\xi_n \geq c + \varepsilon) + P(\xi_n \leq c - \varepsilon) \\ &= 1 - P(\xi_n < c + \varepsilon) + P(\xi_n \leq c - \varepsilon) \\ &= 1 - F_n(c + \varepsilon - 0) + F_n(c - \varepsilon) \\ &\rightarrow 0 \quad (n \rightarrow \infty). \end{aligned}$$

The proof is complete.

Example 1 Let $\{\xi_n\}$ be a sequence of i.i.d. random variables with the common uniform distribution in $[0, a]$. Let $\eta_n = \max\{\xi_1, \xi_2, \dots, \xi_n\}$. Prove that $\eta_n \xrightarrow{P} a$.

Proof.

Example 1 Let $\{\xi_n\}$ be a sequence of i.i.d. random variables with the common uniform distribution in $[0, a]$. Let $\eta_n = \max\{\xi_1, \xi_2, \dots, \xi_n\}$. Prove that $\eta_n \xrightarrow{P} a$.

Proof. Let $F(x)$ be the distribution function of ξ_k . then the distribution function of η_n is $G_n(x) = (F(x))^n$.

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Now the distribution function of ξ_k is

$$F(x) = \begin{cases} 0, & x < 0, \\ x/a, & 0 \leq x < a, \\ 1, & x \geq a. \end{cases}$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Now the distribution function of ξ_k is

$$F(x) = \begin{cases} 0, & x < 0, \\ x/a, & 0 \leq x < a, \\ 1, & x \geq a. \end{cases}$$

Hence

$$\begin{aligned} G_n(x) &= \begin{cases} 0, & x < 0, \\ (x/a)^n, & 0 \leq x < a, \\ 1, & x \geq a, \end{cases} \\ &\rightarrow D(x-a) = \begin{cases} 0, & x < a, \\ 1, & x \geq a, \end{cases} \quad \text{as } n \rightarrow \infty. \end{aligned}$$

So $\eta_n \xrightarrow{d} a$ and a is a constant. So $\eta_n \xrightarrow{P} a$.

- 2 Let $\{\xi, \xi_n, n \geq 1\}$ be a sequence of random variables defined on the probability space (Ω, \mathcal{F}, P) . Prove that
- (1) If $\xi_n \xrightarrow{P} \xi$, $\xi_n \xrightarrow{P} \eta$, then $P(\xi = \eta) = 1$.
 - (2) If $\xi_n \xrightarrow{P} \xi$, f is the continuous function on $(-\infty, \infty)$, then $f(\xi_n) \xrightarrow{P} f(\xi)$.

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

- 2 Let $\{\xi, \xi_n, n \geq 1\}$ be a sequence of random variables defined on the probability space (Ω, \mathcal{F}, P) . Prove that
- (1) If $\xi_n \xrightarrow{P} \xi$, $\xi_n \xrightarrow{P} \eta$, then $P(\xi = \eta) = 1$.
 - (2) If $\xi_n \xrightarrow{P} \xi$, f is the continuous function on $(-\infty, \infty)$, then $f(\xi_n) \xrightarrow{P} f(\xi)$.

In general, if $\xi_n \xrightarrow{P} \xi$, $\eta_n \xrightarrow{P} \eta$ and $f(x, y)$ is a continuous function, then

$$f(\xi_n, \eta_n) \xrightarrow{P} f(\xi, \eta).$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Proof. (1) For any $\varepsilon > 0$, we have

$$(|\xi - \eta| \geq \varepsilon) \subseteq (|\xi_n - \xi| \geq \frac{\varepsilon}{2}) \cup (|\xi_n - \eta| \geq \frac{\varepsilon}{2}).$$

Thus

$$P(|\xi - \eta| \geq \varepsilon) \leq P(|\xi_n - \xi| \geq \frac{\varepsilon}{2}) + P(|\xi_n - \eta| \geq \frac{\varepsilon}{2}).$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Proof. (1) For any $\varepsilon > 0$, we have

$$(|\xi - \eta| \geq \varepsilon) \subseteq (|\xi_n - \xi| \geq \frac{\varepsilon}{2}) \cup (|\xi_n - \eta| \geq \frac{\varepsilon}{2}).$$

Thus

$$P(|\xi - \eta| \geq \varepsilon) \leq P(|\xi_n - \xi| \geq \frac{\varepsilon}{2}) + P(|\xi_n - \eta| \geq \frac{\varepsilon}{2}).$$

Note that the left side of the above inequality is independent of n and $\xi_n \xrightarrow{P} \xi$, $\xi_n \xrightarrow{P} \eta$ as $n \rightarrow \infty$. Therefore

$$P(|\xi - \eta| \geq \varepsilon) = 0.$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Proof. (1) For any $\varepsilon > 0$, we have

$$(|\xi - \eta| \geq \varepsilon) \subseteq (|\xi_n - \xi| \geq \frac{\varepsilon}{2}) \cup (|\xi_n - \eta| \geq \frac{\varepsilon}{2}).$$

Thus

$$P(|\xi - \eta| \geq \varepsilon) \leq P(|\xi_n - \xi| \geq \frac{\varepsilon}{2}) + P(|\xi_n - \eta| \geq \frac{\varepsilon}{2}).$$

Note that the left side of the above inequality is independent of n and $\xi_n \xrightarrow{P} \xi$, $\xi_n \xrightarrow{P} \eta$ as $n \rightarrow \infty$. Therefore

$P(|\xi - \eta| \geq \varepsilon) = 0$. Furthermore,

$$\begin{aligned} P(|\xi - \eta| > 0) &= P\left(\bigcup_{n=1}^{\infty} (|\xi - \eta| \geq \frac{1}{n})\right) \\ &\leq \sum_{n=1}^{\infty} P(|\xi - \eta| \geq \frac{1}{n}) = 0, \end{aligned}$$

i.e., $P(\xi = \eta) = 1$.

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

(2) For any given $\varepsilon' > 0$, there exists an $M > 0$ satisfying

$$P(|\xi| \geq M) \leq P(|\xi| \geq \frac{M}{2}) \leq \frac{\varepsilon'}{4}. \quad (1)$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

(2) For any given $\varepsilon' > 0$, there exists an $M > 0$ satisfying

$$P(|\xi| \geq M) \leq P(|\xi| \geq \frac{M}{2}) \leq \frac{\varepsilon'}{4}. \quad (1)$$

Since $\xi_n \xrightarrow{P} \xi$, when $n \geq N_1$ for some $N_1 \geq 1$,

$$P(|\xi_n - \xi| \geq \frac{M}{2}) \leq \frac{\varepsilon'}{4}.$$

Hence

$$\begin{aligned} P(|\xi_n| \geq M) &\leq P(|\xi_n - \xi| \geq \frac{M}{2}) + P(|\xi| \geq \frac{M}{2}) \\ &\leq \frac{\varepsilon'}{4} + \frac{\varepsilon'}{4} = \frac{\varepsilon'}{2}. \end{aligned} \quad (2)$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

(2) For any given $\varepsilon' > 0$, there exists an $M > 0$ satisfying

$$P(|\xi| \geq M) \leq P(|\xi| \geq \frac{M}{2}) \leq \frac{\varepsilon'}{4}. \quad (1)$$

Since $\xi_n \xrightarrow{P} \xi$, when $n \geq N_1$ for some $N_1 \geq 1$,

$$P(|\xi_n - \xi| \geq \frac{M}{2}) \leq \frac{\varepsilon'}{4}.$$

Hence

$$\begin{aligned} P(|\xi_n| \geq M) &\leq P(|\xi_n - \xi| \geq \frac{M}{2}) + P(|\xi| \geq \frac{M}{2}) \\ &\leq \frac{\varepsilon'}{4} + \frac{\varepsilon'}{4} = \frac{\varepsilon'}{2}. \end{aligned} \quad (2)$$

And for $f(x)$ is continuous function on $(-\infty, \infty)$, then $f(x)$ is uniformly continuous in $[-M, M]$. For given $\varepsilon > 0$, there exists $\delta > 0$, when $|x - y| < \delta$, $|f(x) - f(y)| < \varepsilon$.

Thus

$$\begin{aligned} & P(|f(\xi_n) - f(\xi)| \geq \varepsilon) \\ \leq & P(|f(\xi_n) - f(\xi)| \geq \varepsilon, |\xi_n - \xi| < \delta, |\xi_n| < M, |\xi| < M) \\ & + P(|f(\xi_n) - f(\xi)| \geq \varepsilon, |\xi_n - \xi| \geq \delta, |\xi_n| < M, |\xi| < M) \\ & + P(|\xi_n| \geq M) + P(|\xi| \geq M) \\ \leq & P(|\xi_n - \xi| \geq \delta) + P(|\xi_n| \geq M) + P(|\xi| \geq M). \end{aligned} \quad (3)$$

Thus

$$\begin{aligned} & P(|f(\xi_n) - f(\xi)| \geq \varepsilon) \\ & \leq P(|f(\xi_n) - f(\xi)| \geq \varepsilon, |\xi_n - \xi| < \delta, |\xi_n| < M, |\xi| < M) \\ & \quad + P(|f(\xi_n) - f(\xi)| \geq \varepsilon, |\xi_n - \xi| \geq \delta, |\xi_n| < M, |\xi| < M) \\ & \quad + P(|\xi_n| \geq M) + P(|\xi| \geq M) \\ & \leq P(|\xi_n - \xi| \geq \delta) + P(|\xi_n| \geq M) + P(|\xi| \geq M). \end{aligned} \quad (3)$$

For the above δ , when $n \geq N_2$ for some $N_2 \geq 1$,

$$P(|\xi_n - \xi| \geq \delta) \leq \frac{\varepsilon'}{4}. \quad (4)$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Thus

$$\begin{aligned} & P(|f(\xi_n) - f(\xi)| \geq \varepsilon) \\ & \leq P(|f(\xi_n) - f(\xi)| \geq \varepsilon, |\xi_n - \xi| < \delta, |\xi_n| < M, |\xi| < M) \\ & \quad + P(|f(\xi_n) - f(\xi)| \geq \varepsilon, |\xi_n - \xi| \geq \delta, |\xi_n| < M, |\xi| < M) \\ & \quad + P(|\xi_n| \geq M) + P(|\xi| \geq M) \\ & \leq P(|\xi_n - \xi| \geq \delta) + P(|\xi_n| \geq M) + P(|\xi| \geq M). \end{aligned} \quad (3)$$

For the above δ , when $n \geq N_2$ for some $N_2 \geq 1$,

$$P(|\xi_n - \xi| \geq \delta) \leq \frac{\varepsilon'}{4}. \quad (4)$$

Combining (1), (2), (3) and (4), we obtain

$$P(|f(\xi_n) - f(\xi)| \geq \varepsilon) \leq \frac{\varepsilon'}{4} + \frac{\varepsilon'}{2} + \frac{\varepsilon'}{4} = \varepsilon'$$

provided $n \geq \max\{N_1, N_2\}$. Thus $f(\xi_n) \xrightarrow{P} f(\xi)$.

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

In general, if

$$\boldsymbol{\xi}_n =: (\xi_{n,1}, \dots, \xi_{n,m}) \xrightarrow{P} \boldsymbol{\xi} := (\xi_1, \dots, \xi_m)$$

(i.e., $\|\boldsymbol{\xi}_n - \boldsymbol{\xi}\| \rightarrow 0$, or equivalently, $\xi_{n,k} \rightarrow \xi_k$,
 $k = 1, \dots, m$)

and $f(\boldsymbol{x})$ is a m -dimensional continuous function,
then

$$f(\boldsymbol{\xi}_n) \xrightarrow{P} f(\boldsymbol{\xi}).$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Proof. For any $\epsilon > 0$ and $M > 0$, choose $0 < \delta < M/2$ such that $|f(\mathbf{x}) - f(\mathbf{y})| < \epsilon$ whenever $\|\mathbf{x} - \mathbf{y}\| < \delta$, $\|\mathbf{x}\| \leq M$, $\|\mathbf{y}\| \leq M$.

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Proof. For any $\epsilon > 0$ and $M > 0$, choose $0 < \delta < M/2$ such that $|f(\mathbf{x}) - f(\mathbf{y})| < \epsilon$ whenever

$\|\mathbf{x} - \mathbf{y}\| < \delta, \|\mathbf{x}\| \leq M, \|\mathbf{y}\| \leq M$. Then

$$\begin{aligned} & \{|f(\mathbf{x}) - f(\mathbf{y})| \geq \epsilon\} \\ & \subset \{\|\mathbf{x} - \mathbf{y}\| \geq \delta\} \cup \{\|\mathbf{x}\| > M\} \cup \{\|\mathbf{y}\| > M\} \\ & \subset \{\|\mathbf{x} - \mathbf{y}\| \geq \delta\} \cup \{\|\mathbf{y}\| > M/2\}. \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Proof. For any $\epsilon > 0$ and $M > 0$, choose $0 < \delta < M/2$ such that $|f(\mathbf{x}) - f(\mathbf{y})| < \epsilon$ whenever

$\|\mathbf{x} - \mathbf{y}\| < \delta, \|\mathbf{x}\| \leq M, \|\mathbf{y}\| \leq M$. Then

$$\begin{aligned} & \{|f(\mathbf{x}) - f(\mathbf{y})| \geq \epsilon\} \\ & \subset \{\|\mathbf{x} - \mathbf{y}\| \geq \delta\} \cup \{\|\mathbf{x}\| > M\} \cup \{\|\mathbf{y}\| > M\} \\ & \subset \{\|\mathbf{x} - \mathbf{y}\| \geq \delta\} \cup \{\|\mathbf{y}\| > M/2\}. \end{aligned}$$

So,

$$\begin{aligned} & P(|f(\boldsymbol{\xi}_n) - f(\boldsymbol{\xi})| \geq \epsilon) \\ & \leq P(\|\boldsymbol{\xi}_n - \boldsymbol{\xi}\| \geq \delta) + P(\|\boldsymbol{\xi}\| > M/2) \rightarrow 0, \end{aligned}$$

as $n \rightarrow \infty$ and then $M \rightarrow \infty$.

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

③ We have

① If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} \eta$, then $\xi_n \pm \eta_n \xrightarrow{P} \xi \pm \eta$;

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

③ We have

- ① If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} \eta$, then $\xi_n \pm \eta_n \xrightarrow{P} \xi \pm \eta$;
- ② If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} \eta$, then $\xi_n \eta_n \xrightarrow{P} \xi \eta$;

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

3 We have

- ① If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} \eta$, then $\xi_n \pm \eta_n \xrightarrow{P} \xi \pm \eta$;
- ② If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} \eta$, then $\xi_n \eta_n \xrightarrow{P} \xi \eta$;
- ③ If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} c$, where c is a constant, both η_n and c are not 0, then $\xi_n / \eta_n \xrightarrow{P} \xi / c$;

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

③ We have

- ① If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} \eta$, then $\xi_n \pm \eta_n \xrightarrow{P} \xi \pm \eta$;
- ② If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} \eta$, then $\xi_n \eta_n \xrightarrow{P} \xi \eta$;
- ③ If $\xi_n \xrightarrow{P} \xi, \eta_n \xrightarrow{P} c$, where c is a constant, both η_n and c are not 0, then $\xi_n / \eta_n \xrightarrow{P} \xi / c$;
- ④ If $\xi_n \xrightarrow{d} \xi, \eta_n \xrightarrow{P} c$, where c is a constant, then $\xi_n + \eta_n \xrightarrow{d} \xi + c, \eta_n \xi_n \xrightarrow{d} c\xi$.

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Proof. We only give a proof of (3) and (4).

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Proof. We only give a proof of (3) and (4). By (2), it is sufficient to show that $\eta_n^{-1} \xrightarrow{P} c^{-1}$.

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Proof. We only give a proof of (3) and (4). By (2), it is sufficient to show that $\eta_n^{-1} \xrightarrow{P} c^{-1}$. For any $\epsilon > 0$, let $\delta = \min\{\epsilon \frac{1}{2} c^2, \frac{1}{2} |c|\}$. If $|\eta_n - c| < \delta$, then $|\eta_n| > |c| - \delta > \frac{1}{2} |c|$, and so

$$|\eta_n^{-1} - c^{-1}| = \frac{|\eta_n - c|}{|\eta_n||c|} < \frac{\epsilon \frac{1}{2} c^2}{\frac{1}{2} |c| \cdot |c|} = \epsilon.$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Proof. We only give a proof of (3) and (4). By (2), it is sufficient to show that $\eta_n^{-1} \xrightarrow{P} c^{-1}$. For any $\epsilon > 0$, let $\delta = \min\{\epsilon \frac{1}{2} c^2, \frac{1}{2} |c|\}$. If $|\eta_n - c| < \delta$, then $|\eta_n| > |c| - \delta > \frac{1}{2} |c|$, and so

$$|\eta_n^{-1} - c^{-1}| = \frac{|\eta_n - c|}{|\eta_n||c|} < \frac{\epsilon \frac{1}{2} c^2}{\frac{1}{2} |c| \cdot |c|} = \epsilon.$$

It follows that

$$P(|\eta_n^{-1} - c^{-1}| \geq \epsilon) \leq P(|\eta_n - c| \geq \delta) \rightarrow 0.$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

For (4), it suffices to show that for any bounded continuous function $g(x, y)$ we have

$$Eg(\xi_n, \eta_n) \rightarrow Eg(\xi, c). \quad (*)$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

For (4), it suffices to show that for any bounded continuous function $g(x, y)$ we have

$$Eg(\xi_n, \eta_n) \rightarrow Eg(\xi, c). \quad (*)$$

If fact, choosing $g(x, y) = e^{it(x+y)}$ and $g(x, y) = e^{itxy}$ yields

$$Ee^{it(\xi_n + \eta_n)} \rightarrow Ee^{it(\xi + c)}, Ee^{it(\xi_n \eta_n)} \rightarrow Ee^{it(c\xi)},$$

respectively, which completes the proof by the inverse limit theorem.

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Now, suppose $g(x, y)$ is a continuous function with $|g(x, y)| \leq M$, then it is uniformly continuous in any bounded area. So for any given $\epsilon > 0$ and any $A > 0$ there exist a $\delta = \delta(A, \epsilon, g) > 0$ such that $|g(\xi_n, \eta_n) - g(\xi_n, c)| \leq \epsilon$ whenever $|\eta_n - c| \leq \delta$ and $|\xi_n| \leq A$.

Then

$$\begin{aligned} & |Eg(\xi_n, \eta_n) - Eg(\xi, c)| \\ \leq & |Eg(\xi_n, \eta_n) - Eg(\xi_n, c)| + |Eg(\xi_n, c) - Eg(\xi, c)| \\ \leq & E[|g(\xi_n, \eta_n) - g(\xi_n, c)|] + |Eg(\xi_n, c) - Eg(\xi, c)| \end{aligned}$$

Then

$$\begin{aligned} & |Eg(\xi_n, \eta_n) - Eg(\xi, c)| \\ & \leq |Eg(\xi_n, \eta_n) - Eg(\xi_n, c)| + |Eg(\xi_n, c) - Eg(\xi, c)| \\ & \leq E[|g(\xi_n, \eta_n) - g(\xi_n, c)|] + |Eg(\xi_n, c) - Eg(\xi, c)| \\ & \leq \epsilon + 2MP(|\eta_n - c| > \delta) \\ & \quad + |Eg(\xi_n, c) - Eg(\xi, c)| + 2MP(|\xi_n| > A). \end{aligned}$$

The second term will converge to zero because

$\eta_n \xrightarrow{P} c$. The third will also converge to zero

because $\xi_n \xrightarrow{d} \xi$ and $g(x, c)$ is a continuous function of x .

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

For the fourth term, we can choose A such that $\pm A$ is continuous points of the distribution function of ξ . Then $2MP(|\xi_n| > A)$ will converges to

$$2MP(|\xi| > A),$$

which can be smaller than the given $\epsilon > 0$ if A is large enough.

Finally, by the arbitrariness of ϵ , $(*)$ is proved.

Markov's inequality. Let ξ be a random variable defined on the probability space (Ω, \mathcal{F}, P) , $f(x)$ be a non-negatively monotonically non-decreasing function on $[0, \infty)$, then for any $x > 0$,

$$P(|\xi| > x) \leq \frac{Ef(|\xi|)}{f(x)}.$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

④ $\xi_n \xrightarrow{P} \xi$ if and only if

$$E \frac{|\xi_n - \xi|^2}{1 + |\xi_n - \xi|^2} \longrightarrow 0.$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

④ $\xi_n \xrightarrow{P} \xi$ if and only if

$$E \frac{|\xi_n - \xi|^2}{1 + |\xi_n - \xi|^2} \longrightarrow 0.$$

Proof. Let $F_n(x)$ denote the distribution function of $\xi_n - \xi$. Sufficiency: We have

$$P(|\xi_n - \xi| > \varepsilon) = \int_{|x| > \varepsilon} dF_n(x)$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

④ $\xi_n \xrightarrow{P} \xi$ if and only if

$$E \frac{|\xi_n - \xi|^2}{1 + |\xi_n - \xi|^2} \longrightarrow 0.$$

Proof. Let $F_n(x)$ denote the distribution function of $\xi_n - \xi$. Sufficiency: We have

$$\begin{aligned} P(|\xi_n - \xi| > \varepsilon) &= \int_{|x| > \varepsilon} dF_n(x) \\ &\leq \int_{|x| > \varepsilon} \frac{1 + \varepsilon^2}{\varepsilon^2} \frac{x^2}{1 + x^2} dF_n(x) \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

④ $\xi_n \xrightarrow{P} \xi$ if and only if

$$E \frac{|\xi_n - \xi|^2}{1 + |\xi_n - \xi|^2} \longrightarrow 0.$$

Proof. Let $F_n(x)$ denote the distribution function of $\xi_n - \xi$. Sufficiency: We have

$$\begin{aligned} P(|\xi_n - \xi| > \varepsilon) &= \int_{|x| > \varepsilon} dF_n(x) \\ &\leq \int_{|x| > \varepsilon} \frac{1 + \varepsilon^2}{\varepsilon^2} \frac{x^2}{1 + x^2} dF_n(x) \\ &\leq \frac{1 + \varepsilon^2}{\varepsilon^2} E \frac{|\xi_n - \xi|^2}{1 + |\xi_n - \xi|^2} \rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

That is $\xi_n \xrightarrow{P} \xi$.

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Necessity: For any $\varepsilon > 0$,

$$\begin{aligned} E \frac{|\xi_n - \xi|^2}{1 + |\xi_n - \xi|^2} &= \int_{-\infty}^{+\infty} \frac{x^2}{1 + x^2} dF_n(x) \\ &= \int_{|x| < \varepsilon} \frac{x^2}{1 + x^2} dF_n(x) + \int_{|x| \geq \varepsilon} \frac{x^2}{1 + x^2} dF_n(x) \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Necessity: For any $\varepsilon > 0$,

$$\begin{aligned} E \frac{|\xi_n - \xi|^2}{1 + |\xi_n - \xi|^2} &= \int_{-\infty}^{+\infty} \frac{x^2}{1 + x^2} dF_n(x) \\ &= \int_{|x| < \varepsilon} \frac{x^2}{1 + x^2} dF_n(x) + \int_{|x| \geq \varepsilon} \frac{x^2}{1 + x^2} dF_n(x) \\ &\leq \frac{\varepsilon^2}{1 + \varepsilon^2} + \int_{|x| \geq \varepsilon} dF_n(x) \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Necessity: For any $\varepsilon > 0$,

$$\begin{aligned} E \frac{|\xi_n - \xi|^2}{1 + |\xi_n - \xi|^2} &= \int_{-\infty}^{+\infty} \frac{x^2}{1 + x^2} dF_n(x) \\ &= \int_{|x| < \varepsilon} \frac{x^2}{1 + x^2} dF_n(x) + \int_{|x| \geq \varepsilon} \frac{x^2}{1 + x^2} dF_n(x) \\ &\leq \frac{\varepsilon^2}{1 + \varepsilon^2} + \int_{|x| \geq \varepsilon} dF_n(x) \\ &= \frac{\varepsilon^2}{1 + \varepsilon^2} + P(|\xi_n - \xi| \geq \varepsilon). \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Necessity: For any $\varepsilon > 0$,

$$\begin{aligned} E \frac{|\xi_n - \xi|^2}{1 + |\xi_n - \xi|^2} &= \int_{-\infty}^{+\infty} \frac{x^2}{1 + x^2} dF_n(x) \\ &= \int_{|x| < \varepsilon} \frac{x^2}{1 + x^2} dF_n(x) + \int_{|x| \geq \varepsilon} \frac{x^2}{1 + x^2} dF_n(x) \\ &\leq \frac{\varepsilon^2}{1 + \varepsilon^2} + \int_{|x| \geq \varepsilon} dF_n(x) \\ &= \frac{\varepsilon^2}{1 + \varepsilon^2} + P(|\xi_n - \xi| \geq \varepsilon). \end{aligned}$$

Since $\xi_n \xrightarrow{P} \xi$, first letting $n \rightarrow \infty$ and then letting $\varepsilon \rightarrow 0$ yield

$$E \frac{|\xi_n - \xi|^2}{1 + |\xi_n - \xi|^2} \longrightarrow 0.$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Let

$$\rho(\xi, \eta) = E \frac{|\xi - \eta|}{1 + |\xi - \eta|}.$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Let

$$\rho(\xi, \eta) = E \frac{|\xi - \eta|}{1 + |\xi - \eta|}.$$

Theorem

$\rho(\cdot, \cdot)$ satisfies

- $\rho(\xi, \eta) = 0$ if and only if $P(\xi = \eta) = 1$;
- $\rho(\xi, \eta) = \rho(\eta, \xi)$;
- $\rho(\xi, \tau) \leq \rho(\xi, \eta) + \rho(\eta, \tau)$.

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Let

$$\mathfrak{R} = \{\xi : \xi \text{ is a random variable on } (\Omega, \mathcal{F}, P)\}.$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Let

$$\mathfrak{R} = \{\xi : \xi \text{ is a random variable on } (\Omega, \mathcal{F}, P)\}.$$

Theorem

- (\mathfrak{R}, ρ) is a metric space;

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Let

$$\mathfrak{R} = \{\xi : \xi \text{ is a random variable on } (\Omega, \mathcal{F}, P)\}.$$

Theorem

- (\mathfrak{R}, ρ) is a metric space;
- $(\mathfrak{R}, \rho) = (\mathfrak{R}, \xrightarrow{P})$;

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Let

$$\mathfrak{R} = \{\xi : \xi \text{ is a random variable on } (\Omega, \mathcal{F}, P)\}.$$

Theorem

- (\mathfrak{R}, ρ) is a metric space;
- $(\mathfrak{R}, \rho) = (\mathfrak{R}, \xrightarrow{P})$;
- (\mathfrak{R}, ρ) is complete, i.e., $\xi_n - \xi_m \xrightarrow{P} 0$ as $n, m \rightarrow \infty$ if and only if there exists a random variable ξ such that $\xi_n \xrightarrow{P} \xi$.

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

5 Suppose $\xi_n \xrightarrow{P} \xi$, $P(|\xi_n| \leq \eta) = 1$ and $E\eta < \infty$. Then

$$E\xi_n \rightarrow E\xi.$$

5 Suppose $\xi_n \xrightarrow{P} \xi$, $P(|\xi_n| \leq \eta) = 1$ and $E\eta < \infty$. Then

$$E\xi_n \rightarrow E\xi.$$

Proof. First, we have $P(|\xi| \leq \eta) = 1$. In fact, for any $\epsilon > 0$,

$$\begin{aligned} P(|\xi| > \eta + \epsilon) &= P(|\xi| > \eta + \epsilon, |\xi_n - \xi| < \epsilon) \\ &\quad + P(|\xi| > \eta + \epsilon, |\xi_n - \xi| \geq \epsilon) \\ &\leq P(|\xi_n - \xi| \geq \epsilon) \rightarrow 0, \end{aligned}$$

which implies $P(|\xi| \leq \eta) = 1$.

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Now, for any $\epsilon > 0$ and $M > 0$, we have

$$\begin{aligned} |\xi_n - \xi| &\leq \epsilon + |\xi_n - \xi|I\{|\xi_n - \xi| \geq \epsilon\} \\ &\leq \epsilon + 2\eta I\{|\xi_n - \xi| \geq \epsilon\} \\ &\leq \epsilon + 2MI\{|\xi_n - \xi| \geq \epsilon\} + 2\eta I\{\eta \geq M\} \quad a.s \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Now, for any $\epsilon > 0$ and $M > 0$, we have

$$\begin{aligned} |\xi_n - \xi| &\leq \epsilon + |\xi_n - \xi| I\{|\xi_n - \xi| \geq \epsilon\} \\ &\leq \epsilon + 2\eta I\{|\xi_n - \xi| \geq \epsilon\} \\ &\leq \epsilon + 2MI\{|\xi_n - \xi| \geq \epsilon\} + 2\eta I\{\eta \geq M\} \quad a.s \end{aligned}$$

For any $\epsilon > 0$, choose $M > 0$ large enough such that

$$E\eta I\{\eta \geq M\} = \int_{y \geq M} y dF_\eta(y) < \epsilon/4.$$

Then choose N large enough such that

$$P(|\xi_n - \xi| \geq \epsilon) < \epsilon/(4M), \quad n \geq N.$$

4.2 Convergence in probability and weak law of large numbers

4.2.1 Convergence in probability

Then for $n \geq N$,

$$\begin{aligned} |E\xi_n - E\xi| &\leq E|\xi_n - \xi| \\ &\leq \epsilon + 2MP(|\xi_n - \xi| \geq \epsilon) + 2E\eta I\{\eta \geq M\} < 2\epsilon. \end{aligned}$$

4.2.2 Weak laws of large numbers

Consider the event A in random trial E . Suppose the probability of occurring A is p ($0 < p < 1$).

Now we experiment independently n times— n -fold Bernoulli trial. Let

$$\xi_i = \begin{cases} 1, & A \text{ occurs at the } i\text{-th trial,} \\ 0, & A \text{ does not occur at the } i\text{-th trial,} \end{cases}$$

$1 \leq i \leq n$. Then $P(\xi_i = 1) = p$,

$P(\xi_i = 0) = 1 - p$. Let $S_n = \sum_{i=1}^n \xi_i$. Then

$$\frac{S_n}{n} = F_n(A) \text{ --- the frequency of } A.$$

What does

$$\frac{S_n}{n} = F_n(A) \approx P(A) = p$$

mean?

What does

$$\frac{S_n}{n} = F_n(A) \approx P(A) = p$$

mean?

For any $\varepsilon > 0$ we can not expect that

$|S_n/n - p| \leq \varepsilon$ holds for all the trials even if n is big enough.

What does

$$\frac{S_n}{n} = F_n(A) \approx P(A) = p$$

mean?

For any $\varepsilon > 0$ we can not expect that

$|S_n/n - p| \leq \varepsilon$ holds for all the trials even if n is big enough.

It is nature to hope that the probability to appear $\{|S_n/n - p| \geq \varepsilon\}$ could be as smaller as possible when n is large enough.

Theorem 4 (Bernoulli) Let $\{\xi_n, n \geq 1\}$ be a sequence of independent and identically distributed random variables with $P(\xi_n = 1) = p$, $P(\xi_n = 0) = 1 - p$, $0 < p < 1$. Put $S_n = \sum_{i=1}^n \xi_i$. Then we have

$$\frac{S_n}{n} \xrightarrow{P} p,$$

i.e., for any $\epsilon > 0$ and $\delta > 0$, there is a $N = N(\epsilon, \delta)$ such that

$$P\left(\left|\frac{S_n}{n} - p\right| \geq \epsilon\right) < \delta, \quad \text{for all } n \geq N.$$

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Theorem 5(Chebyshev) Let $\{\xi_n, n \geq 1\}$ be a sequence of **independent** (or **pairwise correlated**) random variables defined on the probability space (Ω, \mathcal{F}, P) with $E\xi_n = \mu_n$ and $Var\xi_n = \sigma_n^2$. If $\sum_{k=1}^n \sigma_k^2 / n^2 \longrightarrow 0$, then $\{\xi_n, n \geq 1\}$ obeys the weak law of large numbers, i.e.,

$$\frac{1}{n} \sum_{k=1}^n \xi_k - \frac{1}{n} \sum_{k=1}^n \mu_k \xrightarrow{P} 0.$$

Using the Chebyshev inequality, we have

$$\begin{aligned} & P\left(\left|\frac{1}{n} \sum_{k=1}^n (\xi_k - \mu_k)\right| \geq \varepsilon\right) \\ & \leq P\left(\left|\frac{1}{n} \sum_{k=1}^n \xi_k - E\frac{1}{n} \sum_{k=1}^n \xi_k\right| \geq \varepsilon\right) \\ & \leq \frac{1}{\varepsilon^2} \text{Var}\left(\frac{1}{n} \sum_{k=1}^n \xi_k\right) \\ & = \frac{1}{\varepsilon^2 n^2} \sum_{k=1}^n \sigma_k^2 \longrightarrow 0 ; \text{ as } n \longrightarrow \infty. \end{aligned}$$

The proof is complete.

Example 8. Suppose that $\xi_k \sim \begin{pmatrix} k^s & -k^s \\ 0.5 & 0.5 \end{pmatrix}$,

where $s < 1/2$ is a constant, and $\{\xi_k, k \geq 1\}$ is indept.. Prove that $\{\xi_k, k \geq 1\}$ obeys the weak LLN.

Proof.

Example 8. Suppose that $\xi_k \sim \begin{pmatrix} k^s & -k^s \\ 0.5 & 0.5 \end{pmatrix}$,

where $s < 1/2$ is a constant, and $\{\xi_k, k \geq 1\}$ is indept.. Prove that $\{\xi_k, k \geq 1\}$ obeys the weak LLN.

Proof. We have $E\xi_k = 0$, $Var\xi_k = k^{2s}$. When $s < 1/2$,

$$\frac{1}{n^2} \sum_{k=1}^n Var\xi_k = \frac{1}{n^2} \sum_{k=1}^n k^{2s} < \frac{1}{n^2} \sum_{k=1}^n n^{2s} = n^{2s-1} \longrightarrow 0.$$

In addition, $\{\xi_k, k \geq 1\}$ is also independent,

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Example 8. Suppose that $\xi_k \sim \begin{pmatrix} k^s & -k^s \\ 0.5 & 0.5 \end{pmatrix}$,

where $s < 1/2$ is a constant, and $\{\xi_k, k \geq 1\}$ is indept.. Prove that $\{\xi_k, k \geq 1\}$ obeys the weak LLN.

Proof. We have $E\xi_k = 0$, $Var\xi_k = k^{2s}$. When $s < 1/2$,

$$\frac{1}{n^2} \sum_{k=1}^n Var\xi_k = \frac{1}{n^2} \sum_{k=1}^n k^{2s} < \frac{1}{n^2} \sum_{k=1}^n n^{2s} = n^{2s-1} \longrightarrow 0.$$

In addition, $\{\xi_k, k \geq 1\}$ is also independent, so $\{\xi_k, k \geq 1\}$ obeys the Chebyshev LLN, i.e.,

$$\frac{1}{n} \sum_{k=1}^n \xi_k \xrightarrow{P} 0.$$

Theorem 6 (Khinchine) Let $\{\xi_n, n \geq 1\}$ be a sequence of independent and identically distributed random variables defined (Ω, \mathcal{F}, P) with $E|\xi_1| < \infty$. Let $E\xi_1 = \mu$, $S_n = \sum_{k=1}^n \xi_k$. Then $\{\xi_n, n \geq 1\}$ obeys the weak LLN, i.e.,

$$\frac{S_n}{n} \xrightarrow{P} \mu \quad \text{as } n \rightarrow \infty.$$

Proof. Let $f(t)$ and $f_n(t)$ be the c.f.s of ξ_1 and S_n/n respectively.

Proof. Let $f(t)$ and $f_n(t)$ be the c.f.s of ξ_1 and S_n/n respectively. Since $\{\xi_n, n \geq 1\}$ is i.i.d., we have $f_n(t) = (f(t/n))^n$. Moreover, from the Taylor expansion formula, we have

$$f(t) = 1 + i\mu t + o(t) \quad \text{as } t \longrightarrow 0,$$

since $E\xi_1 = \mu$.

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Proof. Let $f(t)$ and $f_n(t)$ be the c.f.s of ξ_1 and S_n/n respectively. Since $\{\xi_n, n \geq 1\}$ is i.i.d., we have $f_n(t) = (f(t/n))^n$. Moreover, from the Taylor expansion formula, we have

$$f(t) = 1 + i\mu t + o(t) \quad \text{as } t \longrightarrow 0,$$

since $E\xi_1 = \mu$. For every $t \in \mathbf{R}$,

$$f(t/n) = 1 + i\frac{\mu t}{n} + o\left(\frac{1}{n}\right) \quad \text{as } n \rightarrow \infty,$$

$$f_n(t) = \left(1 + i\frac{\mu t}{n} + o\left(\frac{1}{n}\right)\right)^n \rightarrow e^{i\mu t}.$$

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Proof. Let $f(t)$ and $f_n(t)$ be the c.f.s of ξ_1 and S_n/n respectively. Since $\{\xi_n, n \geq 1\}$ is i.i.d., we have $f_n(t) = (f(t/n))^n$. Moreover, from the Taylor expansion formula, we have

$$f(t) = 1 + i\mu t + o(t) \quad \text{as } t \longrightarrow 0,$$

since $E\xi_1 = \mu$. For every $t \in \mathbf{R}$,

$$f(t/n) = 1 + i\frac{\mu t}{n} + o\left(\frac{1}{n}\right) \quad \text{as } n \rightarrow \infty,$$

$$f_n(t) = \left(1 + i\frac{\mu t}{n} + o\left(\frac{1}{n}\right)\right)^n \rightarrow e^{i\mu t}.$$

By the inverse limit theorem in we know that $S_n/n \xrightarrow{d} \mu$. So, we have $S_n/n \xrightarrow{P} \mu$. The proof is complete.

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

Proof (2): For $M > 0$, let $\eta_k = \xi_k I\{|\xi_k| \leq M\}$,
 $\zeta_k = \xi_k I\{|\xi_k| > M\}$. Then

$$\begin{aligned} & P\left(\frac{|\sum_{k=1}^n (\eta_k - E\eta_k)|}{n} \geq \epsilon/2\right) \\ & \leq \frac{4}{\epsilon^2 n^2} \sum_{k=1}^n \text{Var}(\eta_k) \\ & \leq \sum_{k=1}^n \frac{4}{\epsilon^2 n^2} E[\xi_k^2 I\{|\xi_k| \leq M\}] \leq \frac{4M^2}{\epsilon^2 n}; \end{aligned}$$

4.2 Convergence in probability and weak law of large numbers

4.2.2 Weak laws of large numbers

$$\begin{aligned} P\left(\frac{|\sum_{k=1}^n(\zeta_k - E\zeta_k)|}{n} \geq \epsilon/2\right) &\leq \frac{2}{\epsilon n} E\left|\sum_{k=1}^n(\zeta_k - E\zeta_k)\right| \\ &\leq \frac{2}{\epsilon n} \sum_{k=1}^n E|\zeta_k - E\zeta_k| \leq 2\frac{2}{\epsilon n} \sum_{k=1}^n E|\zeta_k| \leq \frac{4}{\epsilon} E[|\xi_1| I\{|\xi_1| > M\}]. \end{aligned}$$

Hence,

$$\begin{aligned} P\left(\frac{|\sum_{k=1}^n(\xi_k - E\xi_k)|}{n} \geq \epsilon\right) \\ \leq \frac{4M^2}{\epsilon^2 n} + \frac{4}{\epsilon} E[|\xi_1| I\{|\xi_1| > M\}] \\ \rightarrow 0 \quad \text{as } n \rightarrow \infty, \text{ and then } M \rightarrow \infty. \end{aligned}$$

Corollary Let $\{\xi_n, n \geq 1\}$ be a sequence of **pairwise independent** and **identically distributed** random variables defined (Ω, \mathcal{F}, P) with $E|\xi_1| < \infty$. Let $E\xi_1 = \mu$, $S_n = \sum_{k=1}^n \xi_k$. Then $\{\xi_n, n \geq 1\}$ obeys the weak LLN, i.e.,

$$\frac{S_n}{n} \xrightarrow{P} \mu \quad \text{as } n \rightarrow \infty.$$

Definition In general, suppose $\{\xi_n, n \geq 1\}$ is a sequence of random variables defined on the probability space (Ω, \mathcal{F}, P) . If there exist constant sequences $\{a_n, n \geq 1\}$ and $\{b_n, n \geq 1\}$ such that

$$\frac{1}{a_n} \sum_{k=1}^n \xi_k - b_n \xrightarrow{P} 0 \quad \text{as } n \rightarrow \infty.$$

Then $\{\xi_n\}$ will be said to obey the weak law of large numbers, in short $\{\xi_n, n \geq 1\}$ obeys LLN.

The applications of LLN

Example

Let $\{\xi_k, k \geq 1\}$ be a sequence of i.i.d. random variables with $E\xi_k = \mu$ and $Var\xi_k = \sigma^2$. Let

$$\bar{\xi}_n = \frac{1}{n} \sum_{k=1}^n \xi_k, \quad \hat{\sigma}_n^2 = \frac{1}{n} \sum_{k=1}^n (\xi_k - \bar{\xi}_n)^2.$$

Prove that $\hat{\sigma}_n^2 \xrightarrow{P} \sigma^2$ and find the asymptotic distribution of $\sqrt{n} \frac{\bar{\xi}_n - \mu}{\hat{\sigma}_n}$.

Proof.

$$\begin{aligned}\widehat{\sigma}_n^2 &= \frac{1}{n} \sum_{k=1}^n (\xi_k - \bar{\xi}_n)^2 \\ &= \frac{1}{n} \sum_{k=1}^n ((\xi_k - \mu) - (\bar{\xi}_n - \mu))^2 \\ &= \frac{1}{n} \sum_{k=1}^n (\xi_k - \mu)^2 - (\bar{\xi}_n - \mu)^2.\end{aligned}$$

By the Khinchine weak LLN, we have $\bar{\xi}_n \xrightarrow{P} \mu$.

Thus $\bar{\xi}_n - \mu \xrightarrow{P} 0$.

4.2 Convergence in probability and weak law of large numbers

The applications of LLN

Moreover, since $\{(\xi_k - \mu)^2, k \geq 1\}$ is i.i.d. and $E(\xi_k - \mu)^2 = Var\xi_k = \sigma^2$, $\{(\xi_k - \mu)^2, k \geq 1\}$ also obeys the Khinchine weak LLN, i.e.

$\sum_{k=1}^n (\xi_k - \mu)^2 / n \xrightarrow{P} \sigma^2$. Hence $\hat{\sigma}_n^2 \xrightarrow{P} \sigma^2$.

4.2 Convergence in probability and weak law of large numbers

The applications of LLN

Moreover, since $\{(\xi_k - \mu)^2, k \geq 1\}$ is i.i.d. and $E(\xi_k - \mu)^2 = \text{Var}\xi_k = \sigma^2$, $\{(\xi_k - \mu)^2, k \geq 1\}$ also obeys the Khinchine weak LLN, i.e.

$\sum_{k=1}^n (\xi_k - \mu)^2 / n \xrightarrow{P} \sigma^2$. Hence $\hat{\sigma}_n^2 \xrightarrow{P} \sigma^2$.

By the Lindeberg-Lévy central limit theorem,

$$\sqrt{n} \frac{\bar{\xi}_n - \mu}{\sigma} = \frac{\sum_{k=1}^n (\xi_k - \mu)}{\sqrt{n\sigma^2}} \xrightarrow{d} N(0, 1).$$

Hence

$$\sqrt{n} \frac{\bar{\xi}_n - \mu}{\hat{\sigma}_n} = \frac{\sigma}{\hat{\sigma}_n} \cdot \sqrt{n} \frac{\bar{\xi}_n - \mu}{\sigma} \xrightarrow{d} N(0, 1).$$

Example Prove that for any $q > p > 0$,

$$\lim_{n \rightarrow \infty} \int_0^1 \cdots \int_0^1 \frac{x_1^q + \cdots + x_n^q}{x_1^p + \cdots + x_n^p} dx_1 \cdots dx_n = \frac{p+1}{q+1}.$$

Example Prove that for any $q > p > 0$,

$$\lim_{n \rightarrow \infty} \int_0^1 \cdots \int_0^1 \frac{x_1^q + \cdots + x_n^q}{x_1^p + \cdots + x_n^p} dx_1 \cdots dx_n = \frac{p+1}{q+1}.$$

Proof. Let $\{\xi_i\}$ i.i.d. $\sim U(0, 1)$, and let

$$\eta_n = \frac{\xi_1^q + \cdots + \xi_n^q}{\xi_1^p + \cdots + \xi_n^p}.$$

Then $0 \leq \eta_n \leq 1$ and

$$\int_0^1 \cdots \int_0^1 \frac{x_1^q + \cdots + x_n^q}{x_1^p + \cdots + x_n^p} dx_1 \cdots dx_n = E\eta_n.$$

4.2 Convergence in probability and weak law of large numbers

The applications of LLN

On the other hand, by WLLN,

$$\frac{1}{n} \sum_{k=1}^n \xi_k^q \xrightarrow{P} E\xi_1^q = \frac{1}{q+1}$$
$$\frac{1}{n} \sum_{k=1}^n \xi_k^p \xrightarrow{P} E\xi_1^p = \frac{1}{p+1}.$$

So,

$$\eta_n \xrightarrow{P} \frac{E\xi_1^q}{E\xi_1^p} = \frac{p+1}{q+1}.$$

Hence

$$\int_0^1 \cdots \int_0^1 \frac{x_1^q + \cdots + x_n^q}{x_1^p + \cdots + x_n^p} dx_1 \cdots dx_n = E\eta_n \rightarrow \frac{p+1}{q+1}.$$

Convergence in mean of order r :

Definition 3 Let $r > 0$, ξ and $\{\xi_n, n \geq 1\}$ be random variables defined on (Ω, \mathcal{F}, P) with $E|\xi|^r < \infty$ and $E|\xi_n|^r < \infty$. If

$$E|\xi_n - \xi|^r \longrightarrow 0,$$

then we say that $\{\xi_n, n \geq 1\}$ converges in mean of order r to ξ , denoted by $\xi_n \xrightarrow{L_r} \xi$.

Convergence in mean of order r :

Definition 3 Let $r > 0$, ξ and $\{\xi_n, n \geq 1\}$ be random variables defined on (Ω, \mathcal{F}, P) with $E|\xi|^r < \infty$ and $E|\xi_n|^r < \infty$. If

$$E|\xi_n - \xi|^r \longrightarrow 0,$$

then we say that $\{\xi_n, n \geq 1\}$ converges in mean of order r to ξ , denoted by $\xi_n \xrightarrow{L_r} \xi$.

$$\xi_n \xrightarrow{L_r} \xi \Rightarrow \xi_n \xrightarrow{P} \xi.$$

4.2 Convergence in probability and weak law of large numbers

Convergence in mean of order r

Convergence in mean of order r :

Definition 3 Let $r > 0$, ξ and $\{\xi_n, n \geq 1\}$ be random variables defined on (Ω, \mathcal{F}, P) with $E|\xi|^r < \infty$ and $E|\xi_n|^r < \infty$. If

$$E|\xi_n - \xi|^r \longrightarrow 0,$$

then we say that $\{\xi_n, n \geq 1\}$ converges in mean of order r to ξ , denoted by $\xi_n \xrightarrow{L_r} \xi$.

$$\xi_n \xrightarrow{L_r} \xi \Rightarrow \xi_n \xrightarrow{P} \xi.$$

$$\xi_n \xrightarrow{L_r} \xi \not\Leftarrow \xi_n \xrightarrow{P} \xi.$$

Example 5. Define ξ_n by

$$P(\xi_n = n) = 1/\log(n+3),$$

$P(\xi_n = 0) = 1 - 1/\log(n+3)$, $n = 1, 2, \dots$. It is easy to know $\xi_n \xrightarrow{P} 0$, but for any $0 < r < \infty$,

$$E|\xi_n|^r = \frac{n^r}{\log(n+3)} \longrightarrow \infty.$$

That is, $\xi_n \xrightarrow{L_r} 0$ does not hold true.