

# REAL ANALYSIS

## LECTURE NOTES

ABSTRACT. The Notes indicate what we do in the lectures, but are not a complete replacement of the book and lectures. The text is from two books of *Real Analysis*:

- [1] Xingwei Zhou & Wenchang Sun: Real Variable Analysis, the third edition, Science Press, 2014.
- [2] E. Stein & R. Shakarchi: Real Analysis, Princeton University Press, 2005.

### 1. SOME OTHER PROPERTIES OF MEASURABLE SETS AND FUNCTIONS

**Proposition 1.1.** *Let  $E_i$  be measurable set of  $\mathbb{R}^{n_i}$ ,  $i = 1, 2$ . Then  $E = E_1 \times E_2$  are measurable set of  $\mathbb{R}^n$  with  $n = n_1 + n_2$ . Moreover,*

$$(1.1) \quad m(E) = m(E_1)m(E_2).^1$$

*Proof.* Since  $E_1$  and  $E_2$  are measurable, we have

$$E_i = \cup_{j \geq 1} E_{i,j}, \quad i = 1, 2,$$

where  $E_{i,j}$  is either closed or measure zero. By a cut-off if necessary, we can further require that  $m(E_{i,j}) < \infty$  for all  $i, j$ . It can be verified that

$$E = E_1 \times E_2 = (\cup_{j \geq 1} E_{1,j}) \times (\cup_{k \geq 1} E_{2,k}) = \cup_{k,j=1}^{\infty} E_{1,j} \times E_{2,k}.$$

Then it suffices to show that

- (1) if  $H$  and  $F$  are both closed, then  $H \times F$  is measurable;
- (2) if  $m(H) = 0$  and  $m(F) < \infty$ , then  $m(H \times F) = 0$ .

Note that (1) follows by the closedness of  $H \times F$ . For (2), given  $\varepsilon > 0$ , we choose closed cubes  $\{Q_j^1\}$  and  $\{Q_k^2\}$  such that

$$H \subset \cup_{j \geq 1} Q_j^1, \quad F \subset \cup_{k \geq 1} Q_k^2, \quad \sum_{j=1}^{\infty} |Q_j^1| \leq \varepsilon \quad \text{and} \quad \sum_{k=1}^{\infty} |Q_k^2| \leq m(F) + \varepsilon.$$

---

<sup>1</sup>We understand this equality by the convention  $0 \cdot \infty = 0$ .

Consequently  $H \times F \subset \cup_{j,k \geq 1} Q_j^1 \times Q_k^2$  and

$$\sum_{j,k=1}^{\infty} |Q_j^1 \times Q_k^2| = \sum_{j,k=1}^{\infty} |Q_j^1| |Q_k^2| \leq \varepsilon(m(F) + \varepsilon),$$

where we used the fact  $Q_j^1 \times Q_k^2$  are rectangles in  $\mathbb{R}^n$  and so  $|Q_j^1 \times Q_k^2| = |Q_j^1| |Q_k^2|$ . Therefore  $m_*(E) = 0$ . Hence (2) follows.

We next show (1.1). Observe the following

(a) If  $m(H) = 0$ , then  $m(H \times F) = 0$ .

*Proof:* This follows by (1) in previous part and the cut-off argument. Namely using  $F = \cup_{j \geq 1} F_j$  with  $F_j = F \cap \{j-1 \leq |x| < j\}$  we see that  $m(H \times F) = \sum_{j \geq 1} m(H \times F_j) = 0$ .

(b) If  $\mathcal{O}_i$  are open sets of  $\mathbb{R}^{n_i}$ ,  $i = 1, 2$ , then  $m(\mathcal{O}_1 \times \mathcal{O}_2) = m(\mathcal{O}_1)m(\mathcal{O}_2)$ .

*Proof:* There are almost disjoint closed cubes  $\{Q_j^1\}$  and  $\{Q_j^2\}$  such that  $\mathcal{O}_i = \cup_{j=1}^{\infty} Q_j^i$ ,  $i = 1, 2$ . Then

$$\mathcal{O}_1 \times \mathcal{O}_2 = \cup_{j,k \geq 1} Q_j^1 \times Q_k^2.$$

Consequently

$$m(\mathcal{O}_1 \times \mathcal{O}_2) = \sum_{j,k=1}^{\infty} |Q_j^1 \times Q_k^2| = \sum_{j,k=1}^{\infty} |Q_j^1| |Q_k^2| = m(\mathcal{O}_1)m(\mathcal{O}_2).$$

(c) If  $G_i$  are  $G_\delta$  sets in  $\mathbb{R}^{n_i}$  with finite measure,  $i = 1, 2$ , then

$$m(G_1 \times G_2) = m(G_1)m(G_2).$$

*Proof:* Suppose  $G_i = \cap_{j=1}^{\infty} \mathcal{O}_j^i$ . Then  $(\cap_{j=1}^N \mathcal{O}_j^1) \times (\cap_{k=1}^N \mathcal{O}_k^2) \searrow G_1 \times G_2$ . Since  $G_1$  and  $G_2$  are of finite measure, we deduce

$$\begin{aligned} m(G_1 \times G_2) &= \lim_{N \rightarrow \infty} m((\cap_{j=1}^N \mathcal{O}_j^1) \times (\cap_{k=1}^N \mathcal{O}_k^2)) \\ &= \lim_{N \rightarrow \infty} m((\cap_{j=1}^N \mathcal{O}_j^1)) m((\cap_{k=1}^N \mathcal{O}_k^2)) \quad [by \text{ Observation (b)}] \\ &= m(G_1)m(G_2). \end{aligned}$$

Suppose  $E_i$  are of finite measure,  $i = 1, 2$ . Recall that  $E_i = G_i \setminus Z_i$  with  $G_\delta$  set  $G_i$  and measure zero set  $Z_i$ . Note that

$$(G_1 \times G_2) \setminus (E_1 \times E_2) = ((G_1 \setminus E_1) \times G_2) \cup (G_1 \times (G_2 \setminus E_2)) = (Z_1 \times G_2) \cup (G_1 \times Z_2).$$

Hence, by virtue of Observation (a),

$$m((G_1 \times G_2) \setminus (E_1 \times E_2)) \leq m(Z_1 \times G_2) + m(G_1 \times Z_2) = 0.$$

Consequently, by Observation (c), we infer that

$$m(E_1 \times E_2) = m(G_1 \times G_2) = m(G_1)m(G_2) = m(E_1)m(E_2).$$

In general, we consider  $E_i = \cup_{k=1}^{\infty} E_{i,k}$  with  $E_{i,k} = E_i \cap \{|x| \leq k\}$ ,  $i = 1, 2$ . Since  $E_{i,k} \nearrow E_i$  and  $E_{1,k} \times E_{2,k} \nearrow E_1 \times E_2$ , one concludes

$$m(E_1 \times E_2) = \lim_{k \rightarrow \infty} m(E_{1,k} \times E_{2,k}) = \lim_{k \rightarrow \infty} m(E_{1,k})m(E_{2,k}) = m(E_1)m(E_2).$$

□

**Proposition 1.2.** *Let  $E$  be a measurable set of  $\mathbb{R}^n$ . Then*

$$E^- = \{(x, y) \in \mathbb{R}^{2n} : x - y \in E\}$$

*is a measurable set of  $\mathbb{R}^{2n}$ .*

*Proof.* We have the following observation:

- (i) if  $H$  is open, then  $H^-$  are open;
- (ii) if  $H$  is closed, then  $H^-$  are closed;
- (iii) if  $R = \prod_{i=1}^n \langle a_i, b_i \rangle$ , then  $R^- = \{(x_1, \dots, x_n, y_1, \dots, y_n) \in \mathbb{R}^{2n} : x_i - y_i \in \langle a_i, b_i \rangle\}$ .  
Setting  $S_i = \{(x, y) \in \mathbb{R}^2 : x - y \in \langle a_i, b_i \rangle\}$ , then

$$R^- \cap [-\ell, \ell]^{2n} = \prod_{i=1}^n (S_i \cap [-\ell, \ell]^2).$$

By Proposition 1.1, we obtain

$$m(R^- \cap [-\ell, \ell]^{2n}) = \prod_{i=1}^n m(S_i \cap [-\ell, \ell]^2) \leq (2\ell)^n \prod_{i=1}^n (b_i - a_i) = (2\ell)^n |R|.$$

Recall that there are open sets  $\mathcal{O}_j$  and closed sets  $F_j$  such that

$$F_j \subset E \subset \mathcal{O}_j, \quad m(E \setminus F_j) \leq \frac{1}{j} \text{ and } m(\mathcal{O}_j \setminus E) \leq \frac{1}{j}.$$

In view of (i) and (ii),  $F_j^-$  and  $\mathcal{O}_j^-$  are measurable.

Note that  $F_j^- \subset E^- \subset \mathcal{O}_j^-$ . Let  $G = \cap_{j \geq 1} \mathcal{O}_j^-$ . Then  $G \setminus E^- \subset \mathcal{O}_j^- \setminus F_j^-$ ,  $\forall j$ . We claim, for any  $\ell > 0$ ,

$$(1.2) \quad m((\mathcal{O}_j^- \setminus F_j^-) \cap [-\ell, \ell]^{2n}) \rightarrow 0 \text{ as } j \rightarrow \infty.$$

Once this is done, we conclude that  $G \setminus E^-$  is measure zero, and so  $E^-$  is measurable.

We shall prove a general result:

if  $H \subset \mathbb{R}^n$  with  $m_*(H) < \infty$ , then  $m_*(H^- \cap [-\ell, \ell]^{2n}) \leq (2\ell)^n m_*(H)$  for each  $\ell > 0$ .

Then (1.2) is a consequence of the above statement as

$$\mathcal{O}_j^- \setminus F_j^- = (\mathcal{O}_j \setminus F_j)^- \implies m((\mathcal{O}_j^- \setminus F_j^-) \cap [\ell, \ell]^{2n}) \leq (2\ell)^n m(\mathcal{O}_j \setminus F_j) \rightarrow 0, \forall \ell > 0.$$

Take open sets  $U_k \supset H$  so that  $m_*(H) = m(G)$  where  $G = \bigcap_{k \geq 1} U_k$ . Let  $G_j = \bigcap_{k=1}^j U_k$ . It is open. Hence  $G_j = \bigcup_{k \geq 1} Q_{j,k}$  for a sequence of almost disjoint closed cubes  $\{Q_{j,k}\}_{k \geq 1}$ . By Observation (iii),

$$m(Q_{j,k}^- \cap [-\ell, \ell]^{2n}) \leq (2\ell)^n |Q_{j,k}| \implies m(G_j^- \cap [-\ell, \ell]^{2n}) \leq (2\ell)^n m(G_j).$$

Since  $G_j^- \searrow G^-$ , we see that by sending  $j \rightarrow \infty$ ,

$$\begin{aligned} m_*(H^- \cap [-\ell, \ell]^{2n}) &\leq m(G^- \cap [-\ell, \ell]^{2n}) \\ &= \lim_{j \rightarrow \infty} m(G_j^- \cap [-\ell, \ell]^{2n}) \\ &\leq (2\ell)^n \lim_{j \rightarrow \infty} m(G_j) \\ &= (2\ell)^n m(G) \\ &= (2\ell)^n m_*(H). \end{aligned}$$

□

**Proposition 1.3.** *Let  $E_i$  be measurable set of  $\mathbb{R}^{n_i}$  and  $f_i$  be (a.e. finite-valued) measurable function on  $E_i$ ,  $i = 1, 2$ . Then*

- (i)  $E = E_1 \times E_2$  is a measurable set of  $\mathbb{R}^n$  with  $n = n_1 + n_2$ ;
- (ii)  $F(x_1, x_2) = f_1(x_1)f_2(x_2)$  is a measurable function on  $E$ .

*Proof.* For part (i), one sees Proposition 1.1.

We show part (ii). Select two sequences of simple functions  $\{f_{1,k}\}_{k \geq 1}$  and  $\{f_{2,j}\}_{j \geq 1}$  such that, for  $i = 1, 2$ ,

$$(1.3) \quad f_i(x) = \lim_{j \rightarrow \infty} f_{i,j}(x) \quad \text{for all } x \in E_i.$$

By part (i),  $F_j(x_1, x_2) := f_{1,j}(x_1)f_{2,j}(x_2)$  are simple functions. It follows by (1.3)

$$F_j(x_1, x_2) \rightarrow F(x_1, x_2) \quad \text{for all } (x_1, x_2) \in E.$$

Consequently  $F(x_1, x_2)$  is measurable.

□

**Proposition 1.4.** *If  $f$  is a finite-valued measurable function on  $E$ . Then  $f(x - y)$  is a measurable function on  $E^- = \{(x, y) \in \mathbb{R}^{2n} : x - y \in E\}$ .*

*Proof.* Exercise.

□

Convergence in measure is preserved under some operations.

**Proposition 1.5.** *Let  $f_k, g_k, f, g$  be measurable functions on  $E \subset \mathbb{R}^n$ . Suppose that  $f_k \rightarrow f$  in measure and  $g_k \rightarrow g$  in measure. Then*

- (i)  $f_k \pm g_k \rightarrow f \pm g$  in measure;
- (ii)  $|f_k| \rightarrow |f|$  in measure;
- (iii)  $\min\{f_k, g_k\} \rightarrow \min\{f, g\}$  in measure, and  $\max\{f_k, g_k\} \rightarrow \max\{f, g\}$  in measure;
- (iv) when  $m(E) < \infty$ ,  $f_k g_k \rightarrow fg$  in measure;

*Proof.* Exercise.

□