

4.3 Almost sure convergence and strong laws of large numbers

4.3.1 Convergence in probability

4.3 Almost sure convergence and strong laws of large numbers

4.3.1 Almost sure convergence

4.3 Almost sure convergence and strong laws of large numbers

4.3.1 Convergence in probability

4.3 Almost sure convergence and strong laws of large numbers

4.3.1 Almost sure convergence

Definition 1 Suppose that ξ and $\{\xi_n, n \geq 1\}$ are random variables defined on a common probability space (Ω, \mathcal{F}, P) .

If there exists a $\Omega_0 \in \mathcal{F}$ such that $P(\Omega_0) = 0$ and for any $\omega \in \Omega \setminus \Omega_0$, $\xi_n(\omega) \rightarrow \xi(\omega)$, $(n \rightarrow \infty)$, then we say that ξ_n converges **with probability one** or **almost surely** to ξ , denoted by $\xi_n \rightarrow \xi$ a.s.

4.3 Almost sure convergence and strong laws of large numbers

4.3.1 Convergence in probability

Theorem 1 Suppose that ξ and $\{\xi_n, n \geq 1\}$ are random variables defined on (Ω, \mathcal{F}, P) .

$\xi_n(\omega) \rightarrow \xi(\omega)$ a.s. iff for any $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} P(\sup_{k \geq n} |\xi_k - \xi| \geq \epsilon) = 0$$

$$\text{i.e., } \lim_{n \rightarrow \infty} P(\bigcup_{k \geq n} (|\xi_k - \xi| \geq \epsilon)) = 0.$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.1 Convergence in probability

Corollary 1

$$\xi_n \rightarrow \xi \text{ a.s.} \Rightarrow \xi_n \xrightarrow{P} \xi.$$

Corollary 1

$$\xi_n \rightarrow \xi \text{ a.s.} \Rightarrow \xi_n \xrightarrow{P} \xi.$$

Corollary 2. If for any $\epsilon > 0$,

$\sum_{n=1}^{\infty} P(|\xi_n - \xi| \geq \epsilon) < \infty$, then

$$\xi_n \rightarrow \xi \text{ a.s.}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.1 Convergence in probability

Corollary 1

$$\xi_n \rightarrow \xi \text{ a.s.} \Rightarrow \xi_n \xrightarrow{P} \xi.$$

Corollary 2. If for any $\epsilon > 0$,
 $\sum_{n=1}^{\infty} P(|\xi_n - \xi| \geq \epsilon) < \infty$, then

$$\xi_n \rightarrow \xi \text{ a.s.}$$

Proof. Note that

$$\begin{aligned} P(|\xi_n - \xi| \geq \epsilon) &\leq P\left(\bigcup_{k \geq n} (|\xi_k - \xi| \geq \epsilon)\right) \\ &\leq \sum_{k=n}^{\infty} P(|\xi_k - \xi| \geq \epsilon). \end{aligned}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.1 Convergence in probability

Proof of Theorem 1. For any $\epsilon > 0$, let

$A_n^\epsilon = \{|\xi_n - \xi| \geq \epsilon\}$ and $A^\epsilon = \bigcap_{n=1}^{\infty} \bigcup_{k \geq n} A_k^\epsilon$. Then

$\xi_n(\omega) \not\rightarrow \xi(\omega)$ is equivalent to that, there is an $\epsilon_0 > 0$ such that for any N there is a $n \geq N$ for which $|\xi_n(\omega) - \xi(\omega)| \geq \epsilon_0$. This is also equivalent to that, there is an m such that for any n there is a $k \geq n$ for which $|\xi_k(\omega) - \xi(\omega)| \geq 1/m$. So

$$\{\xi_n \not\rightarrow \xi\} = \bigcup_{\epsilon > 0} A^\epsilon = \bigcup_{m=1}^{\infty} A^{\frac{1}{m}}.$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.1 Convergence in probability

Proof of Theorem 1. For any $\epsilon > 0$, let

$A_n^\epsilon = \{|\xi_n - \xi| \geq \epsilon\}$ and $A^\epsilon = \bigcap_{n=1}^{\infty} \bigcup_{k \geq n} A_k^\epsilon$. Then

$\xi_n(\omega) \not\rightarrow \xi(\omega)$ is equivalent to that, there is an $\epsilon_0 > 0$ such that for any N there is a $n \geq N$ for which $|\xi_n(\omega) - \xi(\omega)| \geq \epsilon_0$. This is also equivalent to that, there is an m such that for any n there is a $k \geq n$ for which $|\xi_k(\omega) - \xi(\omega)| \geq 1/m$. So

$$\{\xi_n \not\rightarrow \xi\} = \bigcup_{\epsilon > 0} A^\epsilon = \bigcup_{m=1}^{\infty} A^{\frac{1}{m}}.$$

By the continuity theorem, we have

$$P(A^\epsilon) = P\left(\bigcap_{n=1}^{\infty} \bigcup_{k \geq n} A_k^\epsilon\right) = \lim_{n \rightarrow \infty} P\left(\bigcup_{k \geq n} A_k^\epsilon\right)$$

which implies that the following relations hold:

4.3 Almost sure convergence and strong laws of large numbers

4.3.1 Convergence in probability

$$\begin{aligned} 0 = P(\{\xi_n \not\rightarrow \xi\}) &\Leftrightarrow P\left(\bigcup_{m=1}^{\infty} A_m^{\frac{1}{m}}\right) = 0 \\ &\Leftrightarrow P(A_m^{\frac{1}{m}}) = 0, \forall m \geq 1 \end{aligned}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.1 Convergence in probability

$$\begin{aligned}0 = P(\{\xi_n \not\rightarrow \xi\}) &\Leftrightarrow P\left(\bigcup_{m=1}^{\infty} A_m^{\frac{1}{m}}\right) = 0 \\&\Leftrightarrow P(A_m^{\frac{1}{m}}) = 0, \forall m \geq 1 \\&\Leftrightarrow P\left(\bigcup_{k \geq n} A_k^{\frac{1}{m}}\right) \rightarrow 0, \forall m \geq 1 \\&\Leftrightarrow P\left(\bigcup_{k \geq n} (|\xi_k - \xi| \geq \frac{1}{m})\right) \rightarrow 0, \forall m \geq 1 \\&\Leftrightarrow P\left(\bigcup_{k \geq n} (|\xi_k - \xi| \geq \epsilon)\right) \rightarrow 0, \forall \epsilon \geq 0.\end{aligned}$$

Corollary

If $\xi_n \xrightarrow{P} \xi$, then there exists a sub-sequence $\{\xi_{n_k}\}$ such that

$$\xi_{n_k} \rightarrow \xi \text{ a.s.}$$

Corollary

If $\xi_n \xrightarrow{P} \xi$, then there exists a sub-sequence $\{\xi_{n_k}\}$ such that

$$\xi_{n_k} \rightarrow \xi \text{ a.s.}$$

Proof. Let $\epsilon_k = 2^{-k}$. For any k , there exists a n_k such that

$$P(|\xi_n - \xi| \geq \epsilon_k) < \epsilon_k \quad \forall n \geq n_k.$$

Without loss of generality, we can assume

$n_1 < n_2 < \cdots < n_k < n_{k+1}$. Then for any $\epsilon > 0$, there is a k_0 such that $\epsilon_k < \epsilon$ for $k \geq k_0$.

4.3 Almost sure convergence and strong laws of large numbers

4.3.1 Convergence in probability

So

$$\sum_{k=k_0}^{\infty} P(|\xi_{n_k} - \xi| \geq \epsilon) \leq \sum_{k=k_0}^{\infty} P(|\xi_{n_k} - \xi| \geq \epsilon_k) \leq \sum_{k=k_0}^{\infty} \epsilon_k < \infty.$$

Hence

$$\xi_{n_k} \rightarrow \xi \text{ a.s.}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.1 Convergence in probability

So

$$\sum_{k=k_0}^{\infty} P(|\xi_{n_k} - \xi| \geq \epsilon) \leq \sum_{k=k_0}^{\infty} P(|\xi_{n_k} - \xi| \geq \epsilon_k) \leq \sum_{k=k_0}^{\infty} \epsilon_k < \infty.$$

Hence

$$\xi_{n_k} \rightarrow \xi \text{ a.s.}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.1 Convergence in probability

Corollary

$\xi_n - \xi_m \xrightarrow{P} 0$ as $n, m \rightarrow \infty$ if and only if

$$\exists \xi, \quad \xi_n \xrightarrow{P} \xi.$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.1 Convergence in probability

Corollary

$\xi_n - \xi_m \xrightarrow{P} 0$ as $n, m \rightarrow \infty$ if and only if

$$\exists \xi, \quad \xi_n \xrightarrow{P} \xi.$$

Proof. The "if" part is obvious. For the "if" part, for each $\epsilon_k = 2^{-k}$ there exists n_k such that

$$P(|\xi_n - \xi_m| \geq \epsilon_k) \leq \epsilon_k, \quad \forall n, m \geq n_k.$$

Corollary

$\xi_n - \xi_m \xrightarrow{P} 0$ as $n, m \rightarrow \infty$ if and only if

$$\exists \xi, \quad \xi_n \xrightarrow{P} \xi.$$

Proof. The "if" part is obvious. For the "if" part, for each $\epsilon_k = 2^{-k}$ there exists n_k such that

$$P(|\xi_n - \xi_m| \geq \epsilon_k) \leq \epsilon_k, \quad \forall n, m \geq n_k.$$

Without loss of generality, assume $n_k < n_{k+1}$. Then

$$P(|\xi_{n_{k+1}} - \xi_{n_k}| \geq \epsilon_k) \leq \epsilon_k.$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.1 Convergence in probability

It follows that

$$\begin{aligned} & P\left(\sum_{k=1}^{\infty} |\xi_{n_{k+1}} - \xi_{n_k}| = \infty\right) \\ &= P\left(\sum_{k=k_0}^{\infty} |\xi_{n_{k+1}} - \xi_{n_k}| = \infty\right) \\ &\leq P\left(\sum_{k=k_0}^{\infty} |\xi_{n_{k+1}} - \xi_{n_k}| \geq \sum_{k=k_0}^{\infty} \epsilon_k\right) \\ &\leq \sum_{k=k_0}^{\infty} \epsilon_k \rightarrow 0 \text{ as } k_0 \rightarrow \infty. \end{aligned}$$

Let $\xi_0 = 0$. For $\omega \in A = \{\sum_{k=1}^{\infty} |\xi_{n_{k+1}} - \xi_{n_k}| < \infty\}$, define

$\xi(\omega) = \sum_{k=0}^{\infty} (\xi_{n_{k+1}}(\omega) - \xi_{n_k}(\omega))$, and for $\omega \in A$, define

$\xi(\omega) = 0$.

4.3 Almost sure convergence and strong laws of large numbers

4.3.1 Convergence in probability

Then

$$\xi_{n_k} \rightarrow \xi \text{ a.s.}$$

So,

$$\xi_{n_k} \xrightarrow{P} \xi.$$

It follows that

$$\xi_n = (\xi_n - \xi_{n_k}) + \xi_{n_k} \xrightarrow{P} \xi.$$

4.3.2 Strong laws of large numbers

Definition 2 Suppose that $\{\xi_n, n \geq 1\}$ is a sequence of random variables defined on (Ω, \mathcal{F}, P) . If there exist constant sequences $\{a_n, n \geq 1\}$ and $\{b_n, n \geq 1\}$ such that

$$\frac{1}{a_n} \sum_{k=1}^n \xi_k - b_n \rightarrow 0 \quad a.s.,$$

we say that $\{\xi_n\}$ obeys the strong law of large numbers (**SLLN**).

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

Theorem 2 (Borel) Suppose that $\{\xi_n\}$ is a sequence of i.i.d. random variables defined on (Ω, \mathcal{F}, P) with $P(\xi_n = 1) = p$, $P(\xi_n = 0) = 1 - p$, $0 < p < 1$. Let $S_n = \sum_{k=1}^n \xi_k$, then

$$\frac{S_n}{n} - p \rightarrow 0 \quad a.s.$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

Proof. We have

$$\begin{aligned} & P\left(\frac{|S_n - np|}{n} \geq \epsilon\right) \\ &= P(|S_n - np|^4 \geq (\epsilon n)^4) \end{aligned}$$

Proof. We have

$$\begin{aligned} & P\left(\frac{|S_n - np|}{n} \geq \epsilon\right) \\ &= P(|S_n - np|^4 \geq (\epsilon n)^4) \\ &\leq \frac{1}{\epsilon^4 n^4} E|S_n - np|^4 \quad (\text{by Markov inequality}). \end{aligned}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

Let $\eta_i = \xi_i - p$. Then

$$\begin{aligned} E|S_n - np|^4 &= E\left|\sum_{i=1}^n \eta_i\right|^4 = \sum_{i,j,l,k} E\eta_i\eta_j\eta_l\eta_k \\ &= \sum_i E\eta_i^4 + \sum_{i \neq j} E\eta_i^2\eta_j^2 \\ &= n(q^4p + p^4q) + n(n-1)(pq)^2 \leq n^2pq. \end{aligned}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

Let $\eta_i = \xi_i - p$. Then

$$\begin{aligned} E|S_n - np|^4 &= E\left|\sum_{i=1}^n \eta_i\right|^4 = \sum_{i,j,l,k} E\eta_i\eta_j\eta_l\eta_k \\ &= \sum_i E\eta_i^4 + \sum_{i \neq j} E\eta_i^2\eta_j^2 \\ &= n(q^4p + p^4q) + n(n-1)(pq)^2 \leq n^2pq. \end{aligned}$$

So,

$$\sum_{n=1}^{\infty} P\left(\frac{|S_n - np|}{n} \geq \epsilon\right) \leq \sum_{n=1}^{\infty} \frac{n^2pq}{\epsilon^4 n^4} < \infty.$$

Hence $S_n/n \rightarrow p$ a.s.

Corollary Suppose that $\{\xi_n\}$ is a sequence of i.i.d. random variables defined on (Ω, \mathcal{F}, P) with $E\xi_n = \mu$, $E\xi_n^4 < \infty$. Let $S_n = \sum_{k=1}^n \xi_k$, then

$$\frac{S_n}{n} \rightarrow \mu \quad a.s.$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

Proof. We have

$$\begin{aligned} & P\left(\frac{|S_n - n\mu|}{n} \geq \epsilon\right) \\ &= P(|S_n - n\mu|^4 \geq (\epsilon n)^4) \end{aligned}$$

Proof. We have

$$\begin{aligned} & P\left(\frac{|S_n - n\mu|}{n} \geq \epsilon\right) \\ &= P(|S_n - n\mu|^4 \geq (\epsilon n)^4) \\ &\leq \frac{1}{\epsilon^4 n^4} E|S_n - n\mu|^4 \quad (\text{by Markov inequality}). \end{aligned}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

Let $\eta_i = \xi_i - \mu$. Then $E\eta_i = 0$,

$$E\eta_1^4 = E(\xi_1 - \mu)^4 < \infty,$$

$$\begin{aligned} E|S_n - n\mu|^4 &= E\left|\sum_{i=1}^n \eta_i\right|^4 = \sum_{i,j,l,k} E\eta_i\eta_j\eta_l\eta_k \\ &= \sum_i E\eta_i^4 + \sum_{i \neq j} E\eta_i^2\eta_j^2 \\ &= nE(\xi_1 - \mu)^4 + n(n-1)(\text{Var}(\xi_1))^2 \\ &\leq n^2 c_0. \end{aligned}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

So,

$$\sum_{n=1}^{\infty} P\left(\frac{|S_n - n\mu|}{n} \geq \epsilon\right) \leq \sum_{n=1}^{\infty} \frac{n^2 c_0}{\epsilon^4 n^4} < \infty.$$

Hence $S_n/n \rightarrow \mu$ a.s.

Theorem 3 (Kolmogorov, 1930) Suppose that $\{\xi_n, n \geq 1\}$ is a sequence of i.i.d. random variables defined on (Ω, \mathcal{F}, P) with $E|\xi_1| < \infty$, $E\xi_1 = \mu$. Let $S_n = \sum_{k=1}^n \xi_k$. Then

$$\frac{S_n}{n} \rightarrow \mu \quad a.s. \quad (1)$$

Theorem 3 (Kolmogorov, 1930) Suppose that $\{\xi_n, n \geq 1\}$ is a sequence of i.i.d. random variables defined on (Ω, \mathcal{F}, P) with $E|\xi_1| < \infty$, $E\xi_1 = \mu$. Let $S_n = \sum_{k=1}^n \xi_k$. Then

$$\frac{S_n}{n} \rightarrow \mu \quad a.s. \quad (1)$$

In fact, the converse of Theorem 2 also holds: if there exists a constant μ such that (1) holds, then the expectation of ξ_1 exists and equals to μ .

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

Theorem 4 Suppose that $\{\xi_n, n \geq 1\}$ is a sequence of independent random variables defined on (Ω, \mathcal{F}, P) with $E\xi_k = \mu_k$, $Var\xi_k < \infty$. Let $S_n = \sum_{k=1}^n \xi_k$. If

$$\sum_{n=1}^{\infty} \frac{Var\xi_n}{n^2} < \infty,$$

then

$$\frac{S_n - ES_n}{n} \rightarrow 0 \quad a.s.$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

Proof Theorem 3 from Theorem 4: Let

$\eta_k = \xi_k I\{|\xi_k| \leq k\}$. Then

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{\text{Var}(\eta_n)}{n^2} &\leq \sum_{n=1}^{\infty} \frac{E[\xi_1^2 I\{|\xi_1| \leq n\}]}{n^2} \\ &= \sum_{n=1}^{\infty} \int_n^{n+1} \frac{E[\xi_1^2 I\{|\xi_1| \leq n\}]}{n^2} dx \\ &\leq 2^2 \sum_{n=1}^{\infty} \int_n^{n+1} \frac{E[\xi_1^2 I\{|\xi_1| \leq x\}]}{x^2} dx \\ &= 2^2 \int_1^{\infty} E \left[\frac{\xi_1^2 I\{|\xi_1| \leq x\}}{x^2} \right] dx \\ &= 2^2 E \left[\int_1^{\infty} \frac{\xi_1^2 I\{|\xi_1| \leq x\}}{x^2} dx \right] \leq 4E[|\xi_1|] < \infty. \end{aligned}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

By Theorem 4,

$$\frac{\sum_{k=1}^n (\eta_k - E\eta_k)}{n} \rightarrow 0 \text{ a.s.}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

By Theorem 4,

$$\frac{\sum_{k=1}^n (\eta_k - E\eta_k)}{n} \rightarrow 0 \text{ a.s.}$$

Also,

$$\begin{aligned} \frac{\sum_{k=1}^n E\eta_k}{n} &= \frac{\sum_{k=1}^n E[\xi_1 I\{|\xi_1| \leq k\}]}{n} \\ &\rightarrow E\xi_1 = \mu. \end{aligned}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

By Theorem 4,

$$\frac{\sum_{k=1}^n (\eta_k - E\eta_k)}{n} \rightarrow 0 \text{ a.s.}$$

Also,

$$\begin{aligned} \frac{\sum_{k=1}^n E\eta_k}{n} &= \frac{\sum_{k=1}^n E[\xi_1 I\{|\xi_1| \leq k\}]}{n} \\ &\rightarrow E\xi_1 = \mu. \end{aligned}$$

It follows that

$$\frac{\sum_{k=1}^n \eta_k}{n} \rightarrow \mu \text{ a.s.}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

Finally,

$$\begin{aligned} P(\eta_k \neq \xi_k \text{ i.o.}) &= P\left(\bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} \{\eta_k \neq \xi_k\}\right) \\ &= \lim_{n \rightarrow \infty} P\left(\bigcup_{k=n}^{\infty} \{|\xi_k| \geq k\}\right) \leq \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(|\xi_k| \geq k) = 0, \end{aligned}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

Finally,

$$\begin{aligned} P(\eta_k \neq \xi_k \text{ i.o.}) &= P\left(\bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} \{\eta_k \neq \xi_k\}\right) \\ &= \lim_{n \rightarrow \infty} P\left(\bigcup_{k=n}^{\infty} \{|\xi_k| \geq k\}\right) \leq \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(|\xi_k| \geq k) = 0, \end{aligned}$$

because

$$\sum_{k=1}^{\infty} P(|\xi_k| \geq k) = \sum_{k=1}^{\infty} P(|\xi_1| \geq k) < \infty.$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

Hence,

$$\lim_{n \rightarrow \infty} \frac{\sum_{k=1}^n \xi_k}{n} = \lim_{n \rightarrow \infty} \frac{\sum_{k=1}^n \eta_k}{n} = \eta \quad a.s.$$

The proof is completed.

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

The necessary part: Suppose

$$\lim_{n \rightarrow \infty} \frac{S_n}{n} = \lim_{n \rightarrow \infty} \frac{\sum_{k=1}^n \xi_k}{n} = \mu \text{ a.s.}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

The necessary part: Suppose

$$\lim_{n \rightarrow \infty} \frac{S_n}{n} = \lim_{n \rightarrow \infty} \frac{\sum_{k=1}^n \xi_k}{n} = \mu \text{ a.s.}$$

Then

$$\frac{\xi_n}{n} = \frac{S_n}{n} - \frac{S_{n-1}}{n} = \frac{S_n}{n} - \frac{S_{n-1}}{n-1} \frac{n-1}{n} \rightarrow 0 \text{ a.s.}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

The necessary part: Suppose

$$\lim_{n \rightarrow \infty} \frac{S_n}{n} = \lim_{n \rightarrow \infty} \frac{\sum_{k=1}^n \xi_k}{n} = \mu \text{ a.s.}$$

Then

$$\frac{\xi_n}{n} = \frac{S_n}{n} - \frac{S_{n-1}}{n} = \frac{S_n}{n} - \frac{S_{n-1}}{n-1} \frac{n-1}{n} \rightarrow 0 \text{ a.s.}$$

So,

$$P(|\xi_n| \geq n \text{ i.o.}) = 0,$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

The necessary part: Suppose

$$\lim_{n \rightarrow \infty} \frac{S_n}{n} = \lim_{n \rightarrow \infty} \frac{\sum_{k=1}^n \xi_k}{n} = \mu \text{ a.s.}$$

Then

$$\frac{\xi_n}{n} = \frac{S_n}{n} - \frac{S_{n-1}}{n} = \frac{S_n}{n} - \frac{S_{n-1}}{n-1} \frac{n-1}{n} \rightarrow 0 \text{ a.s.}$$

So,

$$P(|\xi_n| \geq n \text{ i.o.}) = 0,$$

which will imply

$$\sum_{n=1}^{\infty} P(|\xi_1| \geq n) = \sum_{n=1}^{\infty} P(|\xi_n| \geq n) < \infty.$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

In fact, if

$$\sum_{n=1}^{\infty} P(|\xi_n| \geq n) = \infty.$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

In fact, if

$$\sum_{n=1}^{\infty} P(|\xi_n| \geq n) < \infty.$$

Then

$$\begin{aligned} P\left(\bigcap_{k=n}^{\infty} \{|\xi_k| < k\}\right) &= \prod_{k=n}^{\infty} P(|\xi_k| < k) = \prod_{k=n}^{\infty} \left(1 - P(|\xi_k| \geq k)\right) \\ &\leq \exp\left\{-\sum_{k=n}^{\infty} P(|\xi_k| \geq k)\right\} = 0. \end{aligned}$$

So,

$$P\left(\{|\xi_n| \geq n \text{ i.o.}\}^C\right) = P\left(\bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} \{|\xi_k| < k\}\right) = 1.$$

Borel-Cantelli Lemma

Lemma

(1) If $\sum_{n=1}^{\infty} P(A_n) < \infty$, then

$$P(A_n \text{ i.o.}) = 0.$$

(2) If $\sum_{n=1}^{\infty} P(A_n) = \infty$ and $\{A_n\}$ are independent events, then

$$P(A_n \text{ i.o.}) = 1.$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

(1)

$$\begin{aligned} P(A_k \text{ i.o.}) &= P\left(\bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k\right) \\ &= \lim_{n \rightarrow \infty} P\left(\bigcup_{k=n}^{\infty} A_k\right) \leq \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = 0. \end{aligned}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

In fact, if

$$\sum_{n=1}^{\infty} P(A_n) = \infty.$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

In fact, if

$$\sum_{n=1}^{\infty} P(A_n) = \infty.$$

Then

$$\begin{aligned} P\left(\bigcap_{k=n}^{\infty} A_k^C\right) &= \prod_{k=n}^{\infty} P(A_k^C) \\ &\leq \exp\left\{-\sum_{k=n}^{\infty} P(A_k)\right\} = 0. \end{aligned}$$

So,

$$P(\{A_n \text{ i.o.}\}^C) = P\left(\bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k^C\right) = 0.$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

Example. (the Monte Carlo method) Let $f(x)$ be a continuous function defined on $[0, 1]$ with values in $[0, 1]$, and let $\xi_1, \eta_1, \xi_2, \eta_2, \dots$ be a sequence of independent random variables with a common uniform distribution in $[0, 1]$. Define

$$\rho_i = \begin{cases} 1, & \text{if } f(\xi_i) \geq \eta_i, \\ 0, & \text{if } f(\xi_i) < \eta_i. \end{cases}$$

Then $\{\rho_i, i \geq 1\}$ are also i.i.d. random variables. Furthermore,

$$E\rho_1 = P(f(\xi_1) \geq \eta_1) = \int \int_{y \leq f(x)} dx dy = \int_0^1 f(x) dx.$$

By Theorem 3, we have

$$\frac{1}{n} \sum_{k=1}^n \rho_k \rightarrow \int_0^1 f(x) dx \quad a.s.$$

Example. (the Monte Carlo method) Suppose

$D \subset \mathbb{R}^d$ is a bounded area, $\int_D |g(\mathbf{x})| d\mathbf{x} < \infty$.

Compute $\int_D g(\mathbf{x}) d\mathbf{x}$.

Example. (the Monte Carlo method) Suppose

$D \subset \mathbb{R}^d$ is a bounded area, $\int_D |g(\mathbf{x})| d\mathbf{x} < \infty$.

Compute $\int_D g(\mathbf{x}) d\mathbf{x}$.

解: Suppose that $D \subset A$ where A is a rectangle, and ξ is a random vector uniformly distributed in A .

Denote

$$I_D(\mathbf{x}) = \begin{cases} 1, & \text{if } \mathbf{x} \in D, \\ 0, & \text{otherwise.} \end{cases}.$$

Example. (the Monte Carlo method) Suppose

$D \subset \mathbb{R}^d$ is a bounded area, $\int_D |g(\mathbf{x})| d\mathbf{x} < \infty$.

Compute $\int_D g(\mathbf{x}) d\mathbf{x}$.

解: Suppose that $D \subset A$ where A is a rectangle, and ξ is a random vector uniformly distributed in A .

Denote

$$I_D(\mathbf{x}) = \begin{cases} 1, & \text{if } \mathbf{x} \in D, \\ 0, & \text{otherwise.} \end{cases}$$

Then

$$E[g(\xi)I_D(\xi)] = \int_A \frac{g(\mathbf{x})I_D(\mathbf{x})}{m(A)} d\mathbf{x} = \frac{1}{m(A)} \int_D g(\mathbf{x}) d\mathbf{x}.$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

Let ξ_1, ξ_2, \dots be i.i.d. copies of ξ .

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

Let ξ_1, ξ_2, \dots be i.i.d. copies of ξ . Then by the strong law of large numbers,

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n g(\xi_i) I_D(\xi_i) &\rightarrow E[g(\xi) I_D(\xi)] \quad a.s. \\ &= \frac{1}{m(A)} \int_D g(\mathbf{x}) d\mathbf{x}. \end{aligned}$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

Let ξ_1, ξ_2, \dots be i.i.d. copies of ξ . Then by the strong law of large numbers,

$$\begin{aligned}\frac{1}{n} \sum_{i=1}^n g(\xi_i) I_D(\xi_i) &\rightarrow E[g(\xi) I_D(\xi)] \quad a.s. \\ &= \frac{1}{m(A)} \int_D g(\mathbf{x}) d\mathbf{x}.\end{aligned}$$

So, for large n ,

$$\int_D g(\mathbf{x}) d\mathbf{x} \approx \frac{m(A)}{n} \sum_{i=1}^n g(\xi_i) I_D(\xi_i).$$

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

If D is not bounded, we choose a probability density function $f(\mathbf{x}) > 0$, for example the d -dimensional standard normal density. $\boldsymbol{\xi}, \boldsymbol{\xi}_1, \boldsymbol{\xi}_2, \dots$ be i.i.d. random vectors with $f(\mathbf{x})$ being the pdf. Then

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \frac{g(\boldsymbol{\xi}_i) I_D(\boldsymbol{\xi}_i)}{f(\boldsymbol{\xi}_i)} &\rightarrow_E \left[\frac{g(\boldsymbol{\xi}) I_D(\boldsymbol{\xi})}{f(\boldsymbol{\xi})} \right] \quad a.s. \\ &= \int \left[\frac{g(\mathbf{x}) I_D(\mathbf{x})}{f(\mathbf{x})} f(\mathbf{x}) \right] d\mathbf{x} \\ &= \int_D g(\mathbf{x}) d\mathbf{x}. \end{aligned}$$

Convergence rate of the SLLN:

Suppose that $\{\xi_i; i \geq 1\}$ be i.i.d. random variables, $E[\xi_1] = \mu$. Then

$$\frac{S_n}{n} \rightarrow \mu \text{ a.s.}$$

The law of the iterated logarithm:

Theorem

Suppose $\text{Var}(\xi_1) = \sigma^2 < \infty$. Then

$$\limsup_{n \rightarrow \infty} \frac{S_n - n\mu}{\sqrt{2n \ln \ln n}} = \sigma \text{ a.s.} \quad (2)$$

On the other hand, if (2) holds for some μ and σ , then we must have $\text{Var}(\xi_1) = \sigma^2$ and $E\xi_1 = \mu$.

4.3 Almost sure convergence and strong laws of large numbers

4.3.2 Strong laws of large numbers

The law of the iterated logarithm tells that

$$\frac{S_n}{n} - \mu = O\left(\sqrt{\frac{\ln \ln n}{n}}\right) \quad a.s.$$

For the MC method, the error is about $\sqrt{\frac{\ln \ln n}{n}}$.