

Chapter 4 Probability Limit Theorems

In the early days, the aim of probability theory is to reveal the inherent rule of random phenomena caused by a large number of random factors.

Bernoulli first recognized the importance to study an infinite sequence of random trials, and established the first limit theorem in probability theory—the law of large numbers.

de Moivre and Laplace presented that the observed error can be regarded as the summation of a large number of independent and slight errors, and proved that the distribution of the observed error is approximated by a normal distribution—the central limit theorem.

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Definition 1 Let F be a cdf, $\{F_n, n \geq 1\}$ a sequence of cdfs. We say that F_n converges weakly to F , denoted by $F_n \xrightarrow{w} F$, if $F_n(x) \rightarrow F(x)$ holds at **every continuity point** x of F as $n \rightarrow \infty$. Let ξ be a r.v., $\{\xi_n, n \geq 1\}$ a sequence of r.v.s, we say $\{\xi_n\}$ converges in distribution to ξ , denoted by $\xi_n \xrightarrow{d} \xi$, if the cdfs ξ_n 's converges weakly to the cdf of ξ .

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Remark 1. The limit function of a sequence of distribution functions is not necessarily a distribution function. For example, let

$$F_n(x) = \begin{cases} 0, & x < n, \\ 1, & x \geq n. \end{cases}$$

This distribution function converges pointwise to 0, but $F(x) \equiv 0$ is not a distribution function.

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Remark 2. The condition that $F_n(x) \longrightarrow F(x)$ for every continuity point x of F is not a strong condition in Definition 1. For example, let

$$F_n(x) = \begin{cases} 0, & x < \frac{1}{n}, \\ 1, & x \geq \frac{1}{n}. \end{cases} \quad F(x) = \begin{cases} 0, & x < 0, \\ 1, & x \geq 0. \end{cases}$$

Then $F_n(x)$ converges pointwise to $F(x)$ except at point $x = 0$, while $x = 0$ is a unique discontinuous point of $F(x)$. Thus it follows from Definition 1 that $F_n \xrightarrow{w} F$.

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Remark 3. Since the set of discontinuous points of a distribution function F is at most countable, $F_n \xrightarrow{w} F$ means that F_n converges everywhere to F in a dense subset of \mathbf{R} .

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Theorem 1 (Helly's first theorem) Let $\{F_n, n \geq 1\}$ be a sequence of distribution functions. Then there exists a non-decreasing right-continuous function F (not necessarily a distribution function) with $0 \leq F(x) \leq 1$, $x \in \mathbf{R}$, and a subsequence F_{n_k} , such that $F_{n_k}(x) \rightarrow F(x)$ for every continuity point x of F as $k \rightarrow \infty$.

Main idea of the proof. For each given x , since $\{F_n(x)\}$ is a bounded sequence, there is a subsequence $\{F_{n_m}(x)\}$ and a number $F(x)$, such that

$$F_{n_m}(x) \rightarrow F(x).$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Main idea of the proof. For each given x , since $\{F_n(x)\}$ is a bounded sequence, there is a subsequence $\{F_{n_m}(x)\}$ and a number $F(x)$, such that

$$F_{n_m}(x) \rightarrow F(x).$$

However, the subsequence $\{n_m\}$ may depend on the value of x , i.e., $n_m = n_m(x)$. What we want to do is to find an "uniform" sequence $\{n_m\}$ which does depend on x such the above convergence holds.

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Proof. Let r_1, r_2, \dots , denote the set of rational numbers. That $0 \leq F(x) \leq 1$ means that $\{F_n(r_1)\}$ is a bounded sequence. So, there exists a convergent subsequence $\{F_{n_m^{(1)}}(r_1)\}$. Denote the limit by

$$G(r_1) = \lim_{m \rightarrow \infty} F_{n_m^{(1)}}(r_1).$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Proof. Let r_1, r_2, \dots , denote the set of rational numbers. That $0 \leq F(x) \leq 1$ means that $\{F_n(r_1)\}$ is a bounded sequence. So, there exists a convergent subsequence $\{F_{n_m^{(1)}}(r_1)\}$. Denote the limit by

$$G(r_1) = \lim_{m \rightarrow \infty} F_{n_m^{(1)}}(r_1).$$

Then, consider the bounded sequence $\{F_{n_m^{(1)}}(r_2)\}$. There exists a further convergent subsequences $\{F_{n_m^{(2)}}(r_2)\}$. Denote the limit by

$$G(r_2) = \lim_{m \rightarrow \infty} F_{n_m^{(2)}}(r_2).$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Repeating the procedure, we obtain

$$\{F_{n_m^{(k)}}\} \subset \{F_{n_m^{(k-1)}}\}, \quad G(r_k) = \lim_{m \rightarrow \infty} F_{n_m^{(k)}}(r_k), \quad k \geq 2.$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Repeating the procedure, we obtain

$$\{F_{n_m^{(k)}}\} \subset \{F_{n_m^{(k-1)}}\}, \quad G(r_k) = \lim_{m \rightarrow \infty} F_{n_m^{(k)}}(r_k), \quad k \geq 2.$$

Now, consider the diagonal sequence $\{F_{n_m^{(m)}}\}$.

Obviously,

$$\lim_{m \rightarrow \infty} F_{n_m^{(m)}}(r_k) = G(r_k), \quad \forall k \geq 1.$$

In addition, $F_n \nearrow \implies G(r) \nearrow$ and also

$$0 \leq G(r) \leq 1.$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Let

$$F(x) = \lim_{r_j \downarrow x} G(r_j) = \inf_{r_j > x} G(r_j), \quad x \in \mathbf{R}.$$

Then $F(x) \nearrow$ and also $0 \leq F(x) \leq 1$, and $F(x)$ is right-continuous. Further, if $r < x < s$ and r, s are rational numbers, then

$$G(r) \leq F(x) \leq G(s).$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Now for any continuous point x of F and $h > 0$, there are $r_i < r_j$ such that

$$x - h < r_i < x < r_j < x + h.$$

It follows that

$$F_{n_m^{(m)}}(r_i) \leq F_{n_m^{(m)}}(x) \leq F_{n_m^{(m)}}(r_j)$$



$$F(x - h) \leq G(r_i)$$

$$G(r_j) \leq F(x + h).$$

Letting $m \rightarrow \infty$ yields

$$\begin{aligned} F(x - h) \leq G(r_i) &= \lim_m F_{n_m^{(m)}}(r_i) \\ &\leq \liminf_m F_{n_m^{(m)}}(x) \\ &\leq \limsup_m F_{n_m^{(m)}}(x) \\ &\leq \limsup_m F_{n_m^{(m)}}(r_j) \\ &= G(r_j) \leq F(x + h). \end{aligned}$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Letting $m \rightarrow \infty$ yields

$$\begin{aligned} F(x-h) \leq G(r_i) &= \lim_m F_{n_m^{(m)}}(r_i) \\ &\leq \liminf_m F_{n_m^{(m)}}(x) \\ &\leq \limsup_m F_{n_m^{(m)}}(x) \\ &\leq \limsup_m F_{n_m^{(m)}}(r_j) \\ &= G(r_j) \leq F(x+h). \end{aligned}$$

Letting $h \rightarrow 0$ yields

$$\liminf_m F_{n_m^{(m)}}(x) = \limsup_m F_{n_m^{(m)}}(x) = F(x).$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Theorem 2 (Helly's second theorem) Let F be a cdf, $\{F_n, n \geq 1\}$ a sequence of cdfs such that $F_n \xrightarrow{w} F$. If $g(x)$ is a bounded continuous function in \mathbf{R} , then

$$\int_{-\infty}^{\infty} g(x) dF_n(x) \longrightarrow \int_{-\infty}^{\infty} g(x) dF(x).$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Main idea of Proof.

$$\begin{aligned} & \int_{-\infty}^{\infty} g(x) dF_n(x) \\ & \approx \sum g(x_{i-1}) [F_n(x_i) - F_n(x_{i-1})] \\ & \rightarrow \sum g(x_{i-1}) [F(x_i) - F(x_{i-1})] \\ & \quad (\text{if } x'_i\text{'s are continuous points of } F) \\ & \approx \int_{-\infty}^{\infty} g(x) dF(x). \end{aligned}$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Proof. Since g is a bounded function, there must exist a constant $c > 0$ such that $|g(x)| < c$, $x \in \mathbf{R}$.

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Proof. Since g is a bounded function, there must exist a constant $c > 0$ such that $|g(x)| < c$, $x \in \mathbf{R}$. For given $\delta > 0$ and $a > 0$ with $\pm a$ being continuous points of F , select $-a = x_0 < x_1 < \cdots < x_m = a$ such that x_i s are continuous points of F and $|\Delta x| =: \max_i |x_i - x_{i-1}| < \delta$.

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Proof. Since g is a bounded function, there must exist a constant $c > 0$ such that $|g(x)| < c$, $x \in \mathbf{R}$. For given $\delta > 0$ and $a > 0$ with $\pm a$ being continuous points of F , select $-a = x_0 < x_1 < \cdots < x_m = a$ such that x_i s are continuous points of F and $|\Delta x| =: \max_i |x_i - x_{i-1}| < \delta$. Let

$$g_m(x) = \begin{cases} g(x_{i-1}), & \text{if } x_{i-1} < x \leq x_i, \\ 0, & \text{if } x \leq -a \text{ or } x > a. \end{cases}$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

It is easily seen that

$$\int_{-\infty}^{\infty} g_m(x) dF(x) = \sum_{i=1}^m g(x_{i-1})(F(x_i) - F(x_{i-1})),$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

It is easily seen that

$$\begin{aligned}\int_{-\infty}^{\infty} g_m(x) dF(x) &= \sum_{i=1}^m g(x_{i-1})(F(x_i) - F(x_{i-1})), \\ &= \sum_{i=1}^m \int_{x_{i-1}}^{x_i} g(x_{i-1}) dF(x),\end{aligned}$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

It is easily seen that

$$\begin{aligned}\int_{-\infty}^{\infty} g_m(x) dF(x) &= \sum_{i=1}^m g(x_{i-1})(F(x_i) - F(x_{i-1})), \\ &= \sum_{i=1}^m \int_{x_{i-1}}^{x_i} g(x_{i-1}) dF(x), \\ \int_{-\infty}^{\infty} g(x) dF(x) &= \int_{-\infty}^{-a} g(x) dF(x) \\ &\quad + \sum_{i=1}^m \int_{x_{i-1}}^{x_i} g(x) dF(x) + \int_a^{\infty} g(x) dF(x).\end{aligned}$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

It follows that

$$\begin{aligned} & \left| \int_{-\infty}^{\infty} g(x) dF(x) - \int_{-\infty}^{\infty} g_m(x) dF(x) \right| \\ & \leq \left| \int_{-\infty}^{-a} g(x) dF(x) \right| + \left| \int_a^{\infty} g(x) dF(x) \right| \\ & \quad + \sum_{i=1}^m \int_{x_{i-1}}^{x_i} |g(x) - g(x_{i-1})| dF(x) \\ & \leq c[F(-a) + 1 - F(a)] + \max_i \max_{x_{i-1} \leq x \leq x_i} |g(x) - g(x_{i-1})|. \end{aligned}$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

It follows that

$$\begin{aligned}
 & \left| \int_{-\infty}^{\infty} g(x) dF(x) - \int_{-\infty}^{\infty} g_m(x) dF(x) \right| \\
 & \leq \left| \int_{-\infty}^{-a} g(x) dF(x) \right| + \left| \int_a^{\infty} g(x) dF(x) \right| \\
 & \quad + \sum_{i=1}^m \int_{x_{i-1}}^{x_i} |g(x) - g(x_{i-1})| dF(x) \\
 & \leq c[F(-a) + 1 - F(a)] + \max_i \max_{x_{i-1} \leq x \leq x_i} |g(x) - g(x_{i-1})|.
 \end{aligned}$$

Similarly,

$$\begin{aligned}
 & \left| \int_{-\infty}^{\infty} g(x) dF_n(x) - \int_{-\infty}^{\infty} g_m(x) dF_n(x) \right| \\
 & \leq c[F_n(-a) + 1 - F_n(a)] + \max_i \max_{x_{i-1} \leq x \leq x_i} |g(x) - g(x_{i-1})|.
 \end{aligned}$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

While

$$\begin{aligned} & \left| \int_{-\infty}^{\infty} g_m(x) dF_n(x) - \int_{-\infty}^{\infty} g_m(x) dF(x) \right| \\ & \leq \sum_{i=0}^m |g(x_{i-1})| [|F_n(x_i) - F(x_i)| + |F_n(x_{i-1}) - F(x_{i-1})|] \\ & \leq 2cm \max_i |F_n(x_i) - F(x_i)|. \end{aligned}$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

It follows that

$$\leq \left| \int_{-\infty}^{\infty} g(x) dF_n(x) - \int_{-\infty}^{\infty} g(x) dF(x) \right|$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

It follows that

$$\begin{aligned} & \left| \int_{-\infty}^{\infty} g(x) dF_n(x) - \int_{-\infty}^{\infty} g(x) dF(x) \right| \\ & \leq c [F_n(-a) + 1 - F_n(a)] + c [F(-a) + 1 - F(a)] \\ & \quad + 2 \max_i \max_{x_{i-1} \leq x \leq x_i} |g(x) - g(x_{i-1})| \\ & \quad + 2cm \max_i |F_n(x_i) - F(x_i)| \\ & \leq \end{aligned}$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

It follows that

$$\begin{aligned}
 & \left| \int_{-\infty}^{\infty} g(x) dF_n(x) - \int_{-\infty}^{\infty} g(x) dF(x) \right| \\
 \leq & c [F_n(-a) + 1 - F_n(a)] + c [F(-a) + 1 - F(a)] \\
 & + 2 \max_i \max_{x_{i-1} \leq x \leq x_i} |g(x) - g(x_{i-1})| \\
 & + 2cm \max_i |F_n(x_i) - F(x_i)| \\
 \leq & 2c [F(-a) + 1 - F(a)] + 2 \max_{\substack{|x-y| < \delta \\ |x|, |y| \leq a}} |g(x) - g(y)| \\
 & + 4cm \max_{i=0, \dots, m} |F_n(x_i) - F(x_i)|.
 \end{aligned}$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Now, for given $\epsilon > 0$, we first choose $a = a(\epsilon) > 0$ ($\pm a$ be continuous points of F) such that

$$F(-a) + 1 - F(a) < \epsilon/(6c).$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Now, for given $\epsilon > 0$, we first choose $a = a(\epsilon) > 0$ ($\pm a$ be continuous points of F) such that

$$F(-a) + 1 - F(a) < \epsilon/(6c).$$

Secondly, choose $\delta > 0$ such that

$$\max_{|x-y|<\delta; |x|,|y|\leq a} |g(x) - g(y)| < \epsilon/6.$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Now, for given $\epsilon > 0$, we first choose $a = a(\epsilon) > 0$ ($\pm a$ be continuous points of F) such that

$$F(-a) + 1 - F(a) < \epsilon/(6c).$$

Secondly, choose $\delta > 0$ such that

$$\max_{|x-y|<\delta; |x|,|y|\leq a} |g(x) - g(y)| < \epsilon/6.$$

Thirdly, choose m and x_i s such that x_i s are continuous points of F and $|x_i - x_{i-1}| < \delta$.

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Now, for given $\epsilon > 0$, we first choose $a = a(\epsilon) > 0$ ($\pm a$ be continuous points of F) such that

$$F(-a) + 1 - F(a) < \epsilon/(6c).$$

Secondly, choose $\delta > 0$ such that

$$\max_{|x-y|<\delta; |x|,|y|\leq a} |g(x) - g(y)| < \epsilon/6.$$

Thirdly, choose m and x_i s such that x_i s are continuous points of F and $|x_i - x_{i-1}| < \delta$. Finally, choose $N = N(\epsilon)$ such that

$$\max_{i=0,\dots,m} |F_n(x_i) - F(x_i)| < \epsilon/(12cm) \quad n \geq N.$$

The proof is now completed.

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

The second proof. Let $U \sim U(0, 1)$. Then

$\xi_n = F_n^{-1}(U) \sim F_n$, $\xi = F^{-1}(U) \sim F$. Then

$$\begin{aligned}\int_{-\infty}^{\infty} g(x) dF_n(x) &= E g(\xi_n) = \int_0^1 g(F_n^{-1}(y)) dy, \\ \int_{-\infty}^{\infty} g(x) dF(x) &= E g(\xi) = \int_0^1 g(F^{-1}(y)) dy.\end{aligned}$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

The second proof. Let $U \sim U(0, 1)$. Then

$\xi_n = F_n^{-1}(U) \sim F_n$, $\xi = F^{-1}(U) \sim F$. Then

$$\begin{aligned}\int_{-\infty}^{\infty} g(x) dF_n(x) &= E g(\xi_n) = \int_0^1 g(F_n^{-1}(y)) dy, \\ \int_{-\infty}^{\infty} g(x) dF(x) &= E g(\xi) = \int_0^1 g(F^{-1}(y)) dy.\end{aligned}$$

It can be shown that

$$F_n \xrightarrow{w} F \iff F_n^{-1}(y) \rightarrow F^{-1}(y) \quad \forall y \in C(F^{-1}),$$

where $C(F^{-1})$ is the set of continuity points of F^{-1} .

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Hence

$$F_n^{-1}(y) \rightarrow F^{-1}(y) \quad a.e. \ L.$$

It follows that

$$g(F_n^{-1}(y)) \rightarrow g(F^{-1}(y)) \quad a.e. \ L.$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Hence

$$F_n^{-1}(y) \rightarrow F^{-1}(y) \quad a.e. \ L.$$

It follows that

$$g(F_n^{-1}(y)) \rightarrow g(F^{-1}(y)) \quad a.e. \ L.$$

So

$$\int_{-\infty}^{\infty} g(x) dF_n(x) \rightarrow \int_{-\infty}^{\infty} g(x) dF(x).$$

Remark.

- ① If $g(x)$ is continuous, and $\{g_t(x)\}$ satisfy $|g_t(x)| \leq c$ and $|g_t(x) - g_t(y)| \leq |g(x) - g(y)|$, then uniformly in t ,

$$\int_{-\infty}^{\infty} g_t(x) dF_n(x) \longrightarrow \int_{-\infty}^{\infty} g_t(x) dF(x).$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Remark.

- ① If $g(x)$ is continuous, and $\{g_t(x)\}$ satisfy $|g_t(x)| \leq c$ and $|g_t(x) - g_t(y)| \leq |g(x) - g(y)|$, then uniformly in t ,

$$\int_{-\infty}^{\infty} g_t(x) dF_n(x) \longrightarrow \int_{-\infty}^{\infty} g_t(x) dF(x).$$

- ② If $F_n(x), F(x) \nearrow$ are right continuous, and for any continuous point x of F , $F_n(x) \rightarrow F(x)$, then for continuous points $A < B$ of F , and continuous $g(\cdot)$,

$$\int_A^B g(x) dF_n(x) \longrightarrow \int_A^B g(x) dF(x).$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Theorem 3 (Lévy's continuity theorem) Let F be a cdf, $\{F_n, n \geq 1\}$ a sequence of cdfs. If $F_n \xrightarrow{w} F$, then the corresponding sequence of characteristic functions $\{f_n(t)\}$ converges to the characteristic function $f(t)$ of F uniformly in t on any finite interval.

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Theorem 3 (Lévy's continuity theorem) Let F be a cdf, $\{F_n, n \geq 1\}$ a sequence of cdfs. If $F_n \xrightarrow{w} F$, then the corresponding sequence of characteristic functions $\{f_n(t)\}$ converges to the characteristic function $f(t)$ of F uniformly in t on any finite interval.

Proof. Let $g_t(x) = e^{itx}$, then

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Theorem 3 (Lévy's continuity theorem) Let F be a cdf, $\{F_n, n \geq 1\}$ a sequence of cdfs. If $F_n \xrightarrow{w} F$, then the corresponding sequence of characteristic functions $\{f_n(t)\}$ converges to the characteristic function $f(t)$ of F uniformly in t on any finite interval.

Proof. Let $g_t(x) = e^{itx}$, then $|g_t(x)| = 1$ and $\sup_{|t| \leq b} |g_t(x) - g_t(y)| \leq b|x - y|$.

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Theorem 4 (The converse limit theorem) Let $f_n(t)$ be characteristic function of distribution function $F_n(x)$, if for every t , $f_n(t) \longrightarrow f(t)$, and $f(t)$ is continuous on $t = 0$, then $f(t)$ must be a characteristic function of some distribution function F , and $F_n \xrightarrow{w} F$.

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Theorem 4 (The converse limit theorem) Let $f_n(t)$ be characteristic function of distribution function $F_n(x)$, if for every t , $f_n(t) \longrightarrow f(t)$, and $f(t)$ is continuous on $t = 0$, then $f(t)$ must be a characteristic function of some distribution function F , and $F_n \xrightarrow{w} F$.

We want to prove that there exists a cdf F such that for any subsequence $\{n'\}$ there is a further subsequence $\{n''\}$ for which

$$F_{n''} \xrightarrow{w} F.$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Proof. It suffices to prove that for any subsequence $\{n'\}$ there is a further subsequence $\{n''\}$ and a cdf $F(x)$ (which may depend on the subsequence) such that

$$F_{n''} \xrightarrow{w} F. \quad (*)$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

In fact, if (*) holds then

$$f_{n''}(t) \rightarrow f_F(t),$$

due to Lévy's continuity theorem, here $f_F(t)$ is the c.f. of F .

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

In fact, if (*) holds then

$$f_{n''}(t) \rightarrow f_F(t),$$

due to Lévy's continuity theorem, here $f_F(t)$ is the c.f. of F . By the assumption of the theorem, we must have $f_F \equiv f$.

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

In fact, if (*) holds then

$$f_{n''}(t) \rightarrow f_F(t),$$

due to Lévy's continuity theorem, here $f_F(t)$ is the c.f. of F . By the assumption of the theorem, we must have $f_F \equiv f$. So, $f(t)$ is a c.f. and $F(x)$ is uniquely determined by $f(t)$ (and hence does not depend on the subsequence). And further, (*) means that $F_n \xrightarrow{w} F$.

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Now, by Helly's first theorem, there exists a non-decreasing right-continuous function F (not necessarily a distribution function) with $0 \leq F(x) \leq 1$, $x \in \mathbf{R}$, and a subsequence $\{n''\} \subset \{n'\}$ such that

$$F_{n''}(x) \rightarrow F(x), \forall \text{ continuous point } x \text{ of } F.$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Now, by Helly's first theorem, there exists a non-decreasing right-continuous function F (not necessarily a distribution function) with $0 \leq F(x) \leq 1$, $x \in \mathbf{R}$, and a subsequence $\{n''\} \subset \{n'\}$ such that

$$F_{n''}(x) \rightarrow F(x), \forall \text{ continuous point } x \text{ of } F.$$

Next, it suffices to show that $F(x)$ is a **cdf**, i.e., $F(+\infty) - F(-\infty) = 1$.

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Notice that if $a > 0$ and $\pm a$ are continuous points of F , then

$$F(a) - F(-a) = \lim_{n''} [F_{n''}(a) - F_{n''}(-a)] .$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Notice that if $a > 0$ and $\pm a$ are continuous points of F , then

$$F(a) - F(-a) = \lim_{n''} [F_{n''}(a) - F_{n''}(-a)].$$

We need only to show that for any given $\epsilon > 0$, if a is sufficiently large, then

$$\limsup_n \int_{|x| \geq a} dF_n(x) \leq \epsilon. \quad (**)$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Now, for any $u > 0$,

$$\frac{1}{2u} \int_{-u}^u (1 - f_n(t)) dt = \frac{1}{2u} \int_{-u}^u \int_{-\infty}^{\infty} (1 - e^{itx}) dF_n(x) dt$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Now, for any $u > 0$,

$$\begin{aligned} \frac{1}{2u} \int_{-u}^u (1 - f_n(t)) dt &= \frac{1}{2u} \int_{-u}^u \int_{-\infty}^{\infty} (1 - e^{itx}) dF_n(x) dt \\ &= \int_{-\infty}^{\infty} \int_{-u}^u \frac{1}{2u} (1 - e^{itx}) dt dF(x) = \int_{-\infty}^{\infty} \left(1 - \frac{\sin ux}{ux} \right) dF_n(x) \end{aligned}$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Now, for any $u > 0$,

$$\begin{aligned} & \frac{1}{2u} \int_{-u}^u (1 - f_n(t)) dt = \frac{1}{2u} \int_{-u}^u \int_{-\infty}^{\infty} (1 - e^{itx}) dF_n(x) dt \\ &= \int_{-\infty}^{\infty} \int_{-u}^u \frac{1}{2u} (1 - e^{itx}) dt dF(x) = \int_{-\infty}^{\infty} \left(1 - \frac{\sin ux}{ux}\right) dF_n(x) \\ &\geq \int_{|x| \geq 2/u} \left(1 - \frac{\sin ux}{ux}\right) dF_n(x) \geq \frac{1}{2} \int_{|x| \geq 2/u} dF_n(x). \end{aligned}$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

So,

$$\begin{aligned} & \limsup_n \int_{|x| \geq 2/u} dF_n(x) \\ & \leq \limsup_n \frac{1}{u} \int_{-u}^u |1 - f_n(t)| dt \\ & \leq \frac{1}{u} \int_{-u}^u |1 - f(t)| dt \end{aligned}$$

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

So,

$$\begin{aligned} & \limsup_n \int_{|x| \geq 2/u} dF_n(x) \\ & \leq \limsup_n \frac{1}{u} \int_{-u}^u |1 - f_n(t)| dt \\ & \leq \frac{1}{u} \int_{-u}^u |1 - f(t)| dt \end{aligned}$$

Since $f(t)$ is continuous at $t = 0$, we can choose $u > 0$ small enough such that $|1 - f(t)| < \epsilon/2$ whenever $|t| \leq u$. And then (**) is proved.

Summary: The following are equivalent:

- ① $F_n \xrightarrow{w} F$ ($\xi_n \xrightarrow{d} \xi$);
- ② $\int g(x)dF_n(x) \rightarrow \int g(x)dF(x)$ for every bounded, continuous function g ;
- ③ $\int g(x)dF_n(x) \rightarrow \int g(x)dF(x)$ for every bounded, uniformly continuous function;
- ④ $\int g(x)dF_n(x) \rightarrow \int g(x)dF(x)$ for every bounded, continuous function g having bounded, continuous derivatives of each order;
- ⑤ $f_n(t) \rightarrow f(t)$ for all t .

Example 1. Prove the Poisson approximation of binomial distributions by the method of characteristic function.

Proof.

Example 1. Prove the Poisson approximation of binomial distributions by the method of characteristic function.

Proof. Let $\xi_n \sim B(n, p_n)$, and $\lim_{n \rightarrow \infty} np_n = \lambda$. Then its c.f. is $f_n(t) = (p_n e^{it} + q_n)^n$.

Example 1. Prove the Poisson approximation of binomial distributions by the method of characteristic function.

Proof. Let $\xi_n \sim B(n, p_n)$, and $\lim_{n \rightarrow \infty} np_n = \lambda$. Then its c.f. is $f_n(t) = (p_n e^{it} + q_n)^n$. So

$$f_n(t) = \left(1 + \frac{n \cdot p_n(e^{it} - 1)}{n}\right)^n$$

Example 1. Prove the Poisson approximation of binomial distributions by the method of characteristic function.

Proof. Let $\xi_n \sim B(n, p_n)$, and $\lim_{n \rightarrow \infty} np_n = \lambda$. Then its c.f. is $f_n(t) = (p_n e^{it} + q_n)^n$. So

$$f_n(t) = \left(1 + \frac{n \cdot p_n (e^{it} - 1)}{n}\right)^n \rightarrow e^{\lambda(e^{it} - 1)}.$$

This is just the c.f. of $P(\lambda)$.

4.1 Convergence in distribution and central limit theorems

4.1.1 Weak convergence of distribution functions

Example 1. Prove the Poisson approximation of binomial distributions by the method of characteristic function.

Proof. Let $\xi_n \sim B(n, p_n)$, and $\lim_{n \rightarrow \infty} np_n = \lambda$. Then its c.f. is $f_n(t) = (p_n e^{it} + q_n)^n$. So

$$f_n(t) = \left(1 + \frac{n \cdot p_n(e^{it} - 1)}{n}\right)^n \rightarrow e^{\lambda(e^{it} - 1)}.$$

This is just the c.f. of $P(\lambda)$. It follows that from the converse limit theorem, the binomial distribution $B(n, p_n)$ converges in distribution to the Poisson distribution $P(\lambda)$.

4.1 Convergence in distribution and central limit theorems

4.1.2 Properties

4.1.2 Properties

4.1.2 Properties

- ① Let $\{F_n, n \geq 1\}$ be a sequence of distribution functions. If $F_n \xrightarrow{d} F$, and F is a continuous distribution function, then

$$\sup_x |F_n(x) - F(x)| \rightarrow 0.$$

(Proof as exercise).

4.1.2 Properties

- ① Let $\{F_n, n \geq 1\}$ be a sequence of distribution functions. If $F_n \xrightarrow{d} F$, and F is a continuous distribution function, then

$$\sup_x |F_n(x) - F(x)| \rightarrow 0.$$

(Proof as exercise).

- ② Let ξ be a random variable, $\{\xi_n, n \geq 1\}$ a sequence of random variables, $g(x)$ a continuous function on \mathbf{R} . If $\xi_n \xrightarrow{d} \xi$, then $g(\xi_n) \xrightarrow{d} g(\xi)$. (Proof.)

- 3 Let $\{a_n, n \geq 1\}$ and $\{b_n, n \geq 1\}$ be two sequences of constants, F a distribution function, $\{F_n, n \geq 1\}$ a sequence of distribution functions. If $a_n \rightarrow a$, $b_n \rightarrow b$, $F_n \xrightarrow{w} F$, then

$$F_n(a_n x + b_n) \rightarrow F(ax + b),$$

where x is such that $ax + b$ is a continuity point of F .

Proof. Let x be s.t. $ax + b$ is a continuity point of F , and let $\varepsilon > 0$ be s.t. F is continuous on the point $ax + b \pm \varepsilon$.

Proof. Let x be s.t. $ax + b$ is a continuity point of F , and let $\varepsilon > 0$ be s.t. F is continuous on the point $ax + b \pm \varepsilon$. Obviously, $a_n x + b_n \rightarrow ax + b$. So, for n large enough,

$$ax + b - \varepsilon \leq a_n x + b_n \leq ax + b + \varepsilon,$$

Proof. Let x be s.t. $ax + b$ is a continuity point of F , and let $\varepsilon > 0$ be s.t. F is continuous on the point $ax + b \pm \varepsilon$. Obviously, $a_n x + b_n \rightarrow ax + b$. So, for n large enough,

$$ax + b - \varepsilon \leq a_n x + b_n \leq ax + b + \varepsilon,$$

which implies

$$F_n(ax + b - \varepsilon) \leq F_n(a_n x + b_n) \leq F_n(ax + b + \varepsilon).$$

Proof. Let x be s.t. $ax + b$ is a continuity point of F , and let $\varepsilon > 0$ be s.t. F is continuous on the point $ax + b \pm \varepsilon$. Obviously, $a_n x + b_n \rightarrow ax + b$. So, for n large enough,

$$ax + b - \varepsilon \leq a_n x + b_n \leq ax + b + \varepsilon,$$

which implies

$$F_n(ax + b - \varepsilon) \leq F_n(a_n x + b_n) \leq F_n(ax + b + \varepsilon).$$

Since $F_n \xrightarrow{w} F$, letting $n \rightarrow \infty$ yields

Proof. Let x be s.t. $ax + b$ is a continuity point of F , and let $\varepsilon > 0$ be s.t. F is continuous on the point $ax + b \pm \varepsilon$. Obviously, $a_n x + b_n \rightarrow ax + b$. So, for n large enough,

$$ax + b - \varepsilon \leq a_n x + b_n \leq ax + b + \varepsilon,$$

which implies

$$F_n(ax + b - \varepsilon) \leq F_n(a_n x + b_n) \leq F_n(ax + b + \varepsilon).$$

Since $F_n \xrightarrow{w} F$, letting $n \rightarrow \infty$ yields

$$\begin{aligned} F(ax + b - \varepsilon) &\leq \liminf_{n \rightarrow \infty} F(a_n x + b_n) \\ &\leq \limsup_{n \rightarrow \infty} F_n(a_n x + b_n) \leq F(ax + b + \varepsilon). \end{aligned}$$

Proof. Let x be s.t. $ax + b$ is a continuity point of F , and let $\varepsilon > 0$ be s.t. F is continuous on the point $ax + b \pm \varepsilon$. Obviously, $a_n x + b_n \rightarrow ax + b$. So, for n large enough,

$$ax + b - \varepsilon \leq a_n x + b_n \leq ax + b + \varepsilon,$$

which implies

$$F_n(ax + b - \varepsilon) \leq F_n(a_n x + b_n) \leq F_n(ax + b + \varepsilon).$$

Since $F_n \xrightarrow{w} F$, letting $n \rightarrow \infty$ yields

$$\begin{aligned} F(ax + b - \varepsilon) &\leq \liminf_{n \rightarrow \infty} F(a_n x + b_n) \\ &\leq \limsup_{n \rightarrow \infty} F_n(a_n x + b_n) \leq F(ax + b + \varepsilon). \end{aligned}$$

Letting $\varepsilon \rightarrow 0$, the proof is complete.

Corollary. If $\xi_n \xrightarrow{d} \xi$, $a_n \rightarrow a$, $b_n \rightarrow b$, then
 $a_n \xi_n + b_n \xrightarrow{d} a\xi + b$, ($a_n, a \neq 0$).

Proof.

Corollary. If $\xi_n \xrightarrow{d} \xi$, $a_n \rightarrow a$, $b_n \rightarrow b$, then $a_n \xi_n + b_n \xrightarrow{d} a\xi + b$, ($a_n, a \neq 0$).

Proof. It suffices to observe that the distribution functions of $a_n \xi_n + b_n$ and $a\xi + b$ are $F_n(\frac{x-b_n}{a_n})$ and $F(\frac{x-b}{a})$ respectively, when $a_n > 0$ and $a > 0$;

Corollary. If $\xi_n \xrightarrow{d} \xi$, $a_n \rightarrow a$, $b_n \rightarrow b$, then $a_n \xi_n + b_n \xrightarrow{d} a\xi + b$, ($a_n, a \neq 0$).

Proof. It suffices to observe that the distribution functions of $a_n \xi_n + b_n$ and $a\xi + b$ are $F_n(\frac{x-b_n}{a_n})$ and $F(\frac{x-b}{a})$ respectively, when $a_n > 0$ and $a > 0$; and are $1 - F_n(\frac{x-b_n}{a_n} - 0)$ and $1 - F(\frac{x-b}{a} - 0)$ respectively, when $a_n < 0$ and $a < 0$.

4.1.3 Central limit theorems

Let S_n denote the number of successes in n Bernoulli trials, then $P(S_n = k) = b(k; n, p)$. In practice, people are usually interested to calculate

$$P(\alpha < S_n \leq \beta) = \sum_{\alpha < k \leq \beta} b(k; n, p).$$

The computation of the right hand side of the equality is generally very complex. However, it is found by de Moivre and Laplace that the binomial distribution can be well approximated by normal distribution when $n \longrightarrow \infty$.

Theorem 5 (de Moivre-Laplace) Let $\Phi(x)$ be the standard normal distribution function. We have for $-\infty < x < \infty$,

$$\lim_{n \rightarrow \infty} P\left(\frac{S_n - np}{\sqrt{npq}} \leq x\right) = \Phi(x),$$

i.e.,

$$\frac{S_n - np}{\sqrt{npq}} \xrightarrow{d} N(0, 1).$$

4.1 Convergence in distribution and central limit theorems

4.1.3 Central limit theorems

When n is big enough, p is moderate, then

$$\begin{aligned} & P(\alpha < S_n \leq \beta) \\ = & P\left(\frac{\alpha - np}{\sqrt{npq}} < \frac{S_n - np}{\sqrt{npq}} \leq \frac{\beta - np}{\sqrt{npq}}\right) \\ \approx & \Phi\left(\frac{\beta - np}{\sqrt{npq}}\right) - \Phi\left(\frac{\alpha - np}{\sqrt{npq}}\right). \end{aligned}$$

Remark. Suppose $S_n \sim B(n, p)$.

- $np_n \rightarrow \lambda$, then

$$S_n \dot{\sim} P(np),$$

(in practical, if p is close to 0 (or 1), and np is not big (or not small), we use $P(\lambda)$ to approximate $B(n, p)$);

- fixed $0 < p < 1$, as $n \rightarrow \infty$,

$$S_n \dot{\sim} N(np, npq),$$

(in practical, if p is moderate, we use the normal distribution to approximate $B(n, p)$).

Example 3. Rolling a fair coin, how many times need one roll to ensure the probability the proportion of heads is between 0.4 and 0.6 is not smaller than 90%.

Solution.

Example 3. Rolling a fair coin, how many times need one roll to ensure the probability the proportion of heads is between 0.4 and 0.6 is not smaller than 90%.

Solution. Let n be the number of times of rolling the coin, S_n the number of times of appearing head, then $S_n \sim B(n, 1/2)$. Note that n is to satisfy

$$P(0.4 < \frac{S_n}{n} \leq 0.6) \geq 0.9.$$

From Theorem 5,

$$\begin{aligned} & P(0.4 < \frac{S_n}{n} \leq 0.6) \\ = & P(\frac{0.4n - n/2}{\sqrt{n/4}} < \frac{S_n - n/2}{\sqrt{n/4}} \leq \frac{0.6n - n/2}{\sqrt{n/4}}) \end{aligned}$$

From Theorem 5,

$$\begin{aligned} & P(0.4 < \frac{S_n}{n} \leq 0.6) \\ &= P(\frac{0.4n - n/2}{\sqrt{n/4}} < \frac{S_n - n/2}{\sqrt{n/4}} \leq \frac{0.6n - n/2}{\sqrt{n/4}}) \\ &\approx \Phi(0.2\sqrt{n}) - \Phi(-0.2\sqrt{n}) \\ &= 2\Phi(0.2\sqrt{n}) - 1. \end{aligned}$$

From Theorem 5,

$$\begin{aligned} & P(0.4 < \frac{S_n}{n} \leq 0.6) \\ &= P\left(\frac{0.4n - n/2}{\sqrt{n/4}} < \frac{S_n - n/2}{\sqrt{n/4}} \leq \frac{0.6n - n/2}{\sqrt{n/4}}\right) \\ &\approx \Phi(0.2\sqrt{n}) - \Phi(-0.2\sqrt{n}) \\ &= 2\Phi(0.2\sqrt{n}) - 1. \end{aligned}$$

Take $n \geq 69$ such that the above equality ≥ 0.9 .

Definition 2 Let $\{\xi_n, n \geq 1\}$ be a sequence of random variables. If there exist two sequences of constants $B_n > 0$ and A_n such that

$$\frac{1}{B_n} \sum_{k=1}^n \xi_k - A_n \xrightarrow{d} N(0, 1),$$

then we say that $\{\xi_n\}$ obeys the central limit theorem.

Theorem 6 (Lindeberg-Lévy) Let $\{\xi_n, n \geq 1\}$ be a sequence of independent and identically distributed random variables. Let $S_n = \sum_{k=1}^n \xi_k$, $E\xi_1 = a$, $Var\xi_1 = \sigma^2$. Then the central limit theorem holds true, i.e.,

$$\frac{S_n - na}{\sqrt{n}\sigma} \xrightarrow{d} N(0, 1) \quad \text{as } n \rightarrow \infty.$$

4.1 Convergence in distribution and central limit theorems

4.1.3 Central limit theorems

Proof.

Proof. Let $f(t)$ and $f_n(t)$ be c.f.s of $\xi_1 - a$ and $\frac{S_n - na}{\sqrt{n}\sigma}$ respectively. Since $\xi_1, \xi_2, \dots, \xi_n$ are i.i.d., we have $f_n(t) = (f(\frac{t}{\sqrt{n}\sigma}))^n$. And note that $E\xi_1 = a$, $Var\xi_1 = \sigma^2$, so the c.f. $f(t)$ has continuous derivative of 2-order, and $f'(0) = 0$, $f''(0) = -\sigma^2$. Using Taylor's expansion for f , we have

Proof. Let $f(t)$ and $f_n(t)$ be c.f.s of $\xi_1 - a$ and $\frac{S_n - na}{\sqrt{n}\sigma}$ respectively. Since $\xi_1, \xi_2, \dots, \xi_n$ are i.i.d., we have $f_n(t) = (f(\frac{t}{\sqrt{n}\sigma}))^n$. And note that $E\xi_1 = a$, $Var\xi_1 = \sigma^2$, so the c.f. $f(t)$ has continuous derivative of 2-order, and $f'(0) = 0$, $f''(0) = -\sigma^2$. Using Taylor's expansion for f , we have

$$\begin{aligned} f(x) &= f(0) + f'(0)x + \frac{1}{2}f''(0)x^2 + o(x^2) \\ &= 1 - \frac{\sigma^2}{2}x^2 + o(x^2) \quad \text{as } x \rightarrow 0. \end{aligned}$$

4.1 Convergence in distribution and central limit theorems

4.1.3 Central limit theorems

For given $t \in \mathbf{R}$,

4.1 Convergence in distribution and central limit theorems

4.1.3 Central limit theorems

For given $t \in \mathbf{R}$,

$$f\left(\frac{t}{\sqrt{n}\sigma}\right) = 1 - \frac{t^2}{2n} + o\left(\frac{1}{n}\right) \quad \text{as } n \rightarrow \infty.$$

For given $t \in \mathbf{R}$,

$$f\left(\frac{t}{\sqrt{n}\sigma}\right) = 1 - \frac{t^2}{2n} + o\left(\frac{1}{n}\right) \quad \text{as } n \rightarrow \infty.$$

Therefore

$$f_n(t) = \left(1 - \frac{t^2}{2n} + o\left(\frac{1}{n}\right)\right)^n \rightarrow e^{-\frac{t^2}{2}}.$$

The later is the c.f. of $N(0, 1)$. Then Theorem 6 follows from Theorem 4.

Example 4. When we do approximate calculation, the original data x_k rounds off to the m -th decimal place. In this way, the rounding error ξ_k can be regarded as a uniformly distributed random variable in $(-0.5 \cdot 10^{-m}, 0.5 \cdot 10^{-m}]$. If we obtain the sum $\sum_{k=1}^n x_k$ of n x'_k s, how about the error according to the rounding principle?

Example 4. When we do approximate calculation, the original data x_k rounds off to the m -th decimal place. In this way, the rounding error ξ_k can be regarded as a uniformly distributed random variable in $(-0.5 \cdot 10^{-m}, 0.5 \cdot 10^{-m}]$. If we obtain the sum $\sum_{k=1}^n x_k$ of n x'_k s, how about the error according to the rounding principle?

One may usually estimate the error of $\sum_{k=1}^n x_k$ by the sum of ξ'_k s upper bounds, that is $0.5 \cdot n \cdot 10^{-m}$. When n is very big, this number is also very big.

In fact, possibility that the error is so big is very small. Since $\{\xi_k\}$ are independent and identically distributed and $E\xi_k = 0$, $Var\xi_k = \sigma^2 = 10^{-2m}/12$, we have

4.1 Convergence in distribution and central limit theorems

4.1.3 Central limit theorems

In fact, possibility that the error is so big is very small. Since $\{\xi_k\}$ are independent and identically distributed and $E\xi_k = 0$, $Var\xi_k = \sigma^2 = 10^{-2m}/12$, we have

$$P(|\sum_{k=1}^n \xi_k| \leq x\sqrt{n}\sigma) \approx 2\Phi(x) - 1$$

by Theorem 6. The above probability is 0.997 when $x = 3$. The probability that the error of the sum exceeds $3\sigma\sqrt{n} = 0.5 \cdot \sqrt{3} \cdot \sqrt{n} \cdot 10^{-m}$ is only 0.003. Obviously, for large n , the error bound is far smaller than $0.5 \cdot n \cdot 10^{-m}$.

4.1 Convergence in distribution and central limit theorems

4.1.3 Central limit theorems

Non i.i.d. case: Let $B_n^2 = \sum_{k=1}^n \text{Var}\xi_k$.

Theorem 6 (Lindeberg-Feller) Suppose that $\{\xi_k, k \geq 1\}$ is a sequence of indept. r.v.s. If the Lindeberg condition is satisfied:

$$\frac{1}{B_n^2} \sum_{k=1}^n \int_{|x - E\xi_k| \geq \varepsilon B_n} (x - E\xi_k)^2 dF_k(x) \rightarrow 0 \quad \forall \varepsilon > 0, \quad (1)$$

then

$$\frac{\sum_{k=1}^n (\xi_k - E\xi_k)}{B_n} \xrightarrow{d} \Phi(x). \quad (2)$$

4.1 Convergence in distribution and central limit theorems

4.1.3 Central limit theorems

Non i.i.d. case: Let $B_n^2 = \sum_{k=1}^n \text{Var}\xi_k$.

Theorem 6 (Lindeberg-Feller) Suppose that $\{\xi_k, k \geq 1\}$ is a sequence of indept. r.v.s. If the Lindeberg condition is satisfied:

$$\frac{1}{B_n^2} \sum_{k=1}^n \int_{|x - E\xi_k| \geq \epsilon B_n} (x - E\xi_k)^2 dF_k(x) \rightarrow 0 \quad \forall \epsilon > 0, \quad (1)$$

then

$$\frac{\sum_{k=1}^n (\xi_k - E\xi_k)}{B_n} \xrightarrow{d} \Phi(x). \quad (2)$$

Conversely, if (2) and

$$\lim_{n \rightarrow \infty} \max_{1 \leq k \leq n} \frac{\text{Var}\xi_k}{B_n^2} = 0 \quad \text{Feller's condition}, \quad (3)$$

then (1) holds.

Theorem 7 (Lyapunov) Suppose that $\{\xi_k, k \geq 1\}$ is a sequence of indep. r.v.s, which satisfy

$$\frac{1}{(\sum_{k=1}^n \text{Var} \xi_k)^{1+\delta/2}} \sum_{k=1}^n E|\xi_k - E\xi_k|^{2+\delta} \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

then the central limit theorem holds true.

Proof.

Theorem 7 (Lyapunov) Suppose that $\{\xi_k, k \geq 1\}$ is a sequence of indep. r.v.s, which satisfy

$$\frac{1}{(\sum_{k=1}^n Var \xi_k)^{1+\delta/2}} \sum_{k=1}^n E|\xi_k - E\xi_k|^{2+\delta} \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

then the central limit theorem holds true.

Proof.

$$\frac{1}{B_n^2} \sum_{k=1}^n \int_{|x - E\xi_k| \geq \varepsilon B_n} (x - E\xi_k)^2 dF_k(x)$$

\leq

4.1 Convergence in distribution and central limit theorems

4.1.3 Central limit theorems

Theorem 7 (Lyapunov) Suppose that $\{\xi_k, k \geq 1\}$ is a sequence of indep. r.v.s, which satisfy

$$\frac{1}{(\sum_{k=1}^n \text{Var} \xi_k)^{1+\delta/2}} \sum_{k=1}^n E|\xi_k - E\xi_k|^{2+\delta} \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

then the central limit theorem holds true.

Proof.

$$\begin{aligned} & \frac{1}{B_n^2} \sum_{k=1}^n \int_{|x - E\xi_k| \geq \varepsilon B_n} (x - E\xi_k)^2 dF_k(x) \\ & \leq \frac{1}{\varepsilon^\delta} \frac{1}{B_n^{2+\delta}} \sum_{k=1}^n E|\xi_k - E\xi_k|^{2+\delta} \rightarrow 0. \end{aligned}$$

Example 7. An insurance company issues two kinds of one-year-term life insurance with random claim amounts 10,000 yuan and 20,000 yuan respectively. The claim probability q_k and the number of insurant n_k are denoted Table below.

Type k	q_k	claim amounts b_k	n_k
1	0.02	1	500
2	0.02	2	500
3	0.10	1	300
4	0.10	2	500

The insurance company hopes that the probability that the sum of claims exceeds the total premium is only 0.05. Now the premium is priced according to the expectation value principle, that is, the premium of policy i is $\pi(X_i) = (1 + \theta)EX_i$, it is required to estimate the value of θ .

Solution. $S = \sum_{i=1}^{1800} X_i$. θ is to satisfy
 $P(S \leq \pi(S)) = 0.95$. While,

$$\begin{aligned} ES &= \sum_{i=1}^{1800} EX_i = \sum_{k=1}^4 n_k b_k q_k \\ &= 500 \cdot 1 \cdot 0.02 + 500 \cdot 2 \cdot 0.02 + 300 \cdot 1 \cdot 0.10 + 500 \cdot 2 \cdot 0.10 \\ &= 160, \end{aligned}$$

4.1 Convergence in distribution and central limit theorems

4.1.3 Central limit theorems

Solution. $S = \sum_{i=1}^{1800} X_i$. θ is to satisfy
 $P(S \leq \pi(S)) = 0.95$. While,

$$\begin{aligned} ES &= \sum_{i=1}^{1800} EX_i = \sum_{k=1}^4 n_k b_k q_k \\ &= 500 \cdot 1 \cdot 0.02 + 500 \cdot 2 \cdot 0.02 + 300 \cdot 1 \cdot 0.10 + 500 \cdot 2 \cdot 0.10 \\ &= 160, \end{aligned}$$

$$\begin{aligned} Var S &= \sum_{i=1}^{1800} Var X_i = \sum_{k=1}^4 n_k b_k^2 q_k (1 - q_k) \\ &= 500 \cdot 1^2 \cdot 0.02 \cdot 0.98 + 500 \cdot 2^2 \cdot 0.02 \cdot 0.98 \\ &\quad + 300 \cdot 1^2 \cdot 0.10 \cdot 0.90 + 500 \cdot 2^2 \cdot 0.10 \cdot 0.90 \\ &= 256. \end{aligned}$$

From these we obtain the sum of premium

$$\pi(S) = (1 + \theta)ES = 160(1 + \theta).$$

From these we obtain the sum of premium

$$\pi(S) = (1 + \theta)ES = 160(1 + \theta).$$

According to the request, we have

$P(S \leq (1 + \theta)ES) = 0.95$, that is

$$P\left(\frac{S - ES}{\sqrt{VarS}} \leq \frac{\theta ES}{\sqrt{VarS}}\right) = P\left(\frac{S - ES}{\sqrt{VarS}} \leq 10\theta\right) = 0.95.$$

One can approximately regard $\frac{S-ES}{\sqrt{VarS}}$ as a standard normal variable. We have $10\theta = 1.645$, that is $\theta = 0.1645$.