

3.3 Characteristic functions

3.3.1 Definitions

3.3 Characteristic functions

3.3.1 Definitions

3.3 Characteristic functions

3.3.1 Definitions

3.3 Characteristic functions

3.3.1 Definitions

Definition 1 Suppose that ξ and η are real random variables, we call $\zeta = \xi + i\eta$ a complex random variable, where $i^2 = -1$. We call $E\zeta = E\xi + iE\eta$ the expectation of ζ .

3.3 Characteristic functions

3.3.1 Definitions

3.3 Characteristic functions

3.3.1 Definitions

Definition 1 Suppose that ξ and η are real random variables, we call $\zeta = \xi + i\eta$ a complex random variable, where $i^2 = -1$. We call $E\zeta = E\xi + iE\eta$ the expectation of ζ .

$E\zeta$ possesses properties similar to that of a real mathematical expectation.

3.3 Characteristic functions

3.3.1 Definitions

Modulus inequality: $|E\zeta| \leq E|\zeta|$.

3.3 Characteristic functions

3.3.1 Definitions

Modulus inequality: $|E\zeta| \leq E|\zeta|$.

证. 记 $\zeta = \xi + i\eta$. 则 $|E\zeta| = \sqrt{(E\xi)^2 + (E\eta)^2}$,
 $E|\zeta| = E\sqrt{\xi^2 + \eta^2}$.

3.3 Characteristic functions

3.3.1 Definitions

Modulus inequality: $|E\zeta| \leq E|\zeta|$.

证. 记 $\zeta = \xi + i\eta$. 则 $|E\zeta| = \sqrt{(E\xi)^2 + (E\eta)^2}$,
 $E|\zeta| = E\sqrt{\xi^2 + \eta^2}$. 设 a, b 为实数, 由初等不等式

$$|a\xi + b\eta| \leq \sqrt{a^2 + b^2} \cdot \sqrt{\xi^2 + \eta^2}$$

得

3.3 Characteristic functions

3.3.1 Definitions

Modulus inequality: $|E\zeta| \leq E|\zeta|$.

证. 记 $\zeta = \xi + i\eta$. 则 $|E\zeta| = \sqrt{(E\xi)^2 + (E\eta)^2}$,
 $E|\zeta| = E\sqrt{\xi^2 + \eta^2}$. 设 a, b 为实数, 由初等不等式

$$|a\xi + b\eta| \leq \sqrt{a^2 + b^2} \cdot \sqrt{\xi^2 + \eta^2}$$

得

$$|aE\xi + bE\eta| \leq E|a\xi + b\eta| \leq \sqrt{a^2 + b^2} \cdot E\sqrt{\xi^2 + \eta^2}.$$

取 $a = E\xi, b = E\eta$ 得

3.3 Characteristic functions

3.3.1 Definitions

$$(E\xi)^2 + (E\eta)^2 \leq \sqrt{(E\xi)^2 + (E\eta)^2} \cdot E\sqrt{\xi^2 + \eta^2}.$$

所以

$$\sqrt{(E\xi)^2 + (E\eta)^2} \leq E\sqrt{\xi^2 + \eta^2}.$$

即

$$|E\zeta| \leq E|\zeta|.$$

3.3 Characteristic functions

3.3.1 Definitions

Definition 2 Suppose ξ is a real random variable, we call

$$f(t) = Ee^{it\xi}, \quad -\infty < t < \infty$$

the characteristic function of ξ .

3.3 Characteristic functions

3.3.1 Definitions

Definition 2 Suppose ξ is a real random variable, we call

$$f(t) = Ee^{it\xi}, \quad -\infty < t < \infty$$

the characteristic function of ξ .

Notice $|e^{it\xi}| = 1$. $Ee^{it\xi}$ exists for all t .

3.3 Characteristic functions

3.3.1 Definitions

Definition 2 Suppose ξ is a real random variable, we call

$$f(t) = Ee^{it\xi}, \quad -\infty < t < \infty$$

the characteristic function of ξ .

Notice $|e^{it\xi}| = 1$. $Ee^{it\xi}$ exists for all t .

$$f(t) = \int_{-\infty}^{\infty} e^{itx} dF(x).$$

3.3 Characteristic functions

3.3.1 Definitions

If ξ is a discrete random variable with $P(\xi = x_n) = p_n$, then

$$f(t) = \sum_{n=1}^{\infty} p_n e^{itx_n}, \quad -\infty < t < \infty.$$

If ξ is a continuous random variable with the density function $p(x)$, then

$$f(t) = \int_{-\infty}^{\infty} e^{itx} p(x) dx, \quad -\infty < t < \infty,$$

which is just the Fourier transformation of $p(x)$.

3.3 Characteristic functions

3.3.1 Definitions

Example 1. The characteristic function of the degenerate distribution $P(\xi = c) = 1$ is

$$f(t) =$$

3.3 Characteristic functions

3.3.1 Definitions

Example 1. The characteristic function of the degenerate distribution $P(\xi = c) = 1$ is

$$f(t) = e^{ict}, \quad -\infty < t < \infty.$$

3.3 Characteristic functions

3.3.1 Definitions

Example 1. The characteristic function of the degenerate distribution $P(\xi = c) = 1$ is

$$f(t) = e^{ict}, \quad -\infty < t < \infty.$$

Example 2. The characteristic function of the binomial distribution $B(n, p)$ is

$$f(t) = \sum_{k=0}^n \binom{n}{k} p^k q^{n-k} e^{itk}$$

3.3 Characteristic functions

3.3.1 Definitions

Example 1. The characteristic function of the degenerate distribution $P(\xi = c) = 1$ is

$$f(t) = e^{ict}, \quad -\infty < t < \infty.$$

Example 2. The characteristic function of the binomial distribution $B(n, p)$ is

$$f(t) = \sum_{k=0}^n \binom{n}{k} p^k q^{n-k} e^{itk} = \sum_{k=0}^n \binom{n}{k} (pe^{it})^k q^{n-k}$$

3.3 Characteristic functions

3.3.1 Definitions

Example 1. The characteristic function of the degenerate distribution $P(\xi = c) = 1$ is

$$f(t) = e^{ict}, \quad -\infty < t < \infty.$$

Example 2. The characteristic function of the binomial distribution $B(n, p)$ is

$$\begin{aligned} f(t) &= \sum_{k=0}^n \binom{n}{k} p^k q^{n-k} e^{itk} = \sum_{k=0}^n \binom{n}{k} (pe^{it})^k q^{n-k} \\ &= (pe^{it} + q)^n, \quad p + q = 1. \end{aligned}$$

3.3 Characteristic functions

3.3.1 Definitions

Example 3. The characteristic function of the Poisson distribution $P(\lambda)$ is

$$f(t) = \sum_{k=0}^{\infty} \frac{\lambda^k}{k!} e^{-\lambda} e^{itk}$$

3.3 Characteristic functions

3.3.1 Definitions

Example 3. The characteristic function of the Poisson distribution $P(\lambda)$ is

$$\begin{aligned} f(t) &= \sum_{k=0}^{\infty} \frac{\lambda^k}{k!} e^{-\lambda} e^{itk} \\ &= \sum_{k=0}^{\infty} \frac{(\lambda e^{it})^k}{k!} e^{-\lambda} = e^{\lambda(e^{it}-1)}. \end{aligned}$$

3.3 Characteristic functions

3.3.1 Definitions

Example 3. The characteristic function of the Poisson distribution $P(\lambda)$ is

$$\begin{aligned} f(t) &= \sum_{k=0}^{\infty} \frac{\lambda^k}{k!} e^{-\lambda} e^{itk} \\ &= \sum_{k=0}^{\infty} \frac{(\lambda e^{it})^k}{k!} e^{-\lambda} = e^{\lambda(e^{it}-1)}. \end{aligned}$$

Example 4. The characteristic function of the uniform distribution $U[a, b]$ is

$$f(t) = \int_a^b \frac{1}{b-a} e^{itx} dx$$

3.3 Characteristic functions

3.3.1 Definitions

Example 3. The characteristic function of the Poisson distribution $P(\lambda)$ is

$$\begin{aligned} f(t) &= \sum_{k=0}^{\infty} \frac{\lambda^k}{k!} e^{-\lambda} e^{itk} \\ &= \sum_{k=0}^{\infty} \frac{(\lambda e^{it})^k}{k!} e^{-\lambda} = e^{\lambda(e^{it}-1)}. \end{aligned}$$

Example 4. The characteristic function of the uniform distribution $U[a, b]$ is

$$f(t) = \int_a^b \frac{1}{b-a} e^{itx} dx = \frac{e^{itb} - e^{ita}}{i(b-a)t}.$$

3.3 Characteristic functions

3.3.1 Definitions

Example 5. The characteristic function of the normal distribution $N(a, \sigma^2)$ is

$$f(t) = \frac{1}{\sqrt{2\pi}\sigma} \int_{-\infty}^{\infty} e^{itx - \frac{(x-a)^2}{2\sigma^2}} dx$$

3.3 Characteristic functions

3.3.1 Definitions

Example 5. The characteristic function of the normal distribution $N(a, \sigma^2)$ is

$$\begin{aligned} f(t) &= \frac{1}{\sqrt{2\pi}\sigma} \int_{-\infty}^{\infty} e^{itx - \frac{(x-a)^2}{2\sigma^2}} dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{ita + it\sigma y - \frac{y^2}{2}} dy \end{aligned}$$

3.3 Characteristic functions

3.3.1 Definitions

Example 5. The characteristic function of the normal distribution $N(a, \sigma^2)$ is

$$\begin{aligned} f(t) &= \frac{1}{\sqrt{2\pi}\sigma} \int_{-\infty}^{\infty} e^{itx - \frac{(x-a)^2}{2\sigma^2}} dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{ita + it\sigma y - \frac{y^2}{2}} dy \\ &= e^{iat - \frac{\sigma^2 t^2}{2}} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{(y-it\sigma)^2}{2}} dy \end{aligned}$$

3.3 Characteristic functions

3.3.1 Definitions

Example 5. The characteristic function of the normal distribution $N(a, \sigma^2)$ is

$$\begin{aligned} f(t) &= \frac{1}{\sqrt{2\pi}\sigma} \int_{-\infty}^{\infty} e^{itx - \frac{(x-a)^2}{2\sigma^2}} dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{ita + it\sigma y - \frac{y^2}{2}} dy \\ &= e^{iat - \frac{\sigma^2 t^2}{2}} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{(y-it\sigma)^2}{2}} dy \\ &= e^{iat - \frac{\sigma^2 t^2}{2}} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{y^2}{2}} dy \quad (\text{围道积分}) \end{aligned}$$

3.3 Characteristic functions


3.3.1 Definitions

Example 5. The characteristic function of the normal distribution $N(a, \sigma^2)$ is

$$\begin{aligned} f(t) &= \frac{1}{\sqrt{2\pi}\sigma} \int_{-\infty}^{\infty} e^{itx - \frac{(x-a)^2}{2\sigma^2}} dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{ita + it\sigma y - \frac{y^2}{2}} dy \\ &= e^{iat - \frac{\sigma^2 t^2}{2}} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{(y-it\sigma)^2}{2}} dy \\ &= e^{iat - \frac{\sigma^2 t^2}{2}} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{y^2}{2}} dy \quad (\text{围道积分}) \\ &= e^{iat - \frac{\sigma^2 t^2}{2}}. \end{aligned}$$

3.3 Characteristic functions

3.3.1 Definitions



3.3 Characteristic functions

3.3.1 Definitions

Solution (2): Let $\eta = (\xi - a)/\sigma$. Then $\eta \sim N(0, 1)$ and

$$f(t) = Ee^{it(a+\sigma\eta)} = e^{ita} f_{\eta}(\sigma t).$$

3.3 Characteristic functions

3.3.1 Definitions

Solution (2): Let $\eta = (\xi - a)/\sigma$. Then $\eta \sim N(0, 1)$ and

$$f(t) = Ee^{it(a+\sigma\eta)} = e^{ita} f_{\eta}(\sigma t).$$

So it is enough to show that $f_{\eta}(t) = e^{-\frac{t^2}{2}}$.

3.3 Characteristic functions

3.3.1 Definitions

It is obvious that

$$f_{\eta}(t) = Ee^{it\eta} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{itx} e^{-\frac{x^2}{2}} dx$$

3.3 Characteristic functions

3.3.1 Definitions

It is obvious that

$$\begin{aligned}f_{\eta}(t) &= Ee^{it\eta} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{itx} e^{-\frac{x^2}{2}} dx \\&= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \cos(tx) e^{-\frac{x^2}{2}} dx + i \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \sin(tx) e^{-\frac{x^2}{2}} dx \\&= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \cos(tx) e^{-\frac{x^2}{2}} dx.\end{aligned}$$

So

$$f'_{\eta}(t)$$

3.3 Characteristic functions

3.3.1 Definitions

It is obvious that

$$\begin{aligned}f_{\eta}(t) &= Ee^{it\eta} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{itx} e^{-\frac{x^2}{2}} dx \\&= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \cos(tx) e^{-\frac{x^2}{2}} dx + i \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \sin(tx) e^{-\frac{x^2}{2}} dx \\&= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \cos(tx) e^{-\frac{x^2}{2}} dx.\end{aligned}$$

So

$$f'_{\eta}(t) = -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} x \sin(tx) e^{-\frac{x^2}{2}} dx$$

3.3 Characteristic functions

3.3.1 Definitions

It is obvious that

$$\begin{aligned}f_{\eta}(t) &= Ee^{it\eta} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{itx} e^{-\frac{x^2}{2}} dx \\&= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \cos(tx) e^{-\frac{x^2}{2}} dx + i \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \sin(tx) e^{-\frac{x^2}{2}} dx \\&= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \cos(tx) e^{-\frac{x^2}{2}} dx.\end{aligned}$$

So

$$\begin{aligned}f'_{\eta}(t) &= -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} x \sin(tx) e^{-\frac{x^2}{2}} dx \\&= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \sin(tx) de^{-\frac{x^2}{2}}\end{aligned}$$

3.3 Characteristic functions

3.3.1 Definitions

It is obvious that

$$\begin{aligned}f_{\eta}(t) &= Ee^{it\eta} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{itx} e^{-\frac{x^2}{2}} dx \\&= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \cos(tx) e^{-\frac{x^2}{2}} dx + i \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \sin(tx) e^{-\frac{x^2}{2}} dx \\&= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \cos(tx) e^{-\frac{x^2}{2}} dx.\end{aligned}$$

So

$$\begin{aligned}f'_{\eta}(t) &= -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} x \sin(tx) e^{-\frac{x^2}{2}} dx \\&= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \sin(tx) d e^{-\frac{x^2}{2}} \\&= -t \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \cos(tx) e^{-\frac{x^2}{2}} dx = -t f_{\eta}(t)\end{aligned}$$

3.3 Characteristic functions

3.3.1 Definitions

Next we need to solve the differential equation

$$f'_\eta(t) + tf_\eta(t) = 0.$$

3.3 Characteristic functions

3.3.1 Definitions

Next we need to solve the differential equation

$$f'_\eta(t) + tf_\eta(t) = 0.$$

We have

$$\frac{d}{dt} \left(f_\eta(t) e^{\frac{t^2}{2}} \right) = f'_\eta(t) e^{\frac{t^2}{2}} + t f_\eta(t) e^{\frac{t^2}{2}} = 0.$$

3.3 Characteristic functions

3.3.1 Definitions

Next we need to solve the differential equation

$$f'_\eta(t) + tf_\eta(t) = 0.$$

We have

$$\frac{d}{dt} \left(f_\eta(t) e^{\frac{t^2}{2}} \right) = f'_\eta(t) e^{\frac{t^2}{2}} + t f_\eta(t) e^{\frac{t^2}{2}} = 0.$$

Hence

$$f_\eta(t) e^{\frac{t^2}{2}} = C$$

3.3 Characteristic functions

3.3.1 Definitions

Next we need to solve the differential equation

$$f'_\eta(t) + tf_\eta(t) = 0.$$

We have

$$\frac{d}{dt} \left(f_\eta(t) e^{\frac{t^2}{2}} \right) = f'_\eta(t) e^{\frac{t^2}{2}} + t f_\eta(t) e^{\frac{t^2}{2}} = 0.$$

Hence

$$f_\eta(t) e^{\frac{t^2}{2}} = C = f_\eta(0) = 1.$$

So

3.3 Characteristic functions

3.3.1 Definitions

Next we need to solve the differential equation

$$f'_\eta(t) + tf_\eta(t) = 0.$$

We have

$$\frac{d}{dt} \left(f_\eta(t) e^{\frac{t^2}{2}} \right) = f'_\eta(t) e^{\frac{t^2}{2}} + t f_\eta(t) e^{\frac{t^2}{2}} = 0.$$

Hence

$$f_\eta(t) e^{\frac{t^2}{2}} = C = f_\eta(0) = 1.$$

So

$$f_\eta(t) = e^{-\frac{t^2}{2}}.$$

3.3 Characteristic functions

3.3.1 Definitions

Solution (3): Let $\eta = (\xi - a)/\sigma$. Then $\eta \sim N(0, 1)$ and

$$f(t) = Ee^{it(a+\sigma\eta)} = e^{ita} f_{\eta}(\sigma t).$$

3.3 Characteristic functions

3.3.1 Definitions

Solution (3): Let $\eta = (\xi - a)/\sigma$. Then $\eta \sim N(0, 1)$ and

$$f(t) = Ee^{it(a+\sigma\eta)} = e^{ita} f_{\eta}(\sigma t).$$

So it is enough to show that $f_{\eta}(t) = e^{-\frac{t^2}{2}}$.

3.3 Characteristic functions

3.3.1 Definitions

$$f_{\eta}(t) = Ee^{it\eta} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{itx} e^{-\frac{x^2}{2}} dx$$

3.3 Characteristic functions

3.3.1 Definitions

$$\begin{aligned}f_{\eta}(t) &= Ee^{it\eta} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{itx} e^{-\frac{x^2}{2}} dx \\&= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \sum_{n=0}^{\infty} \frac{(itx)^n}{n!} e^{-\frac{x^2}{2}} dx \\&= \sum_{n=0}^{\infty} \frac{(it)^n}{n!} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} x^n e^{-\frac{x^2}{2}} dx\end{aligned}$$

3.3 Characteristic functions

3.3.1 Definitions

$$\begin{aligned}f_{\eta}(t) &= Ee^{it\eta} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{itx} e^{-\frac{x^2}{2}} dx \\&= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \sum_{n=0}^{\infty} \frac{(itx)^n}{n!} e^{-\frac{x^2}{2}} dx \\&= \sum_{n=0}^{\infty} \frac{(it)^n}{n!} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} x^n e^{-\frac{x^2}{2}} dx \\&= \sum_{n=0}^{\infty} \frac{(it)^{2n}}{(2n)!} E\eta^{2n} = \sum_{n=0}^{\infty} \frac{(it)^{2n}}{(2n)!} (2n-1)!!\end{aligned}$$

3.3 Characteristic functions

3.3.1 Definitions

$$\begin{aligned}f_{\eta}(t) &= Ee^{it\eta} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{itx} e^{-\frac{x^2}{2}} dx \\&= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \sum_{n=0}^{\infty} \frac{(itx)^n}{n!} e^{-\frac{x^2}{2}} dx \\&= \sum_{n=0}^{\infty} \frac{(it)^n}{n!} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} x^n e^{-\frac{x^2}{2}} dx \\&= \sum_{n=0}^{\infty} \frac{(it)^{2n}}{(2n)!} E\eta^{2n} = \sum_{n=0}^{\infty} \frac{(it)^{2n}}{(2n)!} (2n-1)!! \\&= \sum_{n=0}^{\infty} \frac{(it)^{2n}}{(2n)!} \frac{(2n)!}{2^n n!} = \sum_{n=0}^{\infty} \left(-\frac{t^2}{2}\right)^n \frac{1}{n!} = e^{-\frac{t^2}{2}}.\end{aligned}$$

3.3 Characteristic functions

3.3.1 Definitions

Example 6. The characteristic function of the Cauchy distribution is

$$f(t) = \int_{-\infty}^{\infty} e^{itx} \frac{1}{\pi(1+x^2)} dx$$

3.3 Characteristic functions

3.3.1 Definitions

Example 6. The characteristic function of the Cauchy distribution is

$$f(t) = \int_{-\infty}^{\infty} e^{itx} \frac{1}{\pi(1+x^2)} dx = e^{-|t|}.$$

3.3 Characteristic functions

3.3.1 Definitions

Example 6. The characteristic function of the Cauchy distribution is


$$f(t) = \int_{-\infty}^{\infty} e^{itx} \frac{1}{\pi(1+x^2)} dx = e^{-|t|}.$$

In fact, when $t > 0$,

$$\begin{aligned} & \int_{-R}^R e^{itx} \frac{1}{\pi(1+x^2)} dx + \int_{\text{semicircle}} \frac{e^{itz}}{\pi(1+z^2)} dz \\ &= 2\pi i \operatorname{Res} \left(\frac{e^{itz}}{\pi(1+z^2)} \text{ at } i \right) \\ &= 2\pi i (z-i) \frac{e^{itz}}{\pi(1+iz)(1-iz)} \Big|_{z=i} = e^{-t}. \end{aligned}$$

3.3 Characteristic functions

3.3.1 Definitions



3.3 Characteristic functions

3.3.1 Definitions

$$\begin{aligned} \left| \int_{\text{semicircle}} \frac{e^{itz}}{\pi(1+z^2)} dz \right| &\leq \int_{\text{semicircle}} \frac{1}{\pi(R^2-1)} dz \\ &= \frac{\pi R}{\pi(R^2-1)} \rightarrow 0. \end{aligned}$$

3.3 Characteristic functions

3.3.2 Properties

3.3.2 Properties

$$\textcircled{1} \quad |f(t)| \leq f(0) = 1; \quad f(-t) = \overline{f(t)}.$$

Proof.

3.3.2 Properties

$$\textcircled{1} \quad |f(t)| \leq f(0) = 1; \quad f(-t) = \overline{f(t)}.$$

Proof. Obviously,

$$|f(t)| = \left| \int_{-\infty}^{\infty} e^{itx} dF(x) \right| \leq \int_{-\infty}^{\infty} |e^{itx}| dF(x) = 1$$

and

3.3.2 Properties

$$\textcircled{1} \quad |f(t)| \leq f(0) = 1; \quad f(-t) = \overline{f(t)}.$$

Proof. Obviously,

$$|f(t)| = \left| \int_{-\infty}^{\infty} e^{itx} dF(x) \right| \leq \int_{-\infty}^{\infty} |e^{itx}| dF(x) = 1$$

and

$$f(0) = \int_{-\infty}^{\infty} e^{i0x} dF(x) = 1.$$

3.3 Characteristic functions

3.3.2 Properties

Also,

$$f(-t) = \int_{-\infty}^{\infty} e^{-itx} dF(x)$$

3.3 Characteristic functions

3.3.2 Properties

Also,

$$f(-t) = \int_{-\infty}^{\infty} e^{-itx} dF(x) = \int_{-\infty}^{\infty} \overline{e^{itx}} dF(x)$$

3.3 Characteristic functions

3.3.2 Properties

Also,

$$\begin{aligned} f(-t) &= \int_{-\infty}^{\infty} e^{-itx} dF(x) = \int_{-\infty}^{\infty} \overline{e^{itx}} dF(x) \\ &= \overline{\int_{-\infty}^{\infty} e^{itx} dF(x)} = \overline{f(t)}, \end{aligned}$$

as desired.

3.3 Characteristic functions

3.3.2 Properties

2 $f(t)$ is uniformly continuous on $(-\infty, \infty)$.

3.3 Characteristic functions

3.3.2 Properties

2 $f(t)$ is uniformly continuous on $(-\infty, \infty)$.

Proof.

3.3 Characteristic functions

3.3.2 Properties

② $f(t)$ is uniformly continuous on $(-\infty, \infty)$.

Proof. For any $t \in (-\infty, \infty)$ and $\varepsilon > 0$,

② $f(t)$ is uniformly continuous on $(-\infty, \infty)$.

Proof. For any $t \in (-\infty, \infty)$ and $\varepsilon > 0$, it follows that

$$\begin{aligned} & |f(t+h) - f(t)| \\ = & \left| \int_{-\infty}^{\infty} (e^{itx} e^{ihx} - e^{itx}) dF(x) \right| \end{aligned}$$

3.3 Characteristic functions

3.3.2 Properties

2 $f(t)$ is uniformly continuous on $(-\infty, \infty)$.

Proof. For any $t \in (-\infty, \infty)$ and $\varepsilon > 0$, it follows that

$$\begin{aligned} & |f(t+h) - f(t)| \\ = & \left| \int_{-\infty}^{\infty} (e^{itx} e^{ihx} - e^{itx}) dF(x) \right| \\ \leq & \int_{-\infty}^{\infty} |e^{ihx} - 1| dF(x) \end{aligned}$$

3.3 Characteristic functions

3.3.2 Properties

② $f(t)$ is uniformly continuous on $(-\infty, \infty)$.

Proof. For any $t \in (-\infty, \infty)$ and $\varepsilon > 0$, it follows that

$$\begin{aligned} & |f(t+h) - f(t)| \\ = & \left| \int_{-\infty}^{\infty} (e^{itx} e^{ihx} - e^{itx}) dF(x) \right| \\ \leq & \int_{-\infty}^{\infty} |e^{ihx} - 1| dF(x) \\ \leq & \left(\int_{|x| \geq A} + \int_{|x| < A} \right) |e^{ihx} - 1| dF(x). \end{aligned}$$

3.3 Characteristic functions

3.3.2 Properties

Note that $|e^{ihx} - 1| \leq 2$ and

$$\begin{aligned}|e^{ihx} - 1| &= |e^{i\frac{h}{2}x}| |e^{i\frac{h}{2}x} - e^{-i\frac{h}{2}x}| = 2 \left| \sin \frac{hx}{2} \right| \\ &\leq |hx| \leq A|h|, \quad \text{when } |x| \leq A.\end{aligned}$$

3.3 Characteristic functions

3.3.2 Properties

Note that $|e^{ihx} - 1| \leq 2$ and

$$\begin{aligned}|e^{ihx} - 1| &= |e^{i\frac{h}{2}x}| |e^{i\frac{h}{2}x} - e^{-i\frac{h}{2}x}| = 2 \left| \sin \frac{hx}{2} \right| \\ &\leq |hx| \leq A|h|, \quad \text{when } |x| \leq A.\end{aligned}$$

We have

$$\begin{aligned}|f(t+h) - f(t)| &\leq 2 \int_{|x| \geq A} dF(x) + A|h| \int_{|x| < A} dF(x) \\ &\leq 2 \int_{|x| \geq A} dF(x) + |h|A.\end{aligned}$$

3.3 Characteristic functions

3.3.2 Properties

Note that $|e^{ihx} - 1| \leq 2$ and

$$\begin{aligned}|e^{ihx} - 1| &= |e^{i\frac{h}{2}x}| |e^{i\frac{h}{2}x} - e^{-i\frac{h}{2}x}| = 2 \left| \sin \frac{hx}{2} \right| \\ &\leq |hx| \leq A|h|, \quad \text{when } |x| \leq A.\end{aligned}$$

We have

$$\begin{aligned}|f(t+h) - f(t)| &\leq 2 \int_{|x| \geq A} dF(x) + A|h| \int_{|x| < A} dF(x) \\ &\leq 2 \int_{|x| \geq A} dF(x) + |h|A.\end{aligned}$$

Choose A such that $\int_{|x| \geq A} dF(x) < \epsilon/4$.

3.3 Characteristic functions

3.3.2 Properties

Note that $|e^{ihx} - 1| \leq 2$ and

$$\begin{aligned}|e^{ihx} - 1| &= |e^{i\frac{h}{2}x}| |e^{i\frac{h}{2}x} - e^{-i\frac{h}{2}x}| = 2 \left| \sin \frac{hx}{2} \right| \\ &\leq |hx| \leq A|h|, \quad \text{when } |x| \leq A.\end{aligned}$$

We have

$$\begin{aligned}|f(t+h) - f(t)| &\leq 2 \int_{|x| \geq A} dF(x) + A|h| \int_{|x| < A} dF(x) \\ &\leq 2 \int_{|x| \geq A} dF(x) + |h|A.\end{aligned}$$

Choose A such that $\int_{|x| \geq A} dF(x) < \epsilon/4$. And then take $\delta = \epsilon/(2A)$. Consequently, $|f(t+h) - f(t)| < \epsilon$ for all t whenever $|h| < \delta$.

3.3 Characteristic functions

3.3.2 Properties

- ③ $f(t)$ is non-negative definite, i.e., for an arbitrary integer n , any real numbers t_1, \dots, t_n and complex numbers $\lambda_1, \dots, \lambda_n$, it follows

$$\sum_{k=1}^n \sum_{j=1}^n f(t_k - t_j) \lambda_k \overline{\lambda_j} \geq 0.$$

Proof.

3.3 Characteristic functions

3.3.2 Properties

Proof.

$$\begin{aligned}& \sum_{k=1}^n \sum_{j=1}^n f(t_k - t_j) \lambda_k \overline{\lambda_j} \\&= \sum_{k=1}^n \sum_{j=1}^n \int_{-\infty}^{\infty} e^{i(t_k - t_j)x} dF(x) \lambda_k \overline{\lambda_j} \\&= \int_{-\infty}^{\infty} \left(\sum_{k=1}^n e^{it_k x} \lambda_k \right) \left(\sum_{j=1}^n e^{-it_j x} \overline{\lambda_j} \right) dF(x)\end{aligned}$$

3.3 Characteristic functions

3.3.2 Properties

Proof.

$$\begin{aligned}& \sum_{k=1}^n \sum_{j=1}^n f(t_k - t_j) \lambda_k \overline{\lambda_j} \\&= \sum_{k=1}^n \sum_{j=1}^n \int_{-\infty}^{\infty} e^{i(t_k - t_j)x} dF(x) \lambda_k \overline{\lambda_j} \\&= \int_{-\infty}^{\infty} \left(\sum_{k=1}^n e^{it_k x} \lambda_k \right) \overline{\left(\sum_{j=1}^n e^{-it_j x} \overline{\lambda_j} \right)} dF(x) \\&= \int_{-\infty}^{\infty} \left(\sum_{k=1}^n e^{it_k x} \lambda_k \right) \left(\sum_{j=1}^n e^{it_j x} \lambda_j \right) dF(x)\end{aligned}$$

3.3 Characteristic functions

3.3.2 Properties

Proof.

$$\begin{aligned}
 & \sum_{k=1}^n \sum_{j=1}^n f(t_k - t_j) \lambda_k \overline{\lambda_j} \\
 = & \sum_{k=1}^n \sum_{j=1}^n \int_{-\infty}^{\infty} e^{i(t_k - t_j)x} dF(x) \lambda_k \overline{\lambda_j} \\
 = & \int_{-\infty}^{\infty} \left(\sum_{k=1}^n e^{it_k x} \lambda_k \right) \overline{\left(\sum_{j=1}^n e^{-it_j x} \overline{\lambda_j} \right)} dF(x) \\
 = & \int_{-\infty}^{\infty} \left(\sum_{k=1}^n e^{it_k x} \lambda_k \right) \left(\sum_{j=1}^n e^{it_j x} \lambda_j \right) dF(x) \\
 = & \int_{-\infty}^{\infty} \left| \sum_{k=1}^n e^{it_k x} \lambda_k \right|^2 dF(x) \geq 0.
 \end{aligned}$$

3.3 Characteristic functions

3.3.2 Properties

Bochner-Khinchine Theorem.

The function $f(t)$ is a characteristic function if and only if $f(t)$ is non-negative definite, continuous and $f(0) = 1$.

3.3 Characteristic functions

3.3.2 Properties

④ Assume that ξ_1, \dots, ξ_n are indept., then

$$f_{\xi_1 + \xi_2 + \dots + \xi_n}(t) = f_{\xi_1}(t) f_{\xi_2}(t) \cdots f_{\xi_n}(t).$$

(Proof?)

3.3 Characteristic functions

3.3.2 Properties

④ Assume that ξ_1, \dots, ξ_n are indept., then

$$f_{\xi_1 + \xi_2 + \dots + \xi_n}(t) = f_{\xi_1}(t) f_{\xi_2}(t) \cdots f_{\xi_n}(t).$$

(Proof?)

⑤ If $E\xi^n$ exists, then $f(t)$ is differentiable of n orders, and when $k \leq n$

$$f^{(k)}(0) = i^k E\xi^k.$$

3.3 Characteristic functions

3.3.2 Properties

④ Assume that ξ_1, \dots, ξ_n are indept., then

$$f_{\xi_1 + \xi_2 + \dots + \xi_n}(t) = f_{\xi_1}(t) f_{\xi_2}(t) \cdots f_{\xi_n}(t).$$

(Proof?)

⑤ If $E\xi^n$ exists, then $f(t)$ is differentiable of n orders, and when $k \leq n$

$$f^{(k)}(0) = i^k E\xi^k.$$

3.3 Characteristic functions

3.3.2 Properties

④ Assume that ξ_1, \dots, ξ_n are indept., then

$$f_{\xi_1+\xi_2+\dots+\xi_n}(t) = f_{\xi_1}(t)f_{\xi_2}(t)\cdots f_{\xi_n}(t).$$

(Proof?)

⑤ If $E\xi^n$ exists, then $f(t)$ is differentiable of n orders, and when $k \leq n$

$$f^{(k)}(0) = i^k E\xi^k.$$

In particular, when $E\xi^2$ exists, $E\xi = -if'(0)$,
 $E\xi^2 = -f''(0)$, $Var\xi = -f''(0) + [f'(0)]^2$.

3.3 Characteristic functions

3.3.2 Properties

Proof. Since

$$\left| \frac{d^k}{dt^k} e^{itx} \right| = |i^k x^k e^{itx}| = |x|^k,$$

3.3 Characteristic functions

3.3.2 Properties

Proof. Since

$$\left| \frac{d^k}{dt^k} e^{itx} \right| = |i^k x^k e^{itx}| = |x|^k,$$
$$\int_{-\infty}^{\infty} |x|^k dF(x) = E|\xi|^k < \infty,$$

Proof. Since

$$\left| \frac{d^k}{dt^k} e^{itx} \right| = |i^k x^k e^{itx}| = |x|^k,$$
$$\int_{-\infty}^{\infty} |x|^k dF(x) = E|\xi|^k < \infty,$$

then $\int_{-\infty}^{\infty} \frac{d^k}{dt^k} e^{itx} dF(x)$ uniformly converges in t .

3.3 Characteristic functions

3.3.2 Properties

Proof. Since

$$\left| \frac{d^k}{dt^k} e^{itx} \right| = |i^k x^k e^{itx}| = |x|^k,$$
$$\int_{-\infty}^{\infty} |x|^k dF(x) = E|\xi|^k < \infty,$$

then $\int_{-\infty}^{\infty} \frac{d^k}{dt^k} e^{itx} dF(x)$ uniformly converges in t . So, $f^{(k)}(t)$ exists and

Proof. Since

$$\left| \frac{d^k}{dt^k} e^{itx} \right| = |i^k x^k e^{itx}| = |x|^k,$$
$$\int_{-\infty}^{\infty} |x|^k dF(x) = E|\xi|^k < \infty,$$

then $\int_{-\infty}^{\infty} \frac{d^k}{dt^k} e^{itx} dF(x)$ uniformly converges in t . So, $f^{(k)}(t)$ exists and

$$f^{(k)}(t) = \int_{-\infty}^{\infty} \frac{d^k}{dt^k} e^{itx} dF(x) = i^k \int_{-\infty}^{\infty} x^k e^{itx} dF(x),$$

Proof. Since

$$\left| \frac{d^k}{dt^k} e^{itx} \right| = |i^k x^k e^{itx}| = |x|^k,$$
$$\int_{-\infty}^{\infty} |x|^k dF(x) = E|\xi|^k < \infty,$$

then $\int_{-\infty}^{\infty} \frac{d^k}{dt^k} e^{itx} dF(x)$ uniformly converges in t . So, $f^{(k)}(t)$ exists and

$$f^{(k)}(t) = \int_{-\infty}^{\infty} \frac{d^k}{dt^k} e^{itx} dF(x) = i^k \int_{-\infty}^{\infty} x^k e^{itx} dF(x),$$

$$f^{(k)}(0) = i^k \int_{-\infty}^{\infty} x^k dF(x) = i^k E\xi^k.$$

3.3 Characteristic functions

3.3.2 Properties

反过来, 若 n 为偶数, 且 $f^{(n)}(0)$ 存在, 则 $E\xi^n$ 存在.

3.3 Characteristic functions

3.3.2 Properties

反过来, 若 n 为偶数, 且 $f^{(n)}(0)$ 存在, 则 $E\xi^n$ 存在.

Proof. 我们用数学归纳法来证明. 当 $n = 2$ 时,

$$\begin{aligned} f''(0) &= \lim_{h \rightarrow 0} \frac{f(h) - 2f(0) + f(-h)}{h^2} \\ &= \lim_{h \rightarrow 0} \int_{-\infty}^{\infty} \frac{e^{ihx} - 2 + e^{-ihx}}{h^2} dF(x) \\ &= - \lim_{h \rightarrow 0} \int_{-\infty}^{\infty} 2 \frac{1 - \cos hx}{h^2} dF(x). \end{aligned}$$

3.3 Characteristic functions

3.3.2 Properties

反过来, 若 n 为偶数, 且 $f^{(n)}(0)$ 存在, 则 $E\xi^n$ 存在.

Proof. 我们用数学归纳法来证明. 当 $n = 2$ 时,

$$\begin{aligned} f''(0) &= \lim_{h \rightarrow 0} \frac{f(h) - 2f(0) + f(-h)}{h^2} \\ &= \lim_{h \rightarrow 0} \int_{-\infty}^{\infty} \frac{e^{ihx} - 2 + e^{-ihx}}{h^2} dF(x) \\ &= - \lim_{h \rightarrow 0} \int_{-\infty}^{\infty} 2 \frac{1 - \cos hx}{h^2} dF(x). \end{aligned}$$

注意到, $0 \leq 2(1 - \cos hx)/h^2 \leq x^2$, 并

且 $\lim_{h \rightarrow 0} 2(1 - \cos hx)/h^2 = x^2$ 关于 x 在任一有限区间内一致成立.

3.3 Characteristic functions

3.3.2 Properties

因此对任意 $a > 0$ 有

$$\begin{aligned} -f''(0) &\geq \lim_{h \rightarrow 0} \int_{-a}^a 2 \frac{1 - \cos hx}{h^2} dF(x) \\ &= \int_{-a}^a \lim_{h \rightarrow 0} 2 \frac{1 - \cos hx}{h^2} dF(x) = \int_{-a}^a x^2 dF(x). \end{aligned}$$

令 $a \rightarrow \infty$ 得 $\int_{-\infty}^{\infty} x^2 dF(x) \leq -f''(0)$, 即 $E\xi^2$ 存在.

3.3 Characteristic functions

3.3.2 Properties

现设 $f^{(2k)}(0)$ 存在, 同时归纳假设 $E\xi^{2k-2}$ 也存在.
由第一部分结论, $f(t)$ 是 $2k-2$ 次可微的, 且

$$\begin{aligned} f^{(2k-2)}(t) &= i^{2k-2} \int_{-\infty}^{\infty} e^{itx} x^{2k-2} dF(x) \\ &= (-1)^{k-1} \int_{-\infty}^{\infty} e^{itx} x^{2k-2} dF(x). \end{aligned}$$

记 $G(y) = \int_{-\infty}^y x^{2k-2} dF(x)$, 其中 $G(\infty) = E\xi^{2k-2}$,
则 $G(y)/G(\infty)$ 为分布函数,

3.3 Characteristic functions

3.3.2 Properties

$G(y)/G(\infty)$ 的特征函数为

$$\begin{aligned} g(t) &= \frac{1}{G(\infty)} \int_{-\infty}^{\infty} e^{ity} dG(y) \\ &= \frac{1}{G(\infty)} \int_{-\infty}^{\infty} e^{ity} y^{2k-2} dF(y) = \frac{(-1)^{k-1}}{G(\infty)} f^{(2k-2)}(t). \end{aligned}$$

从而 $g''(0) = (-1)^{k-1} f^{(2k)}(0)/G(\infty)$ 存在. 由已证的 $n = 2$ 时的结论知

$$\frac{1}{G(\infty)} \int_{-\infty}^{\infty} x^{2k} dF(x) = \frac{1}{G(\infty)} \int_{-\infty}^{\infty} y^2 dG(y)$$

存在. 即 $E\xi^{2k}$ 存在, 结论得证.

3.3 Characteristic functions

3.3.2 Properties

6 Let $\eta = a\xi + b$, where a, b are arbitrary constants. Then

$$f_{\eta}(t) = e^{ibt} f(at).$$

3.3 Characteristic functions

3.3.2 Properties

6 Let $\eta = a\xi + b$, where a, b are arbitrary constants. Then

$$f_{\eta}(t) = e^{ibt} f(at).$$

Proof.

$$Ee^{i(a\xi+b)t} = Ee^{iat\xi} \cdot e^{ibt} = e^{ibt} f(at).$$

Example 7. Are the following functions characteristic functions of some random variables?

(1) $f(t) = \sin t$;

(2) $f(t) = \ln(e + |t|)$;

(3) $f(t) = 0$ when $t < 0$; $f(t) = 1$ when $t \geq 0$.

Solution.....

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

3.3.3 Inverse formula and uniqueness theorem

Theorem 1 (Inverse formula) Suppose that $f(t)$ is a c.f. corresponding to cdf $F(x)$. Let x_1, x_2 be two continuity points of $F(x)$, then

$$F(x_2) - F(x_1) = \lim_{T \rightarrow \infty} \frac{1}{2\pi} \int_{-T}^T \frac{e^{-itx_1} - e^{-itx_2}}{it} f(t) dt.$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Proof. Suppose $\xi \sim F(x)$. Without loss of generality, we assume that $x_1 < x_2$.

$$\frac{1}{2\pi} \int_{-T}^T \frac{e^{-itx_1} - e^{-itx_2}}{it} f(t) dt$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Proof. Suppose $\xi \sim F(x)$. Without loss of generality, we assume that $x_1 < x_2$.

$$\begin{aligned} & \frac{1}{2\pi} \int_{-T}^T \frac{e^{-itx_1} - e^{-itx_2}}{it} f(t) dt \\ = & \frac{1}{2\pi} \int_{-T}^T \int_{-\infty}^{\infty} \frac{e^{-itx_1} - e^{-itx_2}}{it} e^{itx} dF(x) dt \end{aligned}$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Proof. Suppose $\xi \sim F(x)$. Without loss of generality, we assume that $x_1 < x_2$.

$$\begin{aligned}
 & \frac{1}{2\pi} \int_{-T}^T \frac{e^{-itx_1} - e^{-itx_2}}{it} f(t) dt \\
 = & \frac{1}{2\pi} \int_{-T}^T \int_{-\infty}^{\infty} \frac{e^{-itx_1} - e^{-itx_2}}{it} e^{itx} dF(x) dt \\
 = & \frac{1}{2\pi} \int_{-\infty}^{\infty} \left\{ \int_0^T \left[\frac{e^{it(x-x_1)} - e^{-it(x-x_1)}}{it} \right. \right. \\
 & \quad \left. \left. - \frac{e^{it(x-x_2)} - e^{-it(x-x_2)}}{it} \right] dt \right\} dF(x)
 \end{aligned}$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Proof. Suppose $\xi \sim F(x)$. Without loss of generality, we assume that $x_1 < x_2$.

$$\begin{aligned}
 & \frac{1}{2\pi} \int_{-T}^T \frac{e^{-itx_1} - e^{-itx_2}}{it} f(t) dt \\
 = & \frac{1}{2\pi} \int_{-T}^T \int_{-\infty}^{\infty} \frac{e^{-itx_1} - e^{-itx_2}}{it} e^{itx} dF(x) dt \\
 = & \frac{1}{2\pi} \int_{-\infty}^{\infty} \left\{ \int_0^T \left[\frac{e^{it(x-x_1)} - e^{-it(x-x_1)}}{it} \right. \right. \\
 & \quad \left. \left. - \frac{e^{it(x-x_2)} - e^{-it(x-x_2)}}{it} \right] dt \right\} dF(x) \\
 = & \frac{1}{\pi} \int_{-\infty}^{\infty} \left\{ \int_0^T \left[\frac{\sin t(x-x_1)}{t} - \frac{\sin t(x-x_2)}{t} \right] dt \right\} dF(x)
 \end{aligned}$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Proof. Suppose $\xi \sim F(x)$. Without loss of generality, we assume that $x_1 < x_2$.

$$\begin{aligned}
 & \frac{1}{2\pi} \int_{-T}^T \frac{e^{-itx_1} - e^{-itx_2}}{it} f(t) dt \\
 = & \frac{1}{2\pi} \int_{-T}^T \int_{-\infty}^{\infty} \frac{e^{-itx_1} - e^{-itx_2}}{it} e^{itx} dF(x) dt \\
 = & \frac{1}{2\pi} \int_{-\infty}^{\infty} \left\{ \int_0^T \left[\frac{e^{it(x-x_1)} - e^{-it(x-x_1)}}{it} \right. \right. \\
 & \quad \left. \left. - \frac{e^{it(x-x_2)} - e^{-it(x-x_2)}}{it} \right] dt \right\} dF(x) \\
 = & \frac{1}{\pi} \int_{-\infty}^{\infty} \left\{ \int_0^T \left[\frac{\sin t(x-x_1)}{t} - \frac{\sin t(x-x_2)}{t} \right] dt \right\} dF(x) \\
 \triangleq & \int_{-\infty}^{\infty} g(T, x, x_1, x_2) dF(x) = Eg(T, \xi, x_1, x_2).
 \end{aligned}$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Notice

$$\int_0^T \frac{\sin at}{t} dt = \int_0^{Ta} \frac{\sin t}{t} dt$$
$$\rightarrow \operatorname{sgn}(a) \int_0^\infty \frac{\sin t}{t} dt = \begin{cases} \frac{\pi}{2}, & \text{if } a > 0, \\ 0, & \text{if } a = 0, \\ -\frac{\pi}{2}, & \text{if } a < 0. \end{cases}$$

and

$\int_0^x \frac{\sin t}{t} dt$ is a bounded function .

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

It follows that

$$\begin{aligned} & \lim_{T \rightarrow \infty} g(T, x, x_1, x_2) \\ = & \lim_{T \rightarrow \infty} \frac{1}{\pi} \left\{ \int_0^T \left[\frac{\sin t(x - x_1)}{t} - \frac{\sin t(x - x_2)}{t} \right] dt \right\} \\ = & \begin{cases} 0, & x < x_1 \text{ or } x > x_2, \\ \frac{1}{2}, & x = x_1 \text{ or } x = x_2, \\ 1, & x_1 < x < x_2. \end{cases} \end{aligned}$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

It follows that

$$\begin{aligned} & \lim_{T \rightarrow \infty} g(T, x, x_1, x_2) \\ = & \lim_{T \rightarrow \infty} \frac{1}{\pi} \left\{ \int_0^T \left[\frac{\sin t(x - x_1)}{t} - \frac{\sin t(x - x_2)}{t} \right] dt \right\} \\ = & \begin{cases} 0, & x < x_1 \text{ or } x > x_2, \\ \frac{1}{2}, & x = x_1 \text{ or } x = x_2, \\ 1, & x_1 < x < x_2. \end{cases} \triangleq g(x) \end{aligned}$$

and

$$|g(T, x, x_1, x_2)| < M.$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

It follows that

$$\begin{aligned} & \lim_{T \rightarrow \infty} \frac{1}{2\pi} \int_{-T}^T \frac{e^{-itx_1} - e^{-itx_2}}{it} f(t) dt \\ &= \lim_{T \rightarrow \infty} E g(T, \xi, x_1, x_2) \end{aligned}$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

It follows that

$$\begin{aligned} & \lim_{T \rightarrow \infty} \frac{1}{2\pi} \int_{-T}^T \frac{e^{-itx_1} - e^{-itx_2}}{it} f(t) dt \\ &= \lim_{T \rightarrow \infty} Eg(T, \xi, x_1, x_2) = Eg(\xi) \end{aligned}$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

It follows that

$$\begin{aligned} & \lim_{T \rightarrow \infty} \frac{1}{2\pi} \int_{-T}^T \frac{e^{-itx_1} - e^{-itx_2}}{it} f(t) dt \\ &= \lim_{T \rightarrow \infty} Eg(T, \xi, x_1, x_2) = Eg(\xi) \\ &= P(x_1 < \xi < x_2) + \frac{1}{2}(P(\xi = x_1) + P(\xi = x_2)) \end{aligned}$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

It follows that

$$\begin{aligned}& \lim_{T \rightarrow \infty} \frac{1}{2\pi} \int_{-T}^T \frac{e^{-itx_1} - e^{-itx_2}}{it} f(t) dt \\&= \lim_{T \rightarrow \infty} Eg(T, \xi, x_1, x_2) = Eg(\xi) \\&= P(x_1 < \xi < x_2) + \frac{1}{2}(P(\xi = x_1) + P(\xi = x_2)) \\&= F(x_2) - F(x_1).\end{aligned}$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Theorem 2. (Uniqueness) A distribution function can be uniquely determined by its characteristic function.

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Theorem 2. (Uniqueness) A distribution function can be uniquely determined by its characteristic function.

Proof.

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Theorem 2. (Uniqueness) A distribution function can be uniquely determined by its characteristic function.

Proof. By inverse formula, if $y < x$ are continuous points of $F(x)$, then

$$F(x) - F(y) = \lim_{T \rightarrow \infty} \frac{1}{2\pi} \int_{-T}^T \frac{e^{-ity} - e^{-itx}}{it} f(t) dt.$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Theorem 2. (Uniqueness) A distribution function can be uniquely determined by its characteristic function.

Proof. By inverse formula, if $y < x$ are continuous points of $F(x)$, then

$$F(x) - F(y) = \lim_{T \rightarrow \infty} \frac{1}{2\pi} \int_{-T}^T \frac{e^{-ity} - e^{-itx}}{it} f(t) dt.$$

Letting $y \rightarrow -\infty$ along continuity points of $F(x)$, we have

$$F(x) = \lim_{y \rightarrow -\infty} \lim_{T \rightarrow \infty} \frac{1}{2\pi} \int_{-T}^T \frac{e^{-ity} - e^{-itx}}{it} f(t) dt.$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Thus it is easy to see that $f(t)$ determines the value of $F(x)$ at its continuity points. As for the discontinuous points, in view of right continuity of $F(x)$, it suffices to take right limits along continuity points. The theorem is proved.

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Theorem 3. (Inverse Fourier transform) Suppose that $f(t)$ is a c.f. and $\int_{-\infty}^{\infty} |f(t)| dt < \infty$, then $F'(x)$ exists and is continuous. Moreover

$$F'(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-itx} f(t) dt.$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Proof. Since $f(t)$ is absolutely integrable and

$$\left| \frac{e^{-itx} - e^{-ity}}{it} \right| \leq |y - x|,$$

it follows that

$$\begin{aligned} F(y) - F(x) &= \lim_{T \rightarrow \infty} \frac{1}{2\pi} \int_{-T}^T \frac{e^{-itx} - e^{-ity}}{it} f(t) dt \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{e^{-itx} - e^{-ity}}{it} f(t) dt \triangleq H(x, y), \end{aligned}$$

whenever x, y are continuous points of $F(x)$.

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Also, $H(x, y)$ is a continuous function of (x, y) . So, $F(x)$ must be a continuous function and the above equality holds for all x, y .

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Also, $H(x, y)$ is a continuous function of (x, y) . So, $F(x)$ must be a continuous function and the above equality holds for all x, y .

Now, applying the dominated convergence theorem yields

$$F'(x) = \lim_{y \rightarrow x} \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{e^{-itx} - e^{-ity}}{it(y-x)} f(t) dt$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Also, $H(x, y)$ is a continuous function of (x, y) . So, $F(x)$ must be a continuous function and the above equality holds for all x, y .

Now, applying the dominated convergence theorem yields

$$\begin{aligned} F'(x) &= \lim_{y \rightarrow x} \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{e^{-itx} - e^{-ity}}{it(y-x)} f(t) dt \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \lim_{y \rightarrow x} \frac{e^{-itx} - e^{-ity}}{it(y-x)} f(t) dt \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-itx} f(t) dt. \end{aligned}$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

For same reason,

$$\lim_{y \rightarrow x} F'(y) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \lim_{y \rightarrow x} e^{-ity} f(t) dt = F'(x).$$

So, $F'(x)$ is continuous.

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Discrete random variables: Assume

$P(\xi = k) = p_k, k = 0, 1, 2, \dots$, then

$$f(t) = \sum_{k=0}^{\infty} p_k e^{itk}.$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Discrete random variables: Assume

$P(\xi = k) = p_k, k = 0, 1, 2, \dots$, then

$$f(t) = \sum_{k=0}^{\infty} p_k e^{itk}.$$

If $f(t)$ is given, then we can multiply both sides by e^{-itk} and integrate. Noting that

$$\int_0^{2\pi} e^{int} dt = \begin{cases} 2\pi, & n = 0, \\ 0, & n \neq 0, \end{cases}$$

we have

$$p_k = \frac{1}{2\pi} \int_0^{2\pi} e^{-itk} f(t) dt.$$

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Example 8. Show $f(t) = \cos t$ is a characteristic function of some random variable, and find its distribution function.

Solution.....

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Example 8. Show $f(t) = \cos t$ is a characteristic function of some random variable, and find its distribution function.

Solution.....

In general, if $f(t)$ can be written as $\sum a_n e^{ix_n t}$, where $a_n > 0$ and $\sum a_n = 1$, then $f(t)$ is a characteristic function, whose corresponding random variable has distribution sequence $P(\xi = x_n) = a_n, n = 1, 2, \dots$.

3.3 Characteristic functions

3.3.3 Inverse formula and uniqueness theorem

Example 9. If $f(t)$ is a characteristic function of some random variable, show so are $\overline{f(t)}$ and $|f(t)|^2$.

Solution.....

3.3 Characteristic functions

3.3.4 Additivity of distribution functions

3.3.4 Additivity of distribution functions

The additivity, also called regenerativity, means that if ξ and η are independent and follow a common type of distributions, then so do their sum $\xi + \eta$ and the parameter is the sum of parameters of ξ and η .

Example 10. If ξ_j ($j = 1, 2, \dots, k$) follows the binomial distribution $B(n_j, p)$ respectively and are indept. of each other. Find the distribution of

$$\sum_{j=1}^k \xi_j.$$

Solution.

3.3 Characteristic functions

3.3.4 Additivity of distribution functions

Example 10. If ξ_j ($j = 1, 2, \dots, k$) follows the binomial distribution $B(n_j, p)$ respectively and are indept. of each other. Find the distribution of $\sum_{j=1}^k \xi_j$.

Solution. The c.f. $f_j(t)$ of ξ_j is $(pe^{it} + q)^{n_j}$. By Property 4, the c.f. of $\sum_{j=1}^k \xi_j$ is

3.3 Characteristic functions

3.3.4 Additivity of distribution functions

Example 10. If ξ_j ($j = 1, 2, \dots, k$) follows the binomial distribution $B(n_j, p)$ respectively and are indept. of each other. Find the distribution of $\sum_{j=1}^k \xi_j$.

Solution. The c.f. $f_j(t)$ of ξ_j is $(pe^{it} + q)^{n_j}$. By Property 4, the c.f. of $\sum_{j=1}^k \xi_j$ is

$$\prod_{j=1}^k f_j(t) = (pe^{it} + q)^{\sum_{j=1}^k n_j}.$$

3.3 Characteristic functions

3.3.4 Additivity of distribution functions

Example 10. If ξ_j ($j = 1, 2, \dots, k$) follows the binomial distribution $B(n_j, p)$ respectively and are indept. of each other. Find the distribution of $\sum_{j=1}^k \xi_j$.

Solution. The c.f. $f_j(t)$ of ξ_j is $(pe^{it} + q)^{n_j}$. By Property 4, the c.f. of $\sum_{j=1}^k \xi_j$ is

$$\prod_{j=1}^k f_j(t) = (pe^{it} + q)^{\sum_{j=1}^k n_j}.$$

In turn, by the uniqueness theorem,

$$\sum_{j=1}^k \xi_j \sim B(\sum_{j=1}^k n_j, p).$$

3.3 Characteristic functions

3.3.4 Additivity of distribution functions

Example 11. Suppose that ξ_1, \dots, ξ_n are indept., and $\xi_k \sim N(a_k, \sigma_k^2)$, $k = 1, \dots, n$. Find the distribution of $\sum_{k=1}^n \xi_k$.

Solution.

3.3 Characteristic functions

3.3.4 Additivity of distribution functions

Example 11. Suppose that ξ_1, \dots, ξ_n are indept., and $\xi_k \sim N(a_k, \sigma_k^2)$, $k = 1, \dots, n$. Find the distribution of $\sum_{k=1}^n \xi_k$.

Solution. It is known that the c.f. of ξ_k is $e^{ia_k t - \sigma_k^2 t^2 / 2}$, so the c.f. of $\sum_{k=1}^n \xi_k$ is

$$\prod_{k=1}^n e^{ia_k t - \frac{\sigma_k^2 t^2}{2}} = \exp\left\{i \sum_{k=1}^n a_k t - \frac{\sum_{k=1}^n \sigma_k^2 t^2}{2}\right\}.$$

3.3 Characteristic functions

3.3.4 Additivity of distribution functions

Example 11. Suppose that ξ_1, \dots, ξ_n are indept., and $\xi_k \sim N(a_k, \sigma_k^2)$, $k = 1, \dots, n$. Find the distribution of $\sum_{k=1}^n \xi_k$.

Solution. It is known that the c.f. of ξ_k is $e^{ia_k t - \sigma_k^2 t^2 / 2}$, so the c.f. of $\sum_{k=1}^n \xi_k$ is

$$\prod_{k=1}^n e^{ia_k t - \frac{\sigma_k^2 t^2}{2}} = \exp\left\{i \sum_{k=1}^n a_k t - \frac{\sum_{k=1}^n \sigma_k^2 t^2}{2}\right\}.$$

Thus $\sum_{k=1}^n \xi_k \sim N\left(\sum_k a_k, \sum_k \sigma_k^2\right)$.

3.3 Characteristic functions

3.3.5 Multivariate characteristic functions

3.3.5 Multivariate characteristic functions

Definition 3 Suppose the random vector

$\xi = (\xi_1, \dots, \xi_n)'$ has distribution function

$F(x_1, \dots, x_n)$, then its characteristic function is defined by

$$\begin{aligned} f(t_1, \dots, t_n) &= Ee^{i(t_1\xi_1 + \dots + t_n\xi_n)} \\ &= \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} e^{i(t_1x_1 + \dots + t_nx_n)} dF(x_1, \dots, x_n). \end{aligned}$$

3.3 Characteristic functions

3.3.5 Multivariate characteristic functions

3.3.5 Multivariate characteristic functions

Definition 3 Suppose the random vector

$\xi = (\xi_1, \dots, \xi_n)'$ has distribution function

$F(x_1, \dots, x_n)$, then its characteristic function is defined by

$$\begin{aligned} f(t_1, \dots, t_n) &= Ee^{i(t_1\xi_1 + \dots + t_n\xi_n)} \\ &= \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} e^{i(t_1x_1 + \dots + t_nx_n)} dF(x_1, \dots, x_n). \end{aligned}$$

$$f(\mathbf{t}) = Ee^{it'\xi} = \int_{\mathbf{R}^n} e^{it'\mathbf{x}} dF(\mathbf{x}),$$

where $\mathbf{t} = (t_1, \dots, t_n)'$, $\mathbf{x} = (x_1, \dots, x_n)'$.

3.3 Characteristic functions

3.3.5 Multivariate characteristic functions

① The c.f. of $\eta = a_1\xi_1 + \cdots + a_n\xi_n$ is

$$\begin{aligned}f_{\eta}(t) &= Ee^{it\eta} = Ee^{it\sum a_k\xi_k} \\&= Ee^{i\sum (a_k t)\xi_k} = f(a_1t, \cdots, a_nt).\end{aligned}$$

② If the c.f. of $(\xi_1, \cdots, \xi_n)'$ is $f(t_1, \cdots, t_n)$, then k -dimensional sub-vector $(\xi_{l_1}, \cdots, \xi_{l_k})'$ has c.f.

$$f(0, \cdots, 0, t_{l_1}, 0, \cdots, 0, t_{l_k}, 0, \cdots, 0).$$

3.3 Characteristic functions

3.3.5 Multivariate characteristic functions

- ③ Assume that ξ_j has c.f. $f_j(t)$, $j = 1, \dots, n$, then ξ_1, \dots, ξ_n are indept. iff the c.f. of $(\xi_1, \dots, \xi_n)'$ is such that

$$f(t_1, \dots, t_n) = f_1(t_1) \cdots f_n(t_n).$$

- ④ $(\xi_1, \dots, \xi_k)'$ and $(\xi_{k+1}, \dots, \xi_n)'$ are indept. iff the product of their c.f.s is just equal to the c.f. of $(\xi_1, \dots, \xi_n)'$.