

## 2.2 Distribution functions and continuous random variables

## 2.2.1 Distribution functions

## 2.2 Distribution functions and continuous random variables

### 2.2.1 Distribution functions

# 1. Definitions

There is no distribution sequence for other types of random variables. If all possible values of a random variable consists of an interval, then we are not able to enumerate all these values and their probabilities.

We usually we want calculate the probability of the type  $P(a < \xi \leq b)$ .

We usually we want calculate the probability of the type  $P(a < \xi \leq b)$ . Notice

$$P(a < \xi \leq b) = P(\xi \leq b) - P(\xi \leq a).$$

We only need to the probability of the type

$$P(\xi \leq x).$$

**Definition** Let  $\xi$  be a random variable on a probability space  $(\Omega, \mathcal{F}, P)$ , We define its distribution function (sometimes, cumulative distribution function (CDF) ) as

$$F(x) = P(\xi \leq x), \quad -\infty < x < \infty.$$

**Definition** Let  $\xi$  be a random variable on a probability space  $(\Omega, \mathcal{F}, P)$ , We define its distribution function (sometimes, cumulative distribution function (CDF) ) as

$$F(x) = P(\xi \leq x), \quad -\infty < x < \infty.$$

Having distribution function, the probability  $P(\xi(\omega) \in B)$  can be expressed in term of it for any Borel set  $B$ . For example:

## 2.2 Distribution functions and continuous random variables

## 2.2.1 Distribution functions

$$P(a < \xi \leq b) = F(b) - F(a)$$

## 2.2 Distribution functions and continuous random variables

## 2.2.1 Distribution functions

$$P(a < \xi \leq b) = F(b) - F(a)$$

$$P(\xi < a) = \lim_{b \rightarrow a-0} P(\xi \leq b) = F(a-0);$$



## 2.2 Distribution functions and continuous random variables

## 2.2.1 Distribution functions

$$P(a < \xi \leq b) = F(b) - F(a)$$

$$P(\xi < a) = \lim_{b \rightarrow a-0} P(\xi \leq b) = F(a-0);$$

$$\begin{aligned} P(\xi = a) &= P(\xi \leq a) - P(\xi < a) \\ &= F(a) - F(a-0); \end{aligned}$$

## 2.2 Distribution functions and continuous random variables

## 2.2.1 Distribution functions

$$P(a < \xi \leq b) = F(b) - F(a)$$

$$P(\xi < a) = \lim_{b \rightarrow a-0} P(\xi \leq b) = F(a-0);$$

$$\begin{aligned} P(\xi = a) &= P(\xi \leq a) - P(\xi < a) \\ &= F(a) - F(a-0); \end{aligned}$$

$$P(\xi > a) = 1 - P(\xi \leq a) = 1 - F(a);$$

## 2.2 Distribution functions and continuous random variables

## 2.2.1 Distribution functions

$$P(a < \xi \leq b) = F(b) - F(a)$$

$$P(\xi < a) = \lim_{b \rightarrow a-0} P(\xi \leq b) = F(a-0);$$

$$\begin{aligned} P(\xi = a) &= P(\xi \leq a) - P(\xi < a) \\ &= F(a) - F(a-0); \end{aligned}$$

$$P(\xi > a) = 1 - P(\xi \leq a) = 1 - F(a);$$

$$\begin{aligned} P(a < \xi < b) &= P(\xi < b) - P(\xi \leq a) \\ &= F(b-0) - F(a). \end{aligned}$$

**Example 1.** Suppose that a random variable  $\xi$  is distributed as Bernoulli distribution:

$$\begin{pmatrix} 0 & 1 \\ q & p \end{pmatrix}, \quad p, q > 0, \quad p + q = 1.$$

Determine its distribution function  $F(x)$ , and calculate  $P(-1 < \xi < 0.5)$ .

**Solution.** When  $x < 0$ ,  $P(\xi \leq x) = 0$  (null event);

**Solution.** When  $x < 0$ ,  $P(\xi \leq x) = 0$  (null event);  
when  $0 \leq x < 1$ ,  $P(\xi \leq x) = P(\xi = 0) = q$ ;

**Solution.** When  $x < 0$ ,  $P(\xi \leq x) = 0$  (null event);  
when  $0 \leq x < 1$ ,  $P(\xi \leq x) = P(\xi = 0) = q$ ;  
when  $x \geq 1$ ,

$$P(\xi \leq x) = P(\xi = 0) + P(\xi = 1) = q + p = 1.$$

Hence we obtain the following distribution function:

$$F(x) = \begin{cases} 0, & x < 0, \\ q, & 0 \leq x < 1, \\ 1, & x \geq 1. \end{cases}$$



Hence we obtain the following distribution function:

$$F(x) = \begin{cases} 0, & x < 0, \\ q, & 0 \leq x < 1, \\ 1, & x \geq 1. \end{cases}$$

Next, let us compute  $P(-1 < \xi < 0.5)$ .

$$P(-1 < \xi < 0.5) = F(0.5 - 0) - F(-1) = q.$$

## 2.2 Distribution functions and continuous random variables

## 2.2.1 Distribution functions

Suppose that  $\xi$  has the distribution sequence


$$\begin{pmatrix} x_1 & x_2 & \cdots & x_n & \cdots \\ p(x_1) & p(x_2) & \cdots & p(x_n) & \cdots \end{pmatrix},$$

where  $x_1 < x_2 < \cdots < x_k < \cdots$ , then

$$F(x) = \sum_{k: x_k \leq x} p(x_k) = \begin{cases} 0, & x < x_1, \\ p(x_1), & x_1 \leq x < x_2, \\ \cdots & \cdots \\ \sum_{i \leq k} p(x_i), & x_k \leq x < x_{k+1}, \\ \cdots & \cdots \end{cases}$$


## 2.2 Distribution functions and continuous random variables

## 2.2.1 Distribution functions



**Example 2.** Choose randomly a point  $P$  in a triangle  $\triangle ABC$ , let  $\xi$  be the distance from  $P$  to edge  $BC$ . Calculate the distribution function of  $\xi$ .

# Solution.



## 2.2 Distribution functions and continuous random variables

## 2.2.1 Distribution functions

Let  $h$  be the height of edge  $BC$ . It is obvious that

$$P(\xi \leq x) = 0 \text{ when } x < 0$$

## 2.2 Distribution functions and continuous random variables

## 2.2.1 Distribution functions

Let  $h$  be the height of edge  $BC$ . It is obvious that  $P(\xi \leq x) = 0$  when  $x < 0$  and  $P(\xi \leq x) = 1$  when  $x \geq h$ .

## 2.2 Distribution functions and continuous random variables

## 2.2.1 Distribution functions

Let  $h$  be the height of edge  $BC$ . It is obvious that  $P(\xi \leq x) = 0$  when  $x < 0$  and  $P(\xi \leq x) = 1$  when  $x \geq h$ . For  $0 \leq x < h$ , make a line segment  $DE$  paralleling edge  $BC$  with distance  $x$  to  $BC$  inside  $\triangle ABC$ . Then  $\xi \leq x$  if and only if  $P$  falls within  $DBCE$ .



Let  $h$  be the height of edge  $BC$ . It is obvious that  $P(\xi \leq x) = 0$  when  $x < 0$  and  $P(\xi \leq x) = 1$  when  $x \geq h$ . For  $0 \leq x < h$ , make a line segment  $DE$  paralleling edge  $BC$  with distance  $x$  to  $BC$  inside  $\triangle ABC$ . Then  $\xi \leq x$  if and only if  $P$  falls within  $DBCE$ . So

$$P(\xi \leq x) = \frac{\text{area of } DBCE}{\text{area of } \triangle ABC} =$$

Let  $h$  be the height of edge  $BC$ . It is obvious that  $P(\xi \leq x) = 0$  when  $x < 0$  and  $P(\xi \leq x) = 1$  when  $x \geq h$ . For  $0 \leq x < h$ , make a line segment  $DE$  paralleling edge  $BC$  with distance  $x$  to  $BC$  inside  $\triangle ABC$ . Then  $\xi \leq x$  if and only if  $P$  falls within  $DBCE$ . So

$$P(\xi \leq x) = \frac{\text{area of } DBCE}{\text{area of } \triangle ABC} = 1 - \left(1 - \frac{x}{h}\right)^2.$$

The distribution function is

$$F(x) = \begin{cases} 0, & x < 0, \\ 1 - \left(1 - \frac{x}{h}\right)^2, & 0 \leq x < h, \\ 1, & x \geq h. \end{cases}$$

## 2. Properties Three fundamental properties:

- ①  $F(x)$  is monotonic non-decreasing in  $x$ , that is, if  $a < b$  then  $F(a) \leq F(b)$ ;
- ②  $\lim_{x \rightarrow -\infty} F(x) = 0, \lim_{x \rightarrow +\infty} F(x) = 1$ ;
- ③  $F(x)$  is right continuous, that is  $F(x+0) = F(x)$ .

## 2.2 Distribution functions and continuous random variables

## 2.2.1 Distribution functions

**Proof.** (1) If  $a < b$ , then  $\{\xi \leq a\} \subset \{\xi \leq b\}$ .

## 2.2 Distribution functions and continuous random variables

## 2.2.1 Distribution functions

**Proof.** (1) If  $a < b$ , then  $\{\xi \leq a\} \subset \{\xi \leq b\}$ . So  
$$F(a) = P(\xi \leq a) \leq P(\xi \leq b) = F(b).$$

**Proof.** (1) If  $a < b$ , then  $\{\xi \leq a\} \subset \{\xi \leq b\}$ . So

$$F(a) = P(\xi \leq a) \leq P(\xi \leq b) = F(b).$$

(2) By the monotonicity of  $F(x)$ , it follows that

$F(-\infty)$  exists and

$$F(-\infty) = \lim_{n \rightarrow \infty} F(-n).$$

**Proof.** (1) If  $a < b$ , then  $\{\xi \leq a\} \subset \{\xi \leq b\}$ . So

$$F(a) = P(\xi \leq a) \leq P(\xi \leq b) = F(b).$$

(2) By the monotonicity of  $F(x)$ , it follows that

$F(-\infty)$  exists and

$$F(-\infty) = \lim_{n \rightarrow \infty} F(-n).$$

Notice that

$$\{\xi \leq -(n+1)\} \subset \{\xi \leq -n\},$$



**Proof.** (1) If  $a < b$ , then  $\{\xi \leq a\} \subset \{\xi \leq b\}$ . So

$$F(a) = P(\xi \leq a) \leq P(\xi \leq b) = F(b).$$

(2) By the monotonicity of  $F(x)$ , it follows that

$F(-\infty)$  exists and

$$F(-\infty) = \lim_{n \rightarrow \infty} F(-n).$$

Notice that

$$\{\xi \leq -(n+1)\} \subset \{\xi \leq -n\}, \quad \bigcap_{n=1}^{\infty} \{\xi \leq -n\} = \emptyset.$$

We have

$$\lim_{n \rightarrow \infty} F(-n) = \lim_{n \rightarrow \infty} P(\xi \leq -n)$$

We have

$$\begin{aligned}\lim_{n \rightarrow \infty} F(-n) &= \lim_{n \rightarrow \infty} P(\xi \leq -n) \\ &= P\left(\bigcap_{n=1}^{\infty} \{\xi \leq -n\}\right) = P(\emptyset) = 0.\end{aligned}$$

We have

$$\begin{aligned}\lim_{n \rightarrow \infty} F(-n) &= \lim_{n \rightarrow \infty} P(\xi \leq -n) \\ &= P\left(\bigcap_{n=1}^{\infty} \{\xi \leq -n\}\right) = P(\emptyset) = 0.\end{aligned}$$

Similarly,

$$F(+\infty) = \lim_{n \rightarrow \infty} F(n) = \lim_{n \rightarrow \infty} P(\xi \leq n)$$

We have

$$\begin{aligned}\lim_{n \rightarrow \infty} F(-n) &= \lim_{n \rightarrow \infty} P(\xi \leq -n) \\ &= P\left(\bigcap_{n=1}^{\infty} \{\xi \leq -n\}\right) = P(\emptyset) = 0.\end{aligned}$$

Similarly,

$$\begin{aligned}F(+\infty) &= \lim_{n \rightarrow \infty} F(n) = \lim_{n \rightarrow \infty} P(\xi \leq n) \\ &= P\left(\lim_{n \rightarrow \infty} \{\xi \leq n\}\right) = P\left(\bigcup_{n=1}^{\infty} \{\xi \leq n\}\right) \\ &= P(\Omega) = 1.\end{aligned}$$

## 2.2 Distribution functions and continuous random variables

## 2.2.1 Distribution functions

(3) By the monotonicity of  $F(x)$  again, it follows that  $F(x+0) = \lim_{n \rightarrow \infty} F(x + 1/n)$ .

(3) By the monotonicity of  $F(x)$  again, it follows that  $F(x+0) = \lim_{n \rightarrow \infty} F(x + 1/n)$ . Now

$$\begin{aligned} & \lim_{n \rightarrow \infty} F(x + 1/n) \\ = & \lim_{n \rightarrow \infty} P(\xi \leq x + 1/n) \end{aligned}$$

## 2.2 Distribution functions and continuous random variables

## 2.2.1 Distribution functions

(3) By the monotonicity of  $F(x)$  again, it follows that  $F(x+0) = \lim_{n \rightarrow \infty} F(x + 1/n)$ . Now

$$\begin{aligned} & \lim_{n \rightarrow \infty} F(x + 1/n) \\ &= \lim_{n \rightarrow \infty} P(\xi \leq x + 1/n) \\ &= P\left(\lim_{n \rightarrow \infty} \{\xi \leq x + 1/n\}\right) \quad (\because \{\xi \leq x + 1/n\} \searrow) \end{aligned}$$



(3) By the monotonicity of  $F(x)$  again, it follows that  $F(x+0) = \lim_{n \rightarrow \infty} F(x + 1/n)$ . Now

$$\begin{aligned} & \lim_{n \rightarrow \infty} F(x + 1/n) \\ &= \lim_{n \rightarrow \infty} P(\xi \leq x + 1/n) \\ &= P\left(\lim_{n \rightarrow \infty} \{\xi \leq x + 1/n\}\right) \quad (\because \{\xi \leq x + 1/n\} \searrow) \\ &= P\left(\bigcap_{n=1}^{\infty} \{\xi \leq x + 1/n\}\right) \end{aligned}$$

## 2.2 Distribution functions and continuous random variables

## 2.2.1 Distribution functions

(3) By the monotonicity of  $F(x)$  again, it follows that  $F(x+0) = \lim_{n \rightarrow \infty} F(x + 1/n)$ . Now

$$\begin{aligned} & \lim_{n \rightarrow \infty} F(x + 1/n) \\ &= \lim_{n \rightarrow \infty} P(\xi \leq x + 1/n) \\ &= P(\lim_{n \rightarrow \infty} \{\xi \leq x + 1/n\}) \quad (\because \{\xi \leq x + 1/n\} \searrow) \\ &= P(\bigcap_{n=1}^{\infty} \{\xi \leq x + 1/n\}) \\ &= P(\xi \leq x) = F(x). \end{aligned}$$

Similarly,

$$\begin{aligned} F(x-0) &= \lim_{n \rightarrow \infty} F(x-1/n) \\ &= \lim_{n \rightarrow \infty} P(\xi \leq x-1/n) \end{aligned}$$

Similarly,

$$\begin{aligned} F(x-0) &= \lim_{n \rightarrow \infty} F(x-1/n) \\ &= \lim_{n \rightarrow \infty} P(\xi \leq x-1/n) \\ &= P\left(\lim_{n \rightarrow \infty} \{\xi \leq x-1/n\}\right) \quad (\because \{\xi \leq x-1/n\} \nearrow) \end{aligned}$$

Similarly,

$$\begin{aligned} F(x-0) &= \lim_{n \rightarrow \infty} F(x-1/n) \\ &= \lim_{n \rightarrow \infty} P(\xi \leq x-1/n) \\ &= P\left(\lim_{n \rightarrow \infty} \{\xi \leq x-1/n\}\right) \quad (\because \{\xi \leq x-1/n\} \nearrow) \\ &= P\left(\bigcup_{n=1}^{\infty} \{\xi \leq x-1/n\}\right) \end{aligned}$$

Similarly,

$$\begin{aligned} F(x-0) &= \lim_{n \rightarrow \infty} F(x-1/n) \\ &= \lim_{n \rightarrow \infty} P(\xi \leq x-1/n) \\ &= P\left(\lim_{n \rightarrow \infty} \{\xi \leq x-1/n\}\right) \quad (\because \{\xi \leq x-1/n\} \nearrow) \\ &= P\left(\bigcup_{n=1}^{\infty} \{\xi \leq x-1/n\}\right) \\ &= P(\xi < x). \end{aligned}$$

**Example 3.** Suppose a random variable has the distribution function as follows:

$$F(x) = \begin{cases} 0, & x \leq -1, \\ a + b \arcsin x, & -1 < x \leq 1, \\ 1, & x > 1. \end{cases}$$

Find constants  $a, b$ .

**Solution.** From the fact that

$$F(-1 + 0) = F(-1) \quad \text{and} \quad F(1 + 0) = F(1),$$



**Solution.** From the fact that

$$F(-1+0) = F(-1) \quad \text{and} \quad F(1+0) = F(1),$$

it follows that

$$a - b\pi/2 = 0 \quad \text{and} \quad a + b\pi/2 = 1.$$

Hence  $a = 1/2$ ,  $b = 1/\pi$ .

## Theorem

*Let  $F(x)$  be a real function satisfying Properties (1), (2) and (3). Then there is an unique probability measure  $P_F: \mathcal{B} \rightarrow [0, 1]$ , such that*

$$P_F\left((-\infty, x]\right) = F(x) \quad \forall x.$$

## Theorem

Let  $F(x)$  be a real function satisfying Properties (1), (2) and (3). Then there is an unique probability measure  $P_F: \mathcal{B} \rightarrow [0, 1]$ , such that

$$P_F\left((-\infty, x]\right) = F(x) \quad \forall x.$$

On the probability space  $(\mathbf{R}, \mathcal{B}, P_F)$ , define  $X(r) = r, r \in \mathbf{R}$ . Then  $X$  is a r.v. variable with

$$P_F(X \leq x) = P_F\left((-\infty, x]\right) = F(x).$$

So  $F$  is the distribution function of  $X$ .

## 2.2.2 Continuous random variables and density functions

**Definition.** Suppose that a random variable  $\xi$  takes all values of an interval (finite or infinite), and that there exists a non-negative integrable function  $p(x)$  such that the distribution function  $F(x)$  can be written as

$$F(x) = \int_{-\infty}^x p(y)dy, \quad -\infty < x < \infty.$$

Then  $\xi$  is called a continuous random variable, and  $p(x)$  is called the probability density function (PDF) of  $\xi$ , or more simply its density function.  $F(x)$ , having the above property, is said to be absolutely continuous.

- ①  $F(x)$  is continuous. And  $F'(x) = p(x)$  if  $x$  is a continuity point of  $p(x)$

①  $F(x)$  is continuous. And  $F'(x) = p(x)$  if  $x$  is a continuity point of  $p(x)$

②

$$\begin{aligned} P(a < \xi \leq b) &= F(b) - F(a) \\ &= \int_{-\infty}^b p(y) dy - \int_{-\infty}^a p(y) dy \\ &= \int_a^b p(y) dy. \end{aligned}$$

①  $F(x)$  is continuous. And  $F'(x) = p(x)$  if  $x$  is a continuity point of  $p(x)$

②

$$\begin{aligned} P(a < \xi \leq b) &= F(b) - F(a) \\ &= \int_{-\infty}^b p(y) dy - \int_{-\infty}^a p(y) dy \\ &= \int_a^b p(y) dy. \end{aligned}$$



①  $F(x)$  is continuous. And  $F'(x) = p(x)$  if  $x$  is a continuity point of  $p(x)$

②

$$\begin{aligned} P(a < \xi \leq b) &= F(b) - F(a) \\ &= \int_{-\infty}^b p(y) dy - \int_{-\infty}^a p(y) dy \\ &= \int_a^b p(y) dy. \end{aligned}$$

$$P(\xi \in B) = \int_B p(x) dx, \quad B \in \mathcal{B}$$

3

$$\begin{aligned}P(\xi = c) &= F(c) - F(c - 0) \\&= \lim_{h \rightarrow 0^+} \int_{c-h}^c p(y) dy = 0.\end{aligned}$$

The density function possesses the following properties.

- $p(x) \geq 0$ ,
- $\int_{-\infty}^{\infty} p(x)dx = 1$ .

On the contrary, if a function defined in  $(-\infty, \infty)$  satisfies these two properties then it can be considered to be a density function of some random variable.

$$F(x) = \begin{cases} 0, & x < 0 \\ (1+x)/2, & 0 \leq x < 1, \\ 1, & x \geq 1 \end{cases}$$

is a distribution function. But it is neither discrete nor continuous (discontinuous).

$$F(x) = \begin{cases} 0, & x < 0 \\ (1+x)/2, & 0 \leq x < 1, \\ 1, & x \geq 1 \end{cases}$$

is a distribution function. But it is neither discrete nor continuous (discontinuous).

In fact, it is a mixture of  $F_1(x)$  and  $F_2(x)$ :

$$F(x) = \frac{F_1(x) + F_2(x)}{2},$$

where  $F_1$  is a degenerate distribution at  $x = 0$  and  $F_2(x)$  is uniform on  $[0, 1]$ .

## 2.2.3 Typical continuous random variables

### 1. The uniform distribution $\xi \sim U(a, b)$

$$p(x) = \begin{cases} 1/(b-a), & a \leq x \leq b, \\ 0, & \text{otherwise.} \end{cases}$$

## 2.2.3 Typical continuous random variables

### 1. The uniform distribution $\xi \sim U(a, b)$

$$p(x) = \begin{cases} 1/(b-a), & a \leq x \leq b, \\ 0, & \text{otherwise.} \end{cases}$$

Distribution function:

$$F(x) = \begin{cases} 0, & x < a, \\ (x-a)/(b-a), & a \leq x < b, \\ 1, & x \geq b. \end{cases}$$

**2. The normal distribution**  $\xi \sim N(\mu, \sigma^2)$ ,  
where  $-\infty < \mu < \infty, \sigma > 0$ .


$$p(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, \quad -\infty < x < \infty,$$

$$F(x) = \frac{1}{\sqrt{2\pi}\sigma} \int_{-\infty}^x e^{-\frac{(t-\mu)^2}{2\sigma^2}} dt, \quad -\infty < x < \infty.$$



## 2.2 Distribution functions and continuous random variables

## 2.2.3 Typical continuous random variables



Verifying  $\int p(x)dx = 1$ : Let

$$I \quad =: \quad \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx$$

Verifying  $\int p(x)dx = 1$ : Let

$$\begin{aligned} I &=: \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx \\ &\stackrel{t=(x-\mu)/\sigma}{=} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{t^2}{2}} dt. \end{aligned}$$

Hence

$$I^2 = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{1}{2\pi} e^{-\frac{x^2+y^2}{2}} dx dy$$

Verifying  $\int p(x)dx = 1$ : Let

$$I =: \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx$$

$$\stackrel{t=(x-\mu)/\sigma}{=} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{t^2}{2}} dt.$$

Hence

$$I^2 = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{1}{2\pi} e^{-\frac{x^2+y^2}{2}} dx dy$$

$$= \int_0^{\infty} \int_0^{2\pi} \frac{1}{2\pi} e^{-\frac{r^2}{2}} r dr d\theta$$

Verifying  $\int p(x)dx = 1$ : Let

$$I =: \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx$$

$$\stackrel{t=(x-\mu)/\sigma}{=} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{t^2}{2}} dt.$$

Hence

$$I^2 = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{1}{2\pi} e^{-\frac{x^2+y^2}{2}} dx dy$$

$$= \int_0^{\infty} \int_0^{2\pi} \frac{1}{2\pi} e^{-\frac{r^2}{2}} r dr d\theta =$$

Verifying  $\int p(x)dx = 1$ : Let

$$I =: \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx$$

$$\stackrel{t=(x-\mu)/\sigma}{=} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{t^2}{2}} dt.$$

Hence

$$I^2 = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{1}{2\pi} e^{-\frac{x^2+y^2}{2}} dx dy$$

$$= \int_0^{\infty} \int_0^{2\pi} \frac{1}{2\pi} e^{-\frac{r^2}{2}} r dr d\theta = -e^{-\frac{r^2}{2}} \Big|_0^{\infty} = 1.$$


## Standard normal distribution: $N(0, 1)$

$$\varphi(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}, \quad \Phi(x) = \int_{-\infty}^x \varphi(t) dt,$$

$$-\infty < x < \infty.$$

## 2.2 Distribution functions and continuous random variables

## 2.2.3 Typical continuous random variables





## Table of normal distribution.

(1)  $\xi \sim N(0, 1)$ .

•  $x > 0$ : .....

•  $x < 0$ :  $\Phi(x) = 1 - \Phi(-x)$ .

## Table of normal distribution.

(1)  $\xi \sim N(0, 1)$ .

•  $x > 0$ : .....

•  $x < 0$ :  $\Phi(x) = 1 - \Phi(-x)$ .

(2)  $\xi \sim N(\mu, \sigma^2)$

## Table of normal distribution.

(1)  $\xi \sim N(0, 1)$ .

- $x > 0$ : .....

- $x < 0$ :  $\Phi(x) = 1 - \Phi(-x)$ .

(2)  $\xi \sim N(\mu, \sigma^2) \implies \eta = \frac{\xi - \mu}{\sigma} \sim N(0, 1)$ .

## Table of normal distribution.

(1)  $\xi \sim N(0, 1)$ .

•  $x > 0$ : .....

•  $x < 0$ :  $\Phi(x) = 1 - \Phi(-x)$ .

(2)  $\xi \sim N(\mu, \sigma^2) \implies \eta = \frac{\xi - \mu}{\sigma} \sim N(0, 1)$ .

(Proof?)

## Table of normal distribution.

(1)  $\xi \sim N(0, 1)$ .

•  $x > 0$ : .....

•  $x < 0$ :  $\Phi(x) = 1 - \Phi(-x)$ .

(2)  $\xi \sim N(\mu, \sigma^2) \implies \eta = \frac{\xi - \mu}{\sigma} \sim N(0, 1)$ .

(Proof?)

So

$$P(\xi \leq x) = \Phi\left(\frac{x - \mu}{\sigma}\right).$$

**Example 5.** Let  $\xi \sim N(0, 1)$ .

(1) Find  $P(-1 < \xi < 3)$ ;

(2) Suppose  $P(\xi < \lambda) = 0.9755$ , find  $\lambda$ .

**Solution.** (1)

$$\begin{aligned} & P(-1 < \xi < 3) \\ &= \Phi(3) - \Phi(-1) \end{aligned}$$

**Solution.** (1)

$$\begin{aligned} & P(-1 < \xi < 3) \\ &= \Phi(3) - \Phi(-1) = \Phi(3) + \Phi(1) - 1 \end{aligned}$$



**Solution.** (1)

$$\begin{aligned} & P(-1 < \xi < 3) \\ &= \Phi(3) - \Phi(-1) = \Phi(3) + \Phi(1) - 1 \\ &= 0.9987 + 0.8413 - 1 = 0.8400. \end{aligned}$$

**Solution.** (1)

$$\begin{aligned} & P(-1 < \xi < 3) \\ &= \Phi(3) - \Phi(-1) = \Phi(3) + \Phi(1) - 1 \\ &= 0.9987 + 0.8413 - 1 = 0.8400. \end{aligned}$$

(2) Note that  $\Phi(\lambda) = 0.9755$ , which lies between  $\Phi(1.96) = 0.9750$  and  $\Phi(1.98) = 0.9762$ .

**Solution.** (1)

$$\begin{aligned} & P(-1 < \xi < 3) \\ &= \Phi(3) - \Phi(-1) = \Phi(3) + \Phi(1) - 1 \\ &= 0.9987 + 0.8413 - 1 = 0.8400. \end{aligned}$$

(2) Note that  $\Phi(\lambda) = 0.9755$ , which lies between  $\Phi(1.96) = 0.9750$  and  $\Phi(1.98) = 0.9762$ . By using a linear interpolation,

$$\begin{aligned} \lambda &\approx 1.96 + \frac{\Phi(\lambda) - \Phi(1.96)}{\Phi(1.98) - \Phi(1.96)} \cdot (1.98 - 1.96) \\ &\approx 1.968. \end{aligned}$$

**Example 6.** Suppose that  $\xi \sim N(2, 9)$ , calculate  $P(5 < \xi < 20)$ .

**Solution.**

**Example 6.** Suppose that  $\xi \sim N(2, 9)$ , calculate  $P(5 < \xi < 20)$ .

**Solution.** Let  $\eta = (\xi - 2)/3$ , then  $\eta \sim N(0, 1)$ .

Hence

**Example 6.** Suppose that  $\xi \sim N(2, 9)$ , calculate  $P(5 < \xi < 20)$ .

**Solution.** Let  $\eta = (\xi - 2)/3$ , then  $\eta \sim N(0, 1)$ .

Hence

$$P(5 < \xi < 20)$$

**Example 6.** Suppose that  $\xi \sim N(2, 9)$ , calculate  $P(5 < \xi < 20)$ .

**Solution.** Let  $\eta = (\xi - 2)/3$ , then  $\eta \sim N(0, 1)$ .

Hence

$$\begin{aligned} & P(5 < \xi < 20) \\ = & P\left(\frac{5-2}{3} < \frac{\xi-2}{3} < \frac{20-2}{3}\right) \end{aligned}$$

**Example 6.** Suppose that  $\xi \sim N(2, 9)$ , calculate  $P(5 < \xi < 20)$ .

**Solution.** Let  $\eta = (\xi - 2)/3$ , then  $\eta \sim N(0, 1)$ .

Hence

$$\begin{aligned} & P(5 < \xi < 20) \\ &= P\left(\frac{5-2}{3} < \frac{\xi-2}{3} < \frac{20-2}{3}\right) \\ &= P(1 < \eta < 6) = \Phi(6) - \Phi(1) \\ &\approx 1 - 0.8413 = 0.1587. \end{aligned}$$



**Example 7.** Suppose that  $\xi \sim N(\mu, \sigma^2)$ , find

$P(|\xi - \mu| < \sigma)$ ,  $P(|\xi - \mu| < 2\sigma)$  and

$P(|\xi - \mu| < 3\sigma)$ .

**Solution.**

**Example 7.** Suppose that  $\xi \sim N(\mu, \sigma^2)$ , find  $P(|\xi - \mu| < \sigma)$ ,  $P(|\xi - \mu| < 2\sigma)$  and  $P(|\xi - \mu| < 3\sigma)$ .

**Solution.** Let  $\eta = (\xi - \mu)/\sigma$ , then  $\eta \sim N(0, 1)$ .

Hence

**Example 7.** Suppose that  $\xi \sim N(\mu, \sigma^2)$ , find

$P(|\xi - \mu| < \sigma)$ ,  $P(|\xi - \mu| < 2\sigma)$  and

$P(|\xi - \mu| < 3\sigma)$ .

**Solution.** Let  $\eta = (\xi - \mu)/\sigma$ , then  $\eta \sim N(0, 1)$ .

Hence

$$P(|\xi - \mu| < \sigma) = P(|\eta| < 1) = 2\Phi(1) - 1 \approx 0.6827.$$

**Example 7.** Suppose that  $\xi \sim N(\mu, \sigma^2)$ , find  $P(|\xi - \mu| < \sigma)$ ,  $P(|\xi - \mu| < 2\sigma)$  and  $P(|\xi - \mu| < 3\sigma)$ .

**Solution.** Let  $\eta = (\xi - \mu)/\sigma$ , then  $\eta \sim N(0, 1)$ .

Hence

$$P(|\xi - \mu| < \sigma) = P(|\eta| < 1) = 2\Phi(1) - 1 \approx 0.6827.$$

Similarly,

$$P(|\xi - \mu| < 2\sigma) = P(|\eta| < 2) \approx 0.9545,$$

$$P(|\xi - \mu| < 3\sigma) = P(|\eta| < 3) \approx 0.9973.$$

## 2.2 Distribution functions and continuous random variables

## 2.2.3 Typical continuous random variables

Some important values of  $\Phi(x)$ 

$x$	1	2	3	4
$\Phi(x)$	0.841345	0.977250	0.998650	0.999968
$2\Phi(x) - 1$	0.6826895	0.9544997	0.9973002	0.9999367

$\Phi(x)$	0.9000	0.9500	0.9750	0.9900	0.9950
$x$	1.2816	1.6449	1.9600	2.3263	2.5758

**例题：**某人被控告为一个新生儿的父亲. 此案鉴定人作证时指出:母亲的怀孕的天数(即从受孕到婴儿出生的时间)近似地服从正态分布, 参数为 $\mu = 270$ ,  $\sigma^2 = 100$ . 有证据表明: 被告在孩子出生前290天出国, 而于出生前240天才回来. ???

**解:** 用 $X$ 表示怀孕期的天数, 如果被告是孩子的父亲, 那么 $X \geq 290$ 或者 $X \leq 240$ ,

**解：**用 $X$ 表示怀孕期的天数，如果被告是孩子的父亲，那么 $X \geq 290$ 或者 $X \leq 240$ ，这一事件发生的概率为

$$\begin{aligned} & P(X \geq 290 \text{ or } X \leq 240) \\ &= P(X \geq 290) + P(X \leq 240) \end{aligned}$$



**解:** 用 $X$ 表示怀孕期的天数, 如果被告是孩子的父亲, 那么 $X \geq 290$ 或者 $X \leq 240$ , 这一事件发生的概率为

$$\begin{aligned} & P(X \geq 290 \text{ or } X \leq 240) \\ &= P(X \geq 290) + P(X \leq 240) \\ &= P\left(\frac{X - 270}{10} \geq 2\right) + P\left(\frac{X - 270}{10} \leq -3\right) \end{aligned}$$

**解:** 用 $X$ 表示怀孕期的天数, 如果被告是孩子的父亲, 那么 $X \geq 290$ 或者 $X \leq 240$ , 这一事件发生的概率为

$$\begin{aligned} & P(X \geq 290 \text{ or } X \leq 240) \\ &= P(X \geq 290) + P(X \leq 240) \\ &= P\left(\frac{X - 270}{10} \geq 2\right) + P\left(\frac{X - 270}{10} \leq -3\right) \\ &= 1 - \Phi(2) + 1 - \Phi(3) \\ &= 1 - 0.9772 + 1 - 0.9987 = 0.0241. \end{aligned}$$

**Example 8.** There are two ways by bus from a city's southern district to a train station located in the city's northern area. The first route is shorter, but the ride encounters heavy traffic, so the time  $\tau$  required is  $N(50, 100)$ ; the second ride is a bit longer, but unexpected traffic jams seldom occur and the time  $\tau$  required is  $N(60, 16)$ .

- ❶ If one has 70 minutes, then what way should be chosen?
- ❷ What is it about if one has 65 minutes ?

# Solution.

**Solution.** (1) For the first route, the probability one can arrive at the train station within 70 minutes is

$$P(\tau \leq 70) = \Phi\left(\frac{70 - 50}{\sqrt{100}}\right) = \Phi(2) = 0.9772;$$

**Solution.** (1) For the first route, the probability one can arrive at the train station within 70 minutes is

$$P(\tau \leq 70) = \Phi\left(\frac{70 - 50}{\sqrt{100}}\right) = \Phi(2) = 0.9772;$$

For the second route, the probability one can arrive at the train station within 70 minutes is

$$P(\tau \leq 70) = \Phi\left(\frac{70 - 60}{\sqrt{16}}\right) = \Phi(2.5) = 0.9938.$$

So, it is better to take the second route.

(2) For the first route, the probability one can arrive at the train station within 65 minutes is

$$P(\tau \leq 65) = \Phi\left(\frac{65 - 50}{\sqrt{100}}\right) = \Phi(1.5) = 0.9332;$$

(2) For the first route, the probability one can arrive at the train station within 65 minutes is

$$P(\tau \leq 65) = \Phi\left(\frac{65 - 50}{\sqrt{100}}\right) = \Phi(1.5) = 0.9332;$$

For the second route, the probability one can arrive at the train station within 65 minutes is

$$P(\tau \leq 65) = \Phi\left(\frac{65 - 60}{\sqrt{16}}\right) = \Phi(1.25) = 0.8944.$$

So, it is better to take the first route.



### 3. The Exponential distribution

Density function:

$$p(x) = \begin{cases} \lambda e^{-\lambda x}, & x \geq 0, \\ 0, & x < 0. \end{cases} \quad (\lambda > 0).$$

Distribution function:

### 3. The Exponential distribution

Density function:

$$p(x) = \begin{cases} \lambda e^{-\lambda x}, & x \geq 0, \\ 0, & x < 0. \end{cases} \quad (\lambda > 0).$$

Distribution function:

$$F(x) = \begin{cases} 1 - e^{-\lambda x}, & x \geq 0, \\ 0, & x < 0. \end{cases}$$

The exponential distribution possesses a **memoryless property**.

Suppose  $\xi \sim E(\lambda)$ . Then for any  $s, t > 0$ ,

$$P(\xi > s + t | \xi > s) = P(\xi > t).$$

**Proof.**

The exponential distribution possesses a **memoryless property**.

Suppose  $\xi \sim E(\lambda)$ . Then for any  $s, t > 0$ ,

$$P(\xi > s + t | \xi > s) = P(\xi > t).$$

**Proof.**  $P(\xi > t) = 1 - F(t) = e^{-\lambda t}$ .

## 2.2 Distribution functions and continuous random variables

## 2.2.3 Typical continuous random variables

$$P(\xi > s + t | \xi > s) = \frac{P(\xi > s + t, \xi > s)}{P(\xi > s)}$$

$$\begin{aligned}P(\xi > s + t | \xi > s) &= \frac{P(\xi > s + t, \xi > s)}{P(\xi > s)} \\&= \frac{P(\xi > s + t)}{P(\xi > s)}\end{aligned}$$

## 2.2 Distribution functions and continuous random variables

## 2.2.3 Typical continuous random variables

$$\begin{aligned}P(\xi > s + t | \xi > s) &= \frac{P(\xi > s + t, \xi > s)}{P(\xi > s)} \\&= \frac{P(\xi > s + t)}{P(\xi > s)} \\&= \frac{e^{-\lambda(t+s)}}{e^{-\lambda s}} \\&= e^{-\lambda t}\end{aligned}$$

## 2.2 Distribution functions and continuous random variables

## 2.2.3 Typical continuous random variables

$$\begin{aligned}P(\xi > s + t | \xi > s) &= \frac{P(\xi > s + t, \xi > s)}{P(\xi > s)} \\&= \frac{P(\xi > s + t)}{P(\xi > s)} \\&= \frac{e^{-\lambda(t+s)}}{e^{-\lambda s}} \\&= e^{-\lambda t} = P(\xi > t).\end{aligned}$$



## Hazard rate functions

Suppose  $X > 0$  (life time of some item) have distribution function  $F$  and density  $p$ . The hazard rate (sometimes called the failure rate) function  $\lambda(t)$  of  $F$  is defined by

$$\lambda(t) = \frac{p(t)}{\overline{F}(t)}, \quad \text{where } \overline{F} = 1 - F.$$

## Hazard rate functions

Suppose  $X > 0$  (life time of some item) have distribution function  $F$  and density  $p$ . The hazard rate (sometimes called the failure rate) function  $\lambda(t)$  of  $F$  is defined by

$$\lambda(t) = \frac{p(t)}{\overline{F}(t)}, \quad \text{where } \overline{F} = 1 - F.$$

Conversely,

$$F(t) = 1 - \exp \left\{ - \int_0^t \lambda(s) ds \right\}.$$

## 2.2 Distribution functions and continuous random variables

## 2.2.3 Typical continuous random variables

$$P(X \in (t, t + dt) | X > t)$$

## 2.2 Distribution functions and continuous random variables

## 2.2.3 Typical continuous random variables

$$\begin{aligned} & P(X \in (t, t + dt) | X > t) \\ = & \frac{P(X \in (t, t + dt), X > t)}{P(X > t)} \\ = & \frac{P(X \in (t, t + dt))}{P(X > t)} \end{aligned}$$

$$\begin{aligned} & P(X \in (t, t + dt) | X > t) \\ = & \frac{P(X \in (t, t + dt), X > t)}{P(X > t)} \\ = & \frac{P(X \in (t, t + dt))}{P(X > t)} \\ \approx & \frac{p(t)}{\bar{F}(t)} dt. \end{aligned}$$

$$\begin{aligned} & P(X \in (t, t + dt) | X > t) \\ = & \frac{P(X \in (t, t + dt), X > t)}{P(X > t)} \\ = & \frac{P(X \in (t, t + dt))}{P(X > t)} \\ \approx & \frac{p(t)}{\bar{F}(t)} dt. \end{aligned}$$

$\lambda(t)$  represents the conditional probability intensity that a  $t$ -unit-old item will fail.

For exponential distribution  $E(\lambda)$ , the hazard function is

$$\begin{aligned}\lambda(t) &= \frac{p(t)}{\bar{F}(t)} \\ &= \frac{\lambda e^{-\lambda t}}{e^{-\lambda t}} = \lambda.\end{aligned}$$

## Example

One often hears that the death rate of a person who smokes is, at each age, twice that of a nonsmoker. What does this mean?



## Example

One often hears that the death rate of a person who smokes is, at each age, twice that of a nonsmoker. What does this mean?

If  $\lambda_s(t)$  denotes the hazard rate of a smoker of age  $t$  and  $\lambda_n(t)$  denotes the hazard rate of a nonsmoker of age  $t$ , then

## Example

One often hears that the death rate of a person who smokes is, at each age, twice that of a nonsmoker. What does this mean?

If  $\lambda_s(t)$  denotes the hazard rate of a smoker of age  $t$  and  $\lambda_n(t)$  denotes the hazard rate of a nonsmoker of age  $t$ , then

$$\lambda_s(t) = 2\lambda_n(t).$$

$$\begin{aligned} p_{non}^{A \rightarrow B} &=: P(\text{A-year-old nonsmoker reaches age B}) \\ &= P\left(\text{nonsmoker's lifetime} > B \right. \\ &\quad \left. | \text{nonsmoker's lifetime} > A \right) \\ &= \frac{1 - F_{non}(B)}{1 - F_{non}(A)} = \frac{\exp \left\{ - \int_0^B \lambda_n(t) dt \right\}}{\exp \left\{ - \int_0^A \lambda_n(t) dt \right\}} \\ &= \exp \left\{ - \int_A^B \lambda_n(t) dt \right\} \end{aligned}$$

$$\begin{aligned} p_s^{A \rightarrow B} &=: P(\text{A-year-old smoker reaches age B}) \\ &= P(\text{smoker's lifetime} > B | \text{smoker's lifetime} > A) \\ &= \exp \left\{ - \int_A^B \lambda_s(t) dt \right\} \\ &= \exp \left\{ -2 \int_A^B \lambda_n(t) dt \right\} \\ &= \left[ \exp \left\{ - \int_A^B \lambda_n(t) dt \right\} \right]^2 = (p_{non}^{A \rightarrow B})^2 \end{aligned}$$

Suppose

$$\lambda_n(t) = \frac{1}{30}, \quad 50 \leq t \leq 60.$$

Then

$$p_{non}^{50 \rightarrow 60} = e^{-1/3} \approx 0.7165,$$

$$p_s^{50 \rightarrow 60} = e^{-2/3} \approx 0.5134.$$

## 4. The Weibull distribution

Density function:

$$p(x) = \begin{cases} \frac{\alpha}{\sigma} \left(\frac{x-\mu}{\sigma}\right)^{\alpha-1} \exp \left\{ - \left(\frac{x-\mu}{\sigma}\right)^{\alpha} \right\}, & x \geq \mu, \\ 0, & x < \mu. \end{cases}.$$

Distribution function:

## 4. The Weibull distribution

Density function:

$$p(x) = \begin{cases} \frac{\alpha}{\sigma} \left(\frac{x-\mu}{\sigma}\right)^{\alpha-1} \exp \left\{ - \left(\frac{x-\mu}{\sigma}\right)^{\alpha} \right\}, & x \geq \mu, \\ 0, & x < \mu. \end{cases}.$$

Distribution function:

$$F(x) = \begin{cases} 1 - \exp \left\{ - \left(\frac{x-\mu}{\sigma}\right)^{\alpha} \right\}, & x \geq \mu, \\ 0, & x < \mu. \end{cases}$$

For Weibull distribution, the hazard function is

$$\lambda(t) = \frac{p(t)}{F(t)} = \frac{\alpha}{\sigma} \left( \frac{t - \mu}{\sigma} \right)^{\alpha-1}.$$

威布尔是瑞典物理学家, 他在1939年研究物质材料的强度时首先提出了这一类分布.



The Weibull distribution is widely used, in the field of life phenomena, as distribution of the lifetime of some object, particularly when the "weakest link" model is appropriate for the object. That is, consider an object consisting of many parts and suppose that the object experiences death (failure) when any of its parts fail. Under these conditions, it has been shown that a Weibull distribution provides a close approximation to the distribution of the lifetime of the item.

## 5. 帕累托(Pareto)分布

若随机变量 $\xi$ 的密度函数为

$$p(x) = \begin{cases} \alpha x_0^\alpha x^{-(\alpha+1)}, & x > x_0, \\ 0, & x \leq x_0, \end{cases}$$

则称 $\xi$ 服从帕累托分布, 其中参数 $x_0 > 0, \alpha > 0$ .

意大利经济学家帕累托首先引入这一分布来描述一个国家中家庭年收入的分布.

## 6. The Gamma distribution

$\xi \sim \Gamma(\lambda, r)$ : Density function


$$p(x) = \begin{cases} \frac{\lambda^r}{\Gamma(r)} x^{r-1} e^{-\lambda x}, & x \geq 0, \\ 0, & x < 0. \end{cases} \quad (\lambda > 0, r > 0)$$

where  $\Gamma(r)$  is the first type of Euler integral. When  $r$  is an integer, we call it an **Erlang distribution**.

$r = 1$ : an exponential distribution.

## 2.2 Distribution functions and continuous random variables

## 2.2.3 Typical continuous random variables



## 7. The Beta distribution

若随机变量 $\xi$ 的密度函数为


$$p(x) = \begin{cases} \frac{1}{B(a,b)} x^{a-1} (1-x)^{b-1}, & 0 \leq x \leq 1, \\ 0, & \text{其它,} \end{cases}$$

则称 $\xi$ 服从参数为 $a$ 和 $b$ 的 $\beta$ 分布, 记作 $\xi \sim \beta(a, b)$ , 其中 $a, b > 0$ ,  $B(a, b) = \int_0^1 x^{a-1} (1-x)^{b-1} dx$  是有名的 $\beta$ 积分, 并且 $B(a, b) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)}$ .

当 $a = b = 1$ 时, 贝塔分布就是区间 $[0, 1]$ 上的均匀分布. 贝塔分布可以用来为取值在有限区间上的随机现象建模.

## 2.2 Distribution functions and continuous random variables

## 2.2.3 Typical continuous random variables



## 8 Cauchy distribution

$$p(x) = \frac{1}{\pi} \frac{1}{1 + (x - \theta)^2}, \quad -\infty < x < \infty.$$