

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

3.1.4 Expectations for functions of random variables

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

3.1.4 Expectations for functions of random variables

The expectation of a function of a discrete random variable:

ξ has the pmf

$$\begin{pmatrix} x_1 & x_2 & \cdots & x_k & \cdots \\ p_1 & p_2 & \cdots & p_k & \cdots \end{pmatrix}.$$

Then for $\eta = f(\xi)$,

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

3.1.4 Expectations for functions of random variables

The expectation of a function of a discrete random variable:

ξ has the pmf

$$\begin{pmatrix} x_1 & x_2 & \cdots & x_k & \cdots \\ p_1 & p_2 & \cdots & p_k & \cdots \end{pmatrix}.$$

Then for $\eta = f(\xi)$,

$$\begin{pmatrix} f(x_1) & f(x_2) & \cdots & f(x_k) & \cdots \\ p_1 & p_2 & \cdots & p_k & \cdots \end{pmatrix}.$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

So the pmf of η is

$$\begin{pmatrix} y_1 & y_2 & \cdots & y_i & \cdots \\ p_1^* & p_2^* & \cdots & p_i^* & \cdots \end{pmatrix}$$

where $p_i^* = \sum_{j:f(x_j)=y_i} p_j$.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

So the pmf of η is

$$\begin{pmatrix} y_1 & y_2 & \cdots & y_i & \cdots \\ p_1^* & p_2^* & \cdots & p_i^* & \cdots \end{pmatrix}$$

where $p_i^* = \sum_{j:f(x_j)=y_i} p_j$. Hence $E\eta$ exists if and only if

$$\sum_i |y_i| p_i^*$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

So the pmf of η is

$$\begin{pmatrix} y_1 & y_2 & \cdots & y_i & \cdots \\ p_1^* & p_2^* & \cdots & p_i^* & \cdots \end{pmatrix}$$

where $p_i^* = \sum_{j:f(x_j)=y_i} p_j$. Hence $E\eta$ exists if and only if

$$\sum_i |y_i| p_i^* = \sum_i \sum_{j:f(x_j)=y_i} |f(x_j)| p_j$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

So the pmf of η is

$$\begin{pmatrix} y_1 & y_2 & \cdots & y_i & \cdots \\ p_1^* & p_2^* & \cdots & p_i^* & \cdots \end{pmatrix}$$

where $p_i^* = \sum_{j:f(x_j)=y_i} p_j$. Hence $E\eta$ exists if and only if

$$\begin{aligned} \sum_i |y_i| p_i^* &= \sum_i \sum_{j:f(x_j)=y_i} |f(x_j)| p_j \\ &= \sum_k |f(x_k)| p_k < \infty. \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Further,

$$E\eta = \sum_i y_i p_i^* = \sum_i \sum_{j: f(x_j)=y_i} f(x_j) p_j = \sum_k f(x_k) p_k.$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Further,

$$E\eta = \sum_i y_i p_i^* = \sum_i \sum_{j: f(x_j)=y_i} f(x_j) p_j = \sum_k f(x_k) p_k.$$

$$Ef(\xi) = \sum_k f(x_k) p_k = \sum_k f(x_k) P(\xi = x_k).$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Theorem Suppose that ξ is a discrete random variable with the distribution $F_\xi(x)$ and

$$P(\xi = x_k) = p_k, \quad k = 1, 2, \dots,$$

$f(x)$ a Borel function on the real line. Let $\eta = f(\xi)$. Then $Ef(\xi)$ exists if and only if

$$\int_{-\infty}^{\infty} |f(x)| dF_\xi(x) = \sum_k |f(x_k)| P(\xi = x_k) < \infty$$

and

$$Ef(\xi) = \sum_k f(x_k) P(\xi = x_k) = \int_{-\infty}^{+\infty} f(x) dF_\xi(x).$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

In general, we have

Theorem 1. Suppose that ξ is a random variable with the distribution $F_\xi(x)$, $f(x)$ a Borel function on the real line. Let $\eta = f(\xi)$. Then

$$Ef(\xi) = \int_{-\infty}^{+\infty} y dF_\eta(y) = \int_{-\infty}^{+\infty} f(x) dF_\xi(x).$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

In general, we have

Theorem 1. Suppose that ξ is a random variable with the distribution $F_\xi(x)$, $f(x)$ a Borel function on the real line. Let $\eta = f(\xi)$. Then

$$Ef(\xi) = \int_{-\infty}^{+\infty} y dF_\eta(y) = \int_{-\infty}^{+\infty} f(x) dF_\xi(x).$$

When ξ has density $p(x)$, then

$$Ef(\xi) = \int_{-\infty}^{+\infty} f(x)p(x)dx.$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

When ξ is a random variable of the general type and $f(x)$ is a general Borel function, the proof of this theorem is due to measure theory. Next, we give a proof for the case when ξ is a continuous type random variable. Let $p(x)$ be the pdf of ξ .

Then

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

When ξ is a random variable of the general type and $f(x)$ is a general Borel function, the proof of this theorem is due to measure theory. Next, we give a proof for the case when ξ is a continuous type random variable. Let $p(x)$ be the pdf of ξ .

Then

$$\begin{aligned} E|\eta| &= \int_0^{\infty} P(|\eta| > y) dy = \int_0^{\infty} P(|f(\xi)| > y) dy \\ &= \int_0^{\infty} \int_{x: |f(x)| > y} p(x) dx dy \\ &= \int_{-\infty}^{\infty} \int_{y: |f(x)| > y, y \geq 0} p(x) dy dx \\ &= \int_{-\infty}^{\infty} |f(x)| p(x) dx. \end{aligned}$$

So $Ef(\xi)$ exists if and only if $\int_{-\infty}^{\infty} |f(x)| p(x) dx < \infty$.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Further,

$$\begin{aligned} E\eta &= \int_0^{\infty} P(\eta > y)dy - \int_0^{\infty} P(-\eta > y)dy \\ &= \int_0^{\infty} \int_{x:f(x)>y} p(x)dx dy - \int_0^{\infty} \int_{x:-f(x)>y} p(x)dx dy \\ &= \int_{-\infty}^{\infty} \left[\int_{y:f(x)>y, y\geq 0} dy - \int_{y:-f(x)>y, y\geq 0} dy \right] p(x)dx \\ &= \int_{-\infty}^{\infty} f(x)p(x)dx. \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Further,

$$\begin{aligned} E\eta &= \int_0^{\infty} P(\eta > y)dy - \int_0^{\infty} P(-\eta > y)dy \\ &= \int_0^{\infty} \int_{x:f(x)>y} p(x)dx dy - \int_0^{\infty} \int_{x:-f(x)>y} p(x)dx dy \\ &= \int_{-\infty}^{\infty} \left[\int_{y:f(x)>y, y\geq 0} dy - \int_{y:-f(x)>y, y\geq 0} dy \right] p(x)dx \\ &= \int_{-\infty}^{\infty} f(x)p(x)dx. \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Theorem 1 tells us that if ξ and η have the same distribution function, then

$$Ef(\xi) = Ef(\eta).$$

On the contrary, if the above equality holds for any bounded continuous function f , then ξ and η have the same distribution function.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

In fact, for any z and $\epsilon > 0$, let $f(x)$ be a continuous function such that $f(x) = 1$, $0 \leq f(x) \leq 1$ and $f(x) = 0$ on $(-\infty, z]$, $(z, z + \epsilon]$ and $(z + \epsilon, \infty)$, respectively. Then

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

In fact, for any z and $\epsilon > 0$, let $f(x)$ be a continuous function such that $f(x) = 1$, $0 \leq f(x) \leq 1$ and $f(x) = 0$ on $(-\infty, z]$, $(z, z + \epsilon]$ and $(z + \epsilon, \infty)$, respectively. Then

$$\begin{aligned} F_{\xi}(z) &= \int_{-\infty}^z f(x) dF_{\xi}(x) \leq \int_{-\infty}^{\infty} f(x) dF_{\xi}(x) \\ &= \int_{-\infty}^{\infty} f(x) dF_{\eta}(x) = \int_{-\infty}^{z+\epsilon} f(x) dF_{\eta}(x) \\ &\leq \int_{-\infty}^{z+\epsilon} dF_{\eta}(x) = F_{\eta}(z + \epsilon). \end{aligned}$$

Letting $\epsilon \rightarrow 0$ yields $F_{\xi}(z) \leq F_{\eta}(z)$.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

In fact, for any z and $\epsilon > 0$, let $f(x)$ be a continuous function such that $f(x) = 1$, $0 \leq f(x) \leq 1$ and $f(x) = 0$ on $(-\infty, z]$, $(z, z + \epsilon]$ and $(z + \epsilon, \infty)$, respectively. Then

$$\begin{aligned} F_{\xi}(z) &= \int_{-\infty}^z f(x) dF_{\xi}(x) \leq \int_{-\infty}^{\infty} f(x) dF_{\xi}(x) \\ &= \int_{-\infty}^{\infty} f(x) dF_{\eta}(x) = \int_{-\infty}^{z+\epsilon} f(x) dF_{\eta}(x) \\ &\leq \int_{-\infty}^{z+\epsilon} dF_{\eta}(x) = F_{\eta}(z + \epsilon). \end{aligned}$$

Letting $\epsilon \rightarrow 0$ yields $F_{\xi}(z) \leq F_{\eta}(z)$. Similarly, $F_{\eta}(z) \leq F_{\xi}(z)$. So ξ and η have the same distribution function.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Example. (Stein's Lemma) (i) Let $\xi \sim N(0, 1)$, and g be differentiable function satisfying $|g(x)| \leq c_1 e^{c_2|x|}$ and $|g'(x)| \leq c_1 e^{c_2|x|}$ for some $c_1 > 0, c_2 > 0$. Prove

$$E[g(\xi)\xi] = E g'(\xi).$$

(ii)* On the contrary, if the above equality holds for any bounded continuous function $g(x)$ with bounded derivation, then $\xi \sim N(0, 1)$.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Proof. For (i), we have

$$\begin{aligned} & E[g(\xi)\xi] \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} g(x)x \exp\left\{-\frac{x^2}{2}\right\} dx \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Proof. For (i), we have

$$\begin{aligned} & E[g(\xi)\xi] \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} g(x)x \exp\left\{-\frac{x^2}{2}\right\} dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} g(x) d\left[-\exp\left\{-\frac{x^2}{2}\right\}\right]. \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Use integration by parts to get

$$\begin{aligned} & E[g(\xi)\xi] \\ &= \frac{1}{\sqrt{2\pi}} \left[-g(x) \exp\left\{-\frac{x^2}{2}\right\} \Big|_{-\infty}^{\infty} \right. \\ & \quad \left. + \int_{-\infty}^{\infty} g'(x) \exp\left\{-\frac{x^2}{2}\right\} dx \right] \\ &= \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Use integration by parts to get

$$\begin{aligned} & E[g(\xi)\xi] \\ &= \frac{1}{\sqrt{2\pi}} \left[-g(x) \exp\left\{-\frac{x^2}{2}\right\} \Big|_{-\infty}^{\infty} \right. \\ & \quad \left. + \int_{-\infty}^{\infty} g'(x) \exp\left\{-\frac{x^2}{2}\right\} dx \right] \\ &= E g'(\xi). \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

For (ii), Let $\eta \sim N(0, 1)$. It is sufficient to show that $Eh(\xi) = Eh(\eta)$ holds for any bounded continuous function $h(x)$.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

For (ii), Let $\eta \sim N(0, 1)$. It is sufficient to show that $Eh(\xi) = Eh(\eta)$ holds for any bounded continuous function $h(x)$. Without loss of generality, we assume $0 \leq h(x) \leq 1$. Let $g(x)$ satisfy

$$h(x) - Eh(\eta) = g'(x) - xg(x).$$

The above equation is called Stein's equation.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

For (ii), Let $\eta \sim N(0, 1)$. It is sufficient to show that $Eh(\xi) = Eh(\eta)$ holds for any bounded continuous function $h(x)$. Without loss of generality, we assume $0 \leq h(x) \leq 1$. Let $g(x)$ satisfy

$$h(x) - Eh(\eta) = g'(x) - xg(x).$$

The above equation is called Stein's equation. It can be verified that

$$g(x) = e^{\frac{x^2}{2}} \int_{-\infty}^x [h(u) - Eh(\eta)] e^{-\frac{u^2}{2}} du$$

is a solution of the equation, and, both $g(x)$ and $g'(x)$ are bounded continuous function.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

So

$$\begin{aligned} Eh(\xi) &= \int_{-\infty}^{\infty} [g'(x) - xg(x) + Eh(\eta)] dF_{\xi}(x) \\ &= \int_{-\infty}^{\infty} g'(x) dF_{\xi}(x) - \int_{-\infty}^{\infty} xg(x) dF_{\xi}(x) + Eh(\eta) \int_{-\infty}^{\infty} dF_{\xi}(x) \\ &= Eg'(\xi) - E[\xi g(\xi)] + Eh(\eta) = Eh(\eta). \end{aligned}$$

The proof is completed. In the second equality above, the linearity of the Stieltjes integral.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Similarly,

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Similarly,

$$E[\xi g(\xi - 1)] = E[\lambda g(\xi)], \quad \forall g(\text{bounded}) \Leftrightarrow \xi \sim P(\lambda).$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Similarly,

$$E[\xi g(\xi - 1)] = E[\lambda g(\xi)], \quad \forall g(\text{bounded}) \Leftrightarrow \xi \sim P(\lambda).$$

Stein-Chen method.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

In general, suppose $(\xi_1, \dots, \xi_n) \sim F(x_1, \dots, x_n)$.

Also, assume that $g(x_1, \dots, x_n)$ is a Borel function, then

$$Eg(\xi_1, \dots, \xi_n) = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} g(x_1, \dots, x_n) dF(x_1, \dots, x_n).$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

In general, suppose $(\xi_1, \dots, \xi_n) \sim F(x_1, \dots, x_n)$. Also, assume that $g(x_1, \dots, x_n)$ is a Borel function, then

$$Eg(\xi_1, \dots, \xi_n) = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} g(x_1, \dots, x_n) dF(x_1, \dots, x_n).$$

If $(\xi_1, \xi_2, \dots, \xi_n)$ has pmf

$P(\xi_1 = x_1(i_1), \xi_2 = x_2(i_2), \dots, \xi_n = x_n(i_n)) = p_{i_1 i_2 \dots i_n}$, then

$$Eg(\xi_1, \xi_2, \dots, \xi_n) = \sum_{i_1, i_2, \dots, i_n} g(x_1(i_1), x_2(i_2), \dots, x_n(i_n)) p_{i_1 i_2 \dots i_n};$$

If $(\xi_1, \xi_2, \dots, \xi_n)$ has pdf $p(x_1, x_2, \dots, x_n)$, then

$$\begin{aligned} & Eg(\xi_1, \xi_2, \dots, \xi_n) \\ &= \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} g(x_1, x_2, \dots, x_n) p(x_1, x_2, \dots, x_n) dx_1 \cdots dx_n. \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Here, the multi-variable Stieltjes integral is defined similarly as in the one-variable case. For example

$$\begin{aligned} & \int_{a_1}^{b_1} \cdots \int_{a_n}^{b_n} g(x_1, \dots, x_n) dF(x_1, \dots, x_n) \\ &= \lim \sum_{k_1, \dots, k_n} g(x_1(k_1), \dots, x_n(k_n)) \Delta F(x_1(k_1), \dots, x_n(k_n)), \end{aligned}$$

where $x_i(1), x_i(2), \dots$ is a partition of $(a_i, b_i]$,

$\Delta F(x_1(k_1), \dots, x_n(k_n))$ is the probability that (ξ_1, \dots, ξ_n) falls in $(x_1(k_1), x_1(k_1 + 1)] \times \cdots \times (x_n(k_n), x_n(k_n + 1)]$.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

In particular, we have

$$E\xi_i = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} x_i dF(x_1, \cdots, x_n) = \int_{-\infty}^{\infty} x dF_i(x),$$

where $F_i(x)$ is the distribution function of ξ_i .

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

For $F(x, y)$ it follows

$$E\xi\eta = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy dF(x, y)$$

and

$$E\xi^2 = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x^2 dF(x, y),$$

etc.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Example. Suppose R and Θ are indept.,
 $\Theta \sim U(0, 2\pi)$, $R \sim \text{Rayleigh}$. Find $Ee^{R \sin \Theta}$.

Solution.

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Example. Suppose R and Θ are indept.,
 $\Theta \sim U(0, 2\pi)$, $R \sim \text{Rayleigh}$. Find $Ee^{R \sin \Theta}$.

Solution.

$$Ee^{R \sin \Theta}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Example. Suppose R and Θ are indept.,
 $\Theta \sim U(0, 2\pi)$, $R \sim \text{Rayleigh}$. Find $Ee^{R \sin \Theta}$.

Solution.

$$Ee^{R \sin \Theta} = \int_0^\infty \int_0^{2\pi} e^{r \sin \theta} \textcolor{red}{r} e^{-r^2/2} \textcolor{blue}{\frac{1}{2\pi}} d\theta dr$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Example. Suppose R and Θ are indept.,
 $\Theta \sim U(0, 2\pi)$, $R \sim \text{Rayleigh}$. Find $Ee^{R \sin \Theta}$.

Solution.

$$\begin{aligned} Ee^{R \sin \Theta} &= \int_0^\infty \int_0^{2\pi} e^{r \sin \theta} \textcolor{red}{r} \textcolor{red}{e^{-r^2/2}} \textcolor{blue}{\frac{1}{2\pi}} d\theta dr \\ &= \iint e^x \frac{1}{2\pi} e^{-\frac{x^2+y^2}{2}} dx dy (\text{极坐标变换}) \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Example. Suppose R and Θ are indept.,
 $\Theta \sim U(0, 2\pi)$, $R \sim \text{Rayleigh}$. Find $Ee^{R \sin \Theta}$.

Solution.

$$\begin{aligned}
 Ee^{R \sin \Theta} &= \int_0^\infty \int_0^{2\pi} e^{r \sin \theta} \color{red}{r} e^{-r^2/2} \color{blue}{\frac{1}{2\pi}} d\theta dr \\
 &= \iint e^x \frac{1}{2\pi} e^{-\frac{x^2+y^2}{2}} dx dy (\text{极坐标变换}) \\
 &= \int_{-\infty}^\infty e^x \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx
 \end{aligned}$$

3.1 Mathematical expectation

3.1.4 Expectations for functions of random variables

Example. Suppose R and Θ are indept.,
 $\Theta \sim U(0, 2\pi)$, $R \sim \text{Rayleigh}$. Find $Ee^{R \sin \Theta}$.

Solution.

$$\begin{aligned}
 Ee^{R \sin \Theta} &= \int_0^\infty \int_0^{2\pi} e^{r \sin \theta} \color{red}{r} e^{-r^2/2} \color{blue}{\frac{1}{2\pi}} d\theta dr \\
 &= \iint e^x \frac{1}{2\pi} e^{-\frac{x^2+y^2}{2}} dx dy (\text{极坐标变换}) \\
 &= \int_{-\infty}^\infty e^x \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx \\
 &= e^{\frac{1}{2}} \int_{-\infty}^\infty \frac{1}{\sqrt{2\pi}} e^{-\frac{(x-1)^2}{2}} dx = e^{\frac{1}{2}}.
 \end{aligned}$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

3.1.5 Basic properties of expectations-examples

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

3.1.5 Basic properties of expectations-examples

$$\textcircled{1} \quad a \leq \xi \leq b \implies a \leq E\xi \leq b.$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

3.1.5 Basic properties of expectations-examples

$$\textcircled{1} \quad a \leq \xi \leq b \implies a \leq E\xi \leq b.$$

$$\textcircled{2} \quad E\xi_1, \dots, E\xi_n \text{ exist} \implies$$

$$E\left(\sum_{i=1}^n c_i \xi_i + b\right) = \sum_{i=1}^n c_i E\xi_i + b.$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

3.1.5 Basic properties of expectations-examples

$$\textcircled{1} \quad a \leq \xi \leq b \implies a \leq E\xi \leq b.$$

$$\textcircled{2} \quad E\xi_1, \dots, E\xi_n \text{ exist} \implies$$

$$E\left(\sum_{i=1}^n c_i \xi_i + b\right) = \sum_{i=1}^n c_i E\xi_i + b.$$

$$\textcircled{3} \quad \xi_1, \dots, \xi_n \text{ indept. and expectations exist} \implies$$

$$E(\xi_1 \cdots \xi_n) = E\xi_1 \cdots E\xi_n.$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

3.1.5 Basic properties of expectations-examples

$$\textcircled{1} \quad a \leq \xi \leq b \implies a \leq E\xi \leq b.$$

$$\textcircled{2} \quad E\xi_1, \dots, E\xi_n \text{ exist} \implies$$

$$E\left(\sum_{i=1}^n c_i \xi_i + b\right) = \sum_{i=1}^n c_i E\xi_i + b.$$

$$\textcircled{3} \quad \xi_1, \dots, \xi_n \text{ indept. and expectations exist} \implies$$

$$E(\xi_1 \cdots \xi_n) = E\xi_1 \cdots E\xi_n.$$

$$\textcircled{4} \quad \text{If } \xi \leq \eta \text{ and the expectations of } \xi \text{ and } \eta \text{ exist,} \\ \text{then } E\xi \leq E\eta.$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

Example 12. Suppose that ξ follows the binomial distribution $B(n, p)$, find $E\xi$.

Example 12. Suppose that ξ follows the binomial distribution $B(n, p)$, find $E\xi$.

Solution. Consider a Bernoulli trial and set $p = P(A)$ and

$$\xi_i = \begin{cases} 1, & A \text{ occurs in the } i\text{-th trial,} \\ 0, & A \text{ does not occur in the } i\text{-th trial.} \end{cases}$$

Example 12. Suppose that ξ follows the binomial distribution $B(n, p)$, find $E\xi$.

Solution. Consider a Bernoulli trial and set $p = P(A)$ and

$$\xi_i = \begin{cases} 1, & A \text{ occurs in the } i\text{-th trial,} \\ 0, & A \text{ does not occur in the } i\text{-th trial.} \end{cases}$$

Thus ξ_i follows 0-1 distribution, $E\xi_i = p$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

Example 12. Suppose that ξ follows the binomial distribution $B(n, p)$, find $E\xi$.

Solution. Consider a Bernoulli trial and set $p = P(A)$ and

$$\xi_i = \begin{cases} 1, & A \text{ occurs in the } i\text{-th trial,} \\ 0, & A \text{ does not occur in the } i\text{-th trial.} \end{cases}$$

Thus ξ_i follows 0-1 distribution, $E\xi_i = p$ and $\xi = \sum_{i=1}^n \xi_i$.

Example 12. Suppose that ξ follows the binomial distribution $B(n, p)$, find $E\xi$.

Solution. Consider a Bernoulli trial and set $p = P(A)$ and

$$\xi_i = \begin{cases} 1, & A \text{ occurs in the } i\text{-th trial,} \\ 0, & A \text{ does not occur in the } i\text{-th trial.} \end{cases}$$

Thus ξ_i follows 0-1 distribution, $E\xi_i = p$ and $\xi = \sum_{i=1}^n \xi_i$. Hence $E\xi = np$.

Example 13. Suppose that

$$P(\xi = m) = \frac{\binom{M}{m} \binom{N-M}{n-m}}{\binom{N}{n}}, \quad m = 0, 1, \dots, n.$$

($n \leq M \leq N$). Find $E\xi$.

Solution. Design a sampling without replacement.

Example 13. Suppose that

$$P(\xi = m) = \frac{\binom{M}{m} \binom{N-M}{n-m}}{\binom{N}{n}}, \quad m = 0, 1, \dots, n.$$

($n \leq M \leq N$). Find $E\xi$.

Solution. Design a sampling without replacement.

Let ξ_i be the number of defective goods in the i -th draw. Then $\xi = \sum_{i=1}^n \xi_i$.

Example 13. Suppose that

$$P(\xi = m) = \frac{\binom{M}{m} \binom{N-M}{n-m}}{\binom{N}{n}}, \quad m = 0, 1, \dots, n.$$

($n \leq M \leq N$). Find $E\xi$.

Solution. Design a sampling without replacement.

Let ξ_i be the number of defective goods in the i -th draw. Then $\xi = \sum_{i=1}^n \xi_i$. It is known that

$P(\xi_i = 1) = M/N$, so $E\xi_i = M/N$.

Example 13. Suppose that

$$P(\xi = m) = \frac{\binom{M}{m} \binom{N-M}{n-m}}{\binom{N}{n}}, \quad m = 0, 1, \dots, n.$$

($n \leq M \leq N$). Find $E\xi$.

Solution. Design a sampling without replacement.

Let ξ_i be the number of defective goods in the i -th draw. Then $\xi = \sum_{i=1}^n \xi_i$. It is known that

$P(\xi_i = 1) = M/N$, so $E\xi_i = M/N$. Hence

$$E\xi = \sum_{i=1}^n E\xi_i = \frac{nM}{N}.$$

Example 14. Suppose that ξ_1, \dots, ξ_n are independent identically distributed positive random variables with a common density function $f(x)$. Show for any $1 \leq k \leq n$,

$$E \frac{\xi_1 + \dots + \xi_k}{\xi_1 + \dots + \xi_n} = \frac{k}{n}.$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

Proof.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

Proof. Notice that $\frac{\xi_k}{\xi_1 + \cdots + \xi_n}$ is positive and

$$\begin{aligned} & E \frac{\xi_k}{\xi_1 + \cdots + \xi_n} \\ = & \int_0^\infty \cdots \int_0^\infty \frac{x_k}{x_1 + \cdots + x_n} f(x_1) \cdots f(x_n) dx_1 \cdots dx_n \end{aligned}$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

Proof. Notice that $\frac{\xi_k}{\xi_1 + \dots + \xi_n}$ is positive and

$$\begin{aligned} & E \frac{\xi_k}{\xi_1 + \dots + \xi_n} \\ = & \int_0^\infty \dots \int_0^\infty \frac{x_k}{x_1 + \dots + x_n} f(x_1) \dots f(x_n) dx_1 \dots dx_n \\ = & \int_0^\infty \dots \int_0^\infty \frac{y_1}{y_1 + \dots + y_n} f(y_1) \dots f(y_n) dy_1 \dots dy_n \end{aligned}$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

Proof. Notice that $\frac{\xi_k}{\xi_1 + \dots + \xi_n}$ is positive and

$$\begin{aligned} & E \frac{\xi_k}{\xi_1 + \dots + \xi_n} \\ = & \int_0^\infty \dots \int_0^\infty \frac{x_k}{x_1 + \dots + x_n} f(x_1) \dots f(x_n) dx_1 \dots dx_n \\ = & \int_0^\infty \dots \int_0^\infty \frac{y_1}{y_1 + \dots + y_n} f(y_1) \dots f(y_n) dy_1 \dots dy_n \\ = & E \frac{\xi_1}{\xi_1 + \dots + \xi_n}. \end{aligned}$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

On the other hand,

$$E \frac{\xi_1}{\xi_1 + \cdots + \xi_n} + \cdots + E \frac{\xi_n}{\xi_1 + \cdots + \xi_n}$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

On the other hand,

$$\begin{aligned} & E \frac{\xi_1}{\xi_1 + \cdots + \xi_n} + \cdots + E \frac{\xi_n}{\xi_1 + \cdots + \xi_n} \\ = & E \frac{\xi_1 + \cdots + \xi_n}{\xi_1 + \cdots + \xi_n} = 1. \end{aligned}$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

On the other hand,

$$\begin{aligned} & E \frac{\xi_1}{\xi_1 + \cdots + \xi_n} + \cdots + E \frac{\xi_n}{\xi_1 + \cdots + \xi_n} \\ &= E \frac{\xi_1 + \cdots + \xi_n}{\xi_1 + \cdots + \xi_n} = 1. \end{aligned}$$

It follows that

$$E \frac{\xi_1}{\xi_1 + \cdots + \xi_n} = \cdots = E \frac{\xi_n}{\xi_1 + \cdots + \xi_n} = \frac{1}{n}.$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

On the other hand,

$$\begin{aligned} & E \frac{\xi_1}{\xi_1 + \cdots + \xi_n} + \cdots + E \frac{\xi_n}{\xi_1 + \cdots + \xi_n} \\ &= E \frac{\xi_1 + \cdots + \xi_n}{\xi_1 + \cdots + \xi_n} = 1. \end{aligned}$$

It follows that

$$E \frac{\xi_1}{\xi_1 + \cdots + \xi_n} = \cdots = E \frac{\xi_n}{\xi_1 + \cdots + \xi_n} = \frac{1}{n}.$$

Hence

$$E \frac{\xi_1 + \cdots + \xi_k}{\xi_1 + \cdots + \xi_n}$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

On the other hand,

$$\begin{aligned} & E \frac{\xi_1}{\xi_1 + \cdots + \xi_n} + \cdots + E \frac{\xi_n}{\xi_1 + \cdots + \xi_n} \\ &= E \frac{\xi_1 + \cdots + \xi_n}{\xi_1 + \cdots + \xi_n} = 1. \end{aligned}$$

It follows that

$$E \frac{\xi_1}{\xi_1 + \cdots + \xi_n} = \cdots = E \frac{\xi_n}{\xi_1 + \cdots + \xi_n} = \frac{1}{n}.$$

Hence

$$\begin{aligned} & E \frac{\xi_1 + \cdots + \xi_k}{\xi_1 + \cdots + \xi_n} \\ &= E \frac{\xi_1}{\xi_1 + \cdots + \xi_n} + \cdots + E \frac{\xi_k}{\xi_1 + \cdots + \xi_n} = \frac{k}{n}. \end{aligned}$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

Example

A grove of 52 trees is arranged in a circular fashion. If a total of 15 chipmunks (花栗鼠) live in these trees, show that there is a group of 7 consecutive trees that together house at least 3 chipmunks.

Example

A grove of 52 trees is arranged in a circular fashion. If a total of 15 chipmunks (花栗鼠) live in these trees, show that there is a group of 7 consecutive trees that together house at least 3 chipmunks.

解：给定树 j , 记它连同它旁边按顺时针方向排列的6颗构成一个邻域 U_j , 生活在 U_j 中的chipmunks个数记为 Y_j .

Example

A grove of 52 trees is arranged in a circular fashion. If a total of 15 chipmunks (花栗鼠) live in these trees, show that there is a group of 7 consecutive trees that together house at least 3 chipmunks.

解: 给定树 j , 记它连同它旁边按顺时针方向排列的6颗构成一个邻域 U_j , 生活在 U_j 中的chipmunks个数记为 Y_j . 我们只要证明 $EY_j > 2 \forall j$, 就说明了存在一个树, 使得至少有3个chipmunks 生活在此树的邻域中.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

事实上, 如果 $Y_j \leq 2 \ \forall j$, 那么 $\sum_{j=1}^{52} Y_j \leq 2 * 52$.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

事实上, 如果 $Y_j \leq 2 \ \forall j$, 那么 $\sum_{j=1}^{52} Y_j \leq 2 * 52$. 从而

$$E \left[\sum_{j=1}^{52} Y_j \right] \leq 2 * 52.$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

事实上, 如果 $Y_j \leq 2 \ \forall j$, 那么 $\sum_{j=1}^{52} Y_j \leq 2 * 52$. 从而

$$E \left[\sum_{j=1}^{52} Y_j \right] \leq 2 * 52.$$

这与

$$\sum_{j=1}^{52} EY_j > 2 * 52$$

矛盾.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

下面求 EY_j .

$$\text{令 } X_i = \begin{cases} 1, & \text{if chipmunk } i \text{ live in } U_j, \\ 0, & \text{otherwise.} \end{cases}$$

$$\text{则 } Y_j = \sum_{i=1}^{15} X_i.$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

下面求 EY_j .

$$\text{令 } X_i = \begin{cases} 1, & \text{if chipmunk } i \text{ live in } U_j, \\ 0, & \text{otherwise.} \end{cases}$$

则 $Y_j = \sum_{i=1}^{15} X_i$. 因为

$$EX_i = P(X_i = 1) = \frac{7}{52}.$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

下面求 EY_j .

$$\text{令 } X_i = \begin{cases} 1, & \text{if chipmunk } i \text{ live in } U_j, \\ 0, & \text{otherwise.} \end{cases}$$

则 $Y_j = \sum_{i=1}^{15} X_i$. 因为

$$EX_i = P(X_i = 1) = \frac{7}{52}.$$

$$\text{所以 } EY_j = \sum_{i=1}^{15} EX_i = \frac{105}{52} > 2.$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

Example 15. Make a census on some kind of disease in a community with large population. Now check blood for N citizens in two ways: (1) each person each time, so need check N times. (2) check the mixture of bloods of a group of k people. If the outcome reports no virus, that means all these k people are not of this disease; while if the outcome reports virus, then each person from this group is checked again, so k people need check $k + 1$ times in this way. Which way may decrease the number of checks?

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

Solution. Consider the second way. Denote by ξ the number of times each person needs check in a group of k people in the second way. Then

$$\xi = \begin{cases} 1/k, & \text{none of } k \text{ people is sick} \\ (k+1)/k, & \text{at least one of } k \text{ people is sick.} \end{cases}$$

So

Solution. Consider the second way. Denote by ξ the number of times each person needs check in a group of k people in the second way. Then

$$\xi = \begin{cases} 1/k, & \text{none of } k \text{ people is sick} \\ (k+1)/k, & \text{at least one of } k \text{ people is sick.} \end{cases}$$

So $P(\xi = \frac{1}{k}) = (1-p)^k$,
 $P(\xi = 1 + \frac{1}{k}) = 1 - (1-p)^k$.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

Solution. Consider the second way. Denote by ξ the number of times each person needs check in a group of k people in the second way. Then

$$\xi = \begin{cases} 1/k, & \text{none of } k \text{ people is sick} \\ (k+1)/k, & \text{at least one of } k \text{ people is sick.} \end{cases}$$

So $P(\xi = \frac{1}{k}) = (1-p)^k$,

$P(\xi = 1 + \frac{1}{k}) = 1 - (1-p)^k$. Hence

$$\begin{aligned} E\xi &= \frac{1}{k}(1-p)^k + (1 + \frac{1}{k})(1 - (1-p)^k) \\ &= 1 - (1-p)^k + \frac{1}{k}. \end{aligned}$$

Basic properties of expectations (continue)

Corollary

Suppose $|\xi| \leq \eta$, $E\eta < \infty$. Then $E\xi$ exists and $|E\xi| \leq E|\xi| \leq E\eta$.

Basic properties of expectations (continue)

Corollary

Suppose $|\xi| \leq \eta$, $E\eta < \infty$. Then $E\xi$ exists and $|E\xi| \leq E|\xi| \leq E\eta$.

Proof. For $M > 0$, let $\xi_M = |\xi|$ if $|\xi| \leq M$, and 0 if $|\xi| > M$. Then $0 \leq \xi_M \leq M$. By Property 1, $0 \leq E\xi_M \leq M$.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

Basic properties of expectations (continue)

Corollary

Suppose $|\xi| \leq \eta$, $E\eta < \infty$. Then $E\xi$ exists and $|E\xi| \leq E|\xi| \leq E\eta$.

Proof. For $M > 0$, let $\xi_M = |\xi|$ if $|\xi| \leq M$, and 0 if $|\xi| > M$. Then $0 \leq \xi_M \leq M$. By Property 1, $0 \leq E\xi_M \leq M$. So $E\xi_M$, $E\eta$ exist, and $\xi_M \leq \eta$. It follows that

$$\int_{-M}^M |x| dF_{\xi}(x) = E\xi_M \leq E\eta$$

by Property 4.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

Basic properties of expectations (continue)

Corollary

Suppose $|\xi| \leq \eta$, $E\eta < \infty$. Then $E\xi$ exists and $|E\xi| \leq E|\xi| \leq E\eta$.

Proof. For $M > 0$, let $\xi_M = |\xi|$ if $|\xi| \leq M$, and 0 if $|\xi| > M$. Then $0 \leq \xi_M \leq M$. By Property 1, $0 \leq E\xi_M \leq M$. So $E\xi_M$, $E\eta$ exist, and $\xi_M \leq \eta$. It follows that

$$\int_{-M}^M |x| dF_{\xi}(x) = E\xi_M \leq E\eta$$

by Property 4. Hence $E|\xi| = \int_{-\infty}^{\infty} |x| dF_{\xi}(x) \leq E\eta < \infty$.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

Basic properties of expectations (continue)

Corollary

Suppose $|\xi| \leq \eta$, $E\eta < \infty$. Then $E\xi$ exists and $|E\xi| \leq E|\xi| \leq E\eta$.

Proof. For $M > 0$, let $\xi_M = |\xi|$ if $|\xi| \leq M$, and 0 if $|\xi| > M$. Then $0 \leq \xi_M \leq M$. By Property 1, $0 \leq E\xi_M \leq M$. So $E\xi_M$, $E\eta$ exist, and $\xi_M \leq \eta$. It follows that

$$\int_{-M}^M |x| dF_{\xi}(x) = E\xi_M \leq E\eta$$

by Property 4. Hence $E|\xi| = \int_{-\infty}^{\infty} |x| dF_{\xi}(x) \leq E\eta < \infty$.

Finally, since $-|\xi| \leq \xi \leq |\xi|$, by Property 4 we have

$-E|\xi| \leq E\xi \leq E|\xi|$. The proof is completed.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

Corollary

Let $p > 1$. If $E|\xi|^p$ exists, then $E|\xi|$ exists.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

Corollary

Let $p > 1$. If $E|\xi|^p$ exists, then $E|\xi|$ exists.

Proof. Since $|\xi| \leq 1 + |\xi|^p$, $E|\xi| \leq 1 + E|\xi|^p$.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

⑤ **Markov inequality:** If $E|\xi|$ exists, then

$$P(|\xi| \geq \epsilon) \leq \frac{E|\xi|}{\epsilon}, \quad \forall \epsilon > 0.$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

⑤ **Markov inequality:** If $E|\xi|$ exists, then

$$P(|\xi| \geq \epsilon) \leq \frac{E|\xi|}{\epsilon}, \quad \forall \epsilon > 0.$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

5 **Markov inequality:** If $E|\xi|$ exists, then

$$P(|\xi| \geq \epsilon) \leq \frac{E|\xi|}{\epsilon}, \quad \forall \epsilon > 0.$$

In fact, let

$$\eta = \begin{cases} 1, & \text{if } |\xi| \geq \epsilon, \\ 0, & \text{for otherwise.} \end{cases} \quad \text{Then } \eta \leq \frac{|\xi|}{\epsilon}.$$

By Property 4, we have

$$P(|\xi| \geq \epsilon) = E\eta \leq E\left[\frac{|\xi|}{\epsilon}\right] = \frac{E|\xi|}{\epsilon}.$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

6 $P(\xi = 0) = 1$ if and only if $E|\xi| = 0$.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

6 $P(\xi = 0) = 1$ if and only if $E|\xi| = 0$.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

⑥ $P(\xi = 0) = 1$ if and only if $E|\xi| = 0$.

In fact, the "only if" part is obvious. For the "if" part, by Property 5 we have

$$P(|\xi| \geq \epsilon) = 0 \text{ for all } \epsilon > 0.$$

So $P(|\xi| > 0) = 0$.

Convergence theorems

7 (Monotone convergence theorem). If $0 \leq \xi_n(\omega) \nearrow \xi(\omega)$, then

$$\lim_{n \rightarrow \infty} E\xi_n = E\xi. \quad (*)$$

If $0 \leq \xi_n(\omega) \searrow 0$, and $E\xi_n$ s are finite, then

$$\lim_{n \rightarrow \infty} E\xi_n = 0.$$

Convergence theorems

- 7 (*Monotone convergence theorem*). If $0 \leq \xi_n(\omega) \nearrow \xi(\omega)$, then

$$\lim_{n \rightarrow \infty} E\xi_n = E\xi. \quad (*)$$

If $0 \leq \xi_n(\omega) \searrow 0$, and $E\xi_n$ s are finite, then

$$\lim_{n \rightarrow \infty} E\xi_n = 0.$$

- 8 (*Dominated convergence theorem*). If $\xi_n(\omega) \rightarrow \xi(\omega)$, $|\xi_n| \leq \eta$ and $E\eta < \infty$, then $(*)$ holds.

Convergence theorems

- 7 (*Monotone convergence theorem*). If $0 \leq \xi_n(\omega) \nearrow \xi(\omega)$, then

$$\lim_{n \rightarrow \infty} E\xi_n = E\xi. \quad (*)$$

If $0 \leq \xi_n(\omega) \searrow 0$, and $E\xi_n$ s are finite, then

$$\lim_{n \rightarrow \infty} E\xi_n = 0.$$

- 8 (*Dominated convergence theorem*). If $\xi_n(\omega) \rightarrow \xi(\omega)$, $|\xi_n| \leq \eta$ and $E\eta < \infty$, then $(*)$ holds.
- 9 (*Bounded convergence theorem*). If $\xi_n(\omega) \rightarrow \xi(\omega)$ and $|\xi_n| \leq M < \infty$, then $(*)$ holds.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

证明: 先证明有界收敛定理.

首先, 已知 $|\xi_n| \leq M$, $|\xi| \leq M$. 由性质1, $E\xi_n$, $E\xi$ 存在, 并且 $|E\xi_n - E\xi| \leq E|\xi_n - \xi|$.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

证明: 先证明有界收敛定理.

首先, 已知 $|\xi_n| \leq M$, $|\xi| \leq M$. 由性质1, $E\xi_n$, $E\xi$ 存在, 并且 $|E\xi_n - E\xi| \leq E|\xi_n - \xi|$. 对任给的 $\epsilon > 0$, 记 $A_n = \{|\xi_n - \xi| > \epsilon\}$. 在 A_n^c 上有 $|\xi_n - \xi| \leq \epsilon$. 而在 A_n 上有 $|\xi_n - \xi| \leq 2M$.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

证明: 先证明有界收敛定理.

首先, 已知 $|\xi_n| \leq M$, $|\xi| \leq M$. 由性质1, $E\xi_n$, $E\xi$ 存在, 并且 $|E\xi_n - E\xi| \leq E|\xi_n - \xi|$. 对任给的 $\epsilon > 0$, 记 $A_n = \{|\xi_n - \xi| > \epsilon\}$. 在 A_n^c 上有 $|\xi_n - \xi| \leq \epsilon$. 而在 A_n 上有 $|\xi_n - \xi| \leq 2M$. 所以

$$|\xi_n - \xi| \leq \epsilon + 2MI_{A_n}.$$

因此

$$|E\xi_n - E\xi| \leq E|\xi_n - \xi| \leq \epsilon + 2MP(A_n).$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

另一方面, 由于 $\xi_n(\omega) \rightarrow \xi(\omega)$, 所以 $\lim_{n \rightarrow \infty} A_n = \emptyset$.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

另一方面, 由于 $\xi_n(\omega) \rightarrow \xi(\omega)$, 所以 $\lim_{n \rightarrow \infty} A_n = \emptyset$. 由概率的连续性得

$$\lim_{n \rightarrow \infty} P(A_n) = P(\lim_{n \rightarrow \infty} A_n) = 0.$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

另一方面, 由于 $\xi_n(\omega) \rightarrow \xi(\omega)$, 所以 $\lim_{n \rightarrow \infty} A_n = \emptyset$. 由概率的连续性得

$$\lim_{n \rightarrow \infty} P(A_n) = P(\lim_{n \rightarrow \infty} A_n) = 0.$$

所以

$$\limsup_{n \rightarrow \infty} |E\xi_n - E\xi| \leq \limsup_{n \rightarrow \infty} E|\xi_n - \xi| \leq \epsilon.$$

由 $\epsilon > 0$ 的任意性得

$$\lim_{n \rightarrow \infty} |E\xi_n - E\xi| = \lim_{n \rightarrow \infty} E|\xi_n - \xi| = 0.$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

现在证明单调收敛定理. 设 $0 \leq \xi_n(\omega) \nearrow \xi(\omega)$. 对任意的 $M > 0$, 令 $\eta_n = \xi_n I\{|\xi_n| \leq M\}$, $\eta = \xi I\{|\xi| \leq M\}$. 则 $\eta_n \leq \xi_n$, $\eta \leq \xi$,

$$\eta_n(\omega) \rightarrow \eta(\omega), \quad \forall \omega$$

并且 $0 \leq \eta_n \leq M$. 由有界收敛定理和数学期望的单调性知,

$$\lim_{n \rightarrow \infty} E\xi_n \geq \lim_{n \rightarrow \infty} E\eta_n = E\eta = \int_0^M x dF_\xi(x).$$

令 $M \rightarrow \infty$ 得

$$\lim_{n \rightarrow \infty} E\xi_n \geq E\xi.$$

如果 $E\xi = \infty$, 则结论已经得证.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

现在证明单调收敛定理. 设 $0 \leq \xi_n(\omega) \nearrow \xi(\omega)$. 对任意的 $M > 0$, 令 $\eta_n = \xi_n I\{|\xi_n| \leq M\}$, $\eta = \xi I\{|\xi| \leq M\}$. 则 $\eta_n \leq \xi_n$, $\eta \leq \xi$,

$$\eta_n(\omega) \rightarrow \eta(\omega), \quad \forall \omega$$

并且 $0 \leq \eta_n \leq M$. 由有界收敛定理和数学期望的单调性知,

$$\lim_{n \rightarrow \infty} E\xi_n \geq \lim_{n \rightarrow \infty} E\eta_n = E\eta = \int_0^M x dF_\xi(x).$$

令 $M \rightarrow \infty$ 得

$$\lim_{n \rightarrow \infty} E\xi_n \geq E\xi.$$

如果 $E\xi = \infty$, 则结论已经得证. 如果 $E\xi < \infty$, 则由于 $\xi_n \leq \xi$, 由单调性得 $E\xi_n \leq E\xi$. 所以

$$\lim_{n \rightarrow \infty} E\xi_n = E\xi.$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

下设 $0 \leq \xi_n(\omega) \searrow 0$, $E\xi_n$ 存在, 这时

$$0 \leq \xi_1 - \xi_n \nearrow \xi_1.$$

所以

$$E(\xi_1 - \xi_n) \rightarrow E\xi_1.$$

所以

$$E\xi_n \rightarrow 0.$$

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

最后证明控制收敛定理. 记

$$\eta_n = \sup_{m \geq n} |\xi_m - \xi|.$$

则 $0 \leq \eta_n(\omega) \searrow 0$.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

最后证明控制收敛定理. 记

$$\eta_n = \sup_{m \geq n} |\xi_m - \xi|.$$

则 $0 \leq \eta_n(\omega) \searrow 0$. 另一方面, 由于 $0 \leq \eta_n \leq 2\eta$, 所以 $0 \leq E\eta_n \leq 2E\eta < \infty$.

3.1 Mathematical expectation

3.1.5 Basic properties of expectations

最后证明控制收敛定理. 记

$$\eta_n = \sup_{m \geq n} |\xi_m - \xi|.$$

则 $0 \leq \eta_n(\omega) \searrow 0$. 另一方面, 由于 $0 \leq \eta_n \leq 2\eta$, 所以 $0 \leq E\eta_n \leq 2E\eta < \infty$. 由单调收敛定理,

$$E\eta_n \rightarrow 0.$$

而

$$|E\xi_n - E\xi| \leq E|\xi_n - \xi| \leq E\eta_n.$$

因此

$$\lim_{n \rightarrow \infty} E\xi_n = E\xi.$$