

Project 4 : West Nile Virus Prediction

General Assembly DSI 14

July 8, 2020

Song Yuan, Qi Wen, Jin, Eng Seng

Problem Statement

The main goal here for our analytics effort is to put systems in place that reduce people's exposure to mosquitoes that carry WNV

- Gene Leynes, Data Scientist

Questions?

How to reduce WNV
incidences?

Where and how does
WNV appear?

How to make prevention
methods more effective?

Cost Benefit?

History of West Nile Virus In Chicago



WNV Appears

West Nile Virus Exist In United States



64 Deaths

Total of 884 Cases Reported



10 & 13 Deaths



12 Deaths

THE GOAL IS 0 DEATHS

TABLE OF CONTENTS

01 About the West Nile Virus Vectors

EDA are worth Thousand Words!

02 Modeling

Modeling Process and Scoring

03 Cost Benefit Analysis

Cost Analysis

04 Conclusion

Key Findings, Recommendations, Limitations & Further Research

OI

About West Nile Virus Vectors


A black and white photograph of a man in a suit sitting in a car. He is looking down at a laptop computer, which is open on his lap. His right hand is on the keyboard, and his left hand is holding a white coffee cup with a lid. He is wearing a dark suit jacket, a light-colored dress shirt, and a patterned tie. A silver watch is visible on his left wrist. The background shows the interior of a car with a window and some blurred lights.

**EDA's ARE
WORTH A
THOUSAND
WORDS**


Mosquito Species Samples Collected




Total Number of Samples Collected for Each Mosquito Species




Number of Mosquitoes in Samples with West-Nile Virus




EDA for WNV Positive Cases, Spray and Trap Mapping




Top 20 Traps With WNV Recorded




Does Temperature
Affect the
Existence of WNV
??




What About Cool Temperature?




**Summer Is the
SEASON OF
INFECTION !!!**



**When the Weather
precipitation is
low , the more
WNV IS !!!**



The higher
humidity the more
frequent WNV
detected!



02

Modeling



How We Built the Models...

- The model trained on data mosquito data from the city of Chicago year, 2007,2009,2011 and 2013
- Tested against data from 2008, 2010, 2012 and 2014
- The key output is score which indicating the risk that a specific point/area could test positive for WNV in particular time.
- With Variety models tested, we keep the features which will improve the predictive ability
- For each trap, the model will predict the probability the WNV would be exist

Modeling Scoring


MODELS (SMOTE)	BEST PARAMS	KAGGLE PRIVATE SCORE	KAGGLE PUBLIC SCORE
Logistic Regression (lbfgs)	C = 1, Penalty = l2, max_iter = 1000	0.5719	0.5022
Logistic Regression (liblinear)	C = 1, Penalty = l1, max_iter = 1000	0.5719	0.5022
Decision Tree	max_depth = 10, min_samples = 4, min_samples_split = 20	0.5505	0.5873
Random Forest	max_depth = 20, max_leaf_nodes = 20, min_samples_leaf = 10, n_estimators = 100	0.5765	0.5882
AdaBoost	learning_rate = 1.0, n_estimators = 100	0.5463	0.5481
XGBoost	eval_metric = 'auc', scale_pos_weight = 18, subsample = 0.5, eta = 0.2	0.68208	0.6974

Results

The most predictive feature

Features	Importance
Month	0.036464
Trap_T095	0.028762
Trap_T225	0.018460
CULEX TERRITANS	0.017817
DATE	0.017134
Trap_T094	0.016628
Trap_T158	0.015939
Trap_T080	0.015407
Trap_T063	0.015346
Trap_T102	0.015210

ROC Curve with AUC = 0.828



 **ROC CURVE
WITH AUC**

**Scoring :
0.828**


03

Cost Benefit Analysis

What are the real cost and benefits
of vector control?



The Confusion Matrix



906 Cases

Reported From Chicago Year 2005 to 2016

\$ 12,977,000

Estimated Medical Cost from Year 2005 to
2016

\$ 702,000

Out of 49 Cases in 2016, costing \$ 702k in medical
expenses and lost productivity



148,510 Acres

Chicago city Area

\$ 137,000 estimated

Estimated for worst case scenario of
spraying whole city

5 Times

Medical Expenses & Loss Productivity cost incurred 5
times more expensive than spraying in the city !

04

Conclusion

Key Findings

- Mosquito numbers follow the natural onset of the season, begin in May and end in October.
- Peak at August and decreasing at October (died / hibernation due to winter)
- Spray Operations by City of Chicago and Department of Public Health do not have noticeable effect.
- Spray operations missed so locations where having most cases
- Strong Correlation between temperature and precipitation with number of WNV positive !
- Most WNV positive samples found between 71F - 80F, zero precipitation.

Recommendations

- Better spraying regime that is informed by weather and trap data.
- Concentrate efforts to combat WNV prone areas which tend to fall into two categories outline. Inner suburbs fairly high income, 1940s - 1960s housing and Low income, young in City.
- More public awareness campaigns. Urging people to minimize exposure with necessary precautions steps.

- The dataset was extremely imbalanced and we had to generate synthetic samples to balance the dataset
- Spraying data was limited to only 2 years with no indications of dosage
- Due to computational power limit, we had only tuned limited combinations of hyperparamters
- Other mosquito control efforts have not been accounted for in our model determine the actual effectiveness of spraying
- For a more complete analysis on economical and costings, we should collect the data in terms of healthcare expenses and mosquito control efforts.

Limitations & Further Research

THANKS!

ANY QUESTIONS ???