



This work is licensed under the terms of the CC BY-NC-SA 4.0 International License (<https://creativecommons.org/licenses/by-nc-sa/4.0>). This license requires that reusers give credit to the creator. It allows reusers to distribute, remix, adapt, and build upon the material in any medium or format, for noncommercial purposes only. If others modify or adapt the material, they must license the modified material under identical terms. *All images except for 'by-nc-sa.png' in this manual are licensed under CC0.*

---

# ABBREVIATION TABLE

## A B

abs	absolute
add	addi(tion)(tive)
adj	adjoint
algo	algorithm
arb	arbitrary
assoc	associa(tive)(tivity)
asum	assum(e)(ption)
becs	because

## E

-ec	-ec(t)(tor)(tion)(tive)
elem	element(s)
ent	entr(y)(ies)
equa	equality
equiv	equivalen(t)(ce)
exa	example
exe	exercise
exis	exist(s)(ing)
existns	existence
expo	exponent
expr	expression

## L

liney	linear(ly)
linity	linearity
len	length
low-	lower-

## R

recurly	recursively
repeti	repetition(s)
repres	represent(s)(ation(s))
req	require(s)(d)/requiring
respectly	respectively
restr	restrict(ion)(ive)(ing)
rev	revers(e(s))(ed)(ing)

## C

closd	closed under
coeff	coefficient
combina	combination
commu	commut(es)(ing)(ativity)
cond	condition
corres	correspond(s)(ing)
conveni	convenience
convly	conversely
count-	counter-
ctradic	contradict(s)(ion)
ctrapos	constrapositive

## F G H

factoriz	factorizaion
fini	finite
finide	finite-dimensional
homo	homogeneity
hypo	hypothesis

## M N

max	maxi(mal(ity))(mum)
min	mini(mal(ity))(mum)
multi	multipl(e)(icati-on/ve)
non0	nonzero
nonC	nonconst
notat	notation(al)

## S

seq	sequence
simlr	similar(ly)
solus	solution
sp	space
stmt	statement
std	standard
supp	suppose
surj	surjectiv(e)(ity)
suth	such that

## D

Ddkd	Dedekind
def	definition
deg	degree
deri	derivative(s)
diff	differentia(l)(ting)(tion)
dim	dimension(al)
disti	distinct
distr	distributive propert(ies)(ty)
div	div(ide)(ision)

## I

id	identity
immed	immediately
induc	induct(ion)(ive)
infily	infinitely
inje	injectiv(e)(ity)
inv	inver(se)(tib-le/ility)
iso	isomorph(ism)(ic)

## O P Q

othws	otherwise
orthog	orthogonal
orthon	orthonormal
poly	polynomial
posi	positive
prod	product
quad	quadratic
quotient	quot

## T U V W X Y Z

uniq	unique
uniques	uniqueness
val	value
-wd	-ward
-ws	-wise
wрто	with respect to

**0.B** NOTE:  $C, D$  are Dedekind cuts. Numbers used here are always rational.

$$\begin{aligned} \bullet \tilde{0} &= \{a : a < 0\} = \mathbf{Q} \setminus \mathbf{Q}^* \Rightarrow -\tilde{0} = \{a : a < -b \leq 0\} = \tilde{0}. \\ D &= \{a : a < -b, b \notin D\} = \{-a : -a < -b \Leftrightarrow a > b, b \notin D\}. \end{aligned}$$

• TIPS: Prove  $\forall \varepsilon > 0, \exists b \notin D$  suth  $b - \varepsilon \in D$ .

SOLUS: Asum  $\exists \varepsilon > 0$  suth  $\nexists b \notin D, b - \varepsilon \in D \Leftrightarrow \forall b \notin D, b - \varepsilon \notin D$ .

Then  $(b - \varepsilon) - \dots - \varepsilon = b - n \cdot \varepsilon \notin D$  for any  $n \in \mathbf{N}^+$ .

For any  $d \in D, \exists n \in \mathbf{N}^+$  suth  $b - n \cdot \varepsilon < d \Rightarrow b - n \cdot \varepsilon \in D$ , ctradict. □

**1** Prove (a)  $D + \tilde{0} = D$ , (b)  $-D$  is Dedekind cut, and  $D + (-D) = \tilde{0}$ .

SOLUS: (a)  $\forall d \in D, \exists \varepsilon > 0, d + \varepsilon \in D \Rightarrow (d + \varepsilon) + (-\varepsilon) \in D + \tilde{0}$ .

(b) Asum  $x \in -D$  is the largest elem of  $-D \Rightarrow \exists b \notin D, x < -b \Rightarrow 0 < -b - x$ .

Let  $\delta = (-b - x)/2 \Rightarrow 0 < \delta < -b - x \Rightarrow x < x + \delta < -b$ .

Thus by def,  $x + \delta \in -D$ , ctradict the max of  $x \in -D$ . Hence  $-D$  is Ddkd cut.

$D + (-D) = \{x + y : x + y < x - b, x \in D, b \notin D\}$ .

Supp  $a \in \tilde{0} \Rightarrow -a > 0$ . By TIPS,  $\exists b \notin D$  suth  $b + a \in D$ .

Note that  $b < b - a \notin D \Rightarrow -b > -b + a \in -D$ . Then  $(-b + a) + (b + a) = 2a < 0$ .

Thus  $\forall a \in \tilde{0}, \exists b \notin D, d = b + \frac{1}{2}a \in D, c = -b + \frac{1}{2}a \in -D \Rightarrow c + d = a \in D + (-D)$ . □

**5** Prove  $D$  is posi  $\Leftrightarrow -D$  is not posi  $\Leftrightarrow 0 \in D$ .

SOLUS:  $0 \notin -D = \{a : a < -b, b \notin D\} \Leftrightarrow \nexists b \notin D, b < 0 \Leftrightarrow \mathbf{Q} \setminus \mathbf{Q}^+ \not\subseteq D \Leftrightarrow 0 \in D$ . □

• Define  $D^+ = \{d \in D : d > 0\} = D \cap \mathbf{Q}^+$ . Then  $D^+ = \emptyset \Leftrightarrow D \not\subseteq \mathbf{Q} \setminus \mathbf{Q}^+ \Leftrightarrow 0 \notin D \Leftrightarrow D$  not posi.

Define  $D^- = \{r \notin D : r \leq 0\} = (\mathbf{Q} \setminus D) \cap (\mathbf{Q} \setminus \mathbf{Q}^+) = \mathbf{Q} \setminus (D \cup \mathbf{Q}^+)$ .

Then  $D^- = \emptyset \Leftrightarrow D \cup \mathbf{Q}^+ = \mathbf{Q} \Leftrightarrow \mathbf{Q} \setminus \mathbf{Q}^+ \subseteq D \Leftrightarrow 0 \in D \Leftrightarrow D$  posi. And  $\tilde{0}^- = \{0\}$ .

•  $(-D)^+ = (-D) \cap \mathbf{Q}^+ = \{a : 0 < a < -b, b \notin D \Leftrightarrow b \in D^- \setminus \{0\}\}$ .

$(-D)^- = (\mathbf{Q} \setminus -D) \cap (\mathbf{Q} \setminus \mathbf{Q}^+) = \{a : 0 \geq a \geq -b, \forall b \notin D\}$ .

Define  $\tilde{1} = \{a : a < 1\} \Rightarrow \tilde{1}^+ = \{a : 0 < a < 1\}, \tilde{1}^- = \{a : a \leq 0\} = \tilde{0} \cup \{0\}$ .

Then  $-\tilde{1} = \{a : -a > 1 \Leftrightarrow a < -1\}$  not posi.

• For  $C, D$  posi, define  $CD = \{a : a \leq cd, c \in C^+, d \in D^+\} = \{cd : c \in C^+, d \in D^+\} \cup (\mathbf{Q} \setminus \mathbf{Q}^+)$ .

Note that  $\{cd : c \in C^+, d \in D^+\} = CD \cap \mathbf{Q}^+ = (CD)^+$ .

• For  $C, D$  not posi, define  $CD = (-C)(-D) = \{cd : c \in (-C)^+, d \in (-D)^+\} \cup (\mathbf{Q} \setminus \mathbf{Q}^+)$ .

$CD = \{0 < cd < (-r)(-s) : r \in C^- \setminus \{0\}, s \in D^- \setminus \{0\}\} \cup (\mathbf{Q} \setminus \mathbf{Q}^+) = \{a : a < rs, r \in C^-, s \in D^-\}$ .

• The intuitive key point is that the prod of cuts is the cut with the endpoint being the prod of endpoints of cuts.

• For  $C$  not posi and  $D$  posi, define  $CD = \{cs : c \in C, s \in D^+\}$ .

• For a posi  $D, \tilde{1}D = \{a : a \leq ij < j, 0 < i < 1, j \in D^+\} \subseteq D$ .

Now  $(\tilde{1}D)^+ \subseteq D^+$ . 又  $\forall d \in D^+, \exists \varepsilon > 0, d + \varepsilon \in D^+ \Rightarrow d = (d + \varepsilon) \frac{d}{d + \varepsilon} \in \tilde{1}D$ .

•

---

**3** Show  $C \subsetneq D \iff D - C = \{d - y : d \in D, y > x, x \notin C\}$  posi.

**SOLUS:** (a) Supp  $C \subsetneq D$ . Let  $d \in D \setminus C \subseteq \mathbf{Q} \setminus C$ . For any  $a \in \tilde{0}$ , let  $y = d - a > d \Rightarrow a = d - y \in D - C$ .

(b) Supp  $D - C$  posi. Becs  $d = y$  for some  $d \in D, y > x \notin C$ .

Note that  $x \notin C \Rightarrow \forall c \in C, c < x < y = d \Rightarrow C \subseteq D$ . □

---

**ENDED**

## 0.C

**5** Supp  $a_1, a_2, \dots$  is a seq in  $\mathbf{Q}$ , and  $\sup\{a_1, a_2, \dots\} = \sqrt{2}$ .

Prove  $\sup\{a_n, a_{n+1}, \dots\} = \sqrt{2}$  for all  $n \in \mathbf{N}^+$ .

**SOLUS:** Becs the sup not in seq  $\Rightarrow$  infily many disti elem.

$\forall a_i, \exists a_j, a_i < a_j < \sqrt{2}$ . For  $a_{n+k}$ , choose  $a_i > a_{n+k}$ . If  $i \in \{1, \dots, n\}$ , then choose  $a_j > a_i$ .

After at most  $(n+1)$  steps, we have  $a_m$  with  $m > n$ . Thus  $\forall a_{n+i}, \exists a_{n+j}, a_{n+i} < a_{n+j} < \sqrt{2}$ .  $\square$

• Supp nonempty  $A \subseteq \mathbf{R}$ .

• **TIPS 1:** Define  $-A = \{-a : a \in A\} \Rightarrow -(-A) = A$ . Prove  $\sup(-A) = -\inf A$ .

**SOLUS:**  $-b$  is an upper bound of  $-A \iff \forall a \in A, -a \leq -b \iff a \geq b \iff b$  is a lower bound of  $A$ .

Thus  $-b_M = \sup(-A) \iff -b_M \leq -b \iff b_M \geq b \iff b_M = \inf A$ .  $\square$

• **TIPS 2:** Show if  $x \in \mathbf{R}$ , (a)  $\sup A > x \Rightarrow \exists a \in A, a > x$ , (b)  $\inf A < x \Rightarrow \exists a \in A, a < x$ .

**SOLUS:** (a)  $\nexists a > x \iff \forall a \in A, a \leq x$ . Then by def of sup.

OR. By (b),  $\inf(-A) = -\sup A < -x \Rightarrow \exists -a \in A, -a < -x$ . Simlr for (b).  $\square$

**6** Supp  $A, B \subseteq \mathbf{R}$  has infily many disti elem, so has  $A + B = \{a + b : a \in A, b \in B\}$ .

Prove  $\sup(A + B) = \sup A + \sup B$ , and  $\inf(A + B) = \inf A + \inf B$ .

**SOLUS:**  $\inf A + \inf B \leq a + b \leq \sup A + \sup B \Rightarrow \sup(A + B) \leq \sup A + \sup B, \inf A + \inf B \leq \inf(A + B)$ .

$\sup A + \sup B > \sup(A + B) \iff \sup A > \sup(A + B) - \sup B$

$\iff \exists a + \sup B > \sup(A + B) \iff \sup B > \sup(A + B) - a \iff \exists a + b > \sup(A + B)$ . Ctradic.

Simlr for  $\inf(A + B) \in A + B$ . OR. Apply to  $-A - B$ , becs  $\sup(-A) = -\inf A$ .  $\square$

**16** Supp  $\mathbf{R}_1, \mathbf{R}_2$  are complete ordered fields. Define  $\varphi_1, \varphi_2$  as in [0.11].

Define  $\mathcal{R}_1(a) = \{q \in \mathbf{Q} : \varphi_1(q) \leq a\} \Rightarrow \sup_{\mathcal{R}_1(a)} \varphi_1 = a$ . Define  $\psi_1(a) = \sup_{\mathcal{R}_1(a)} \varphi_2$ .

(a) Show  $\psi = \psi_1 : \mathbf{R}_1 \rightarrow \mathbf{R}_2$  is one-to-one. (b) Show  $\psi(0) = 0, \psi(1) = 1$ .

(c) Show  $\psi(a \pm b) = \psi(a) \pm \psi(b)$ , and  $\psi(ab^{-1}) = \psi(a)\psi(b)^{-1}$ .

(d) Supp  $a \in \mathbf{R}_1$ . Show  $a > 0 \iff \psi(a) > 0$ .

**SOLUS:** (a) Define  $\mathcal{R}_2(c) = \{q \in \mathbf{Q} : \varphi_2(q) \leq c\} \Rightarrow \sup_{\mathcal{R}_2(c)} \varphi_2 = c$ .

Define  $\psi_2(c) = \sup_{\mathcal{R}_2(c)} \varphi_1$ . Then  $\psi_2 : \mathbf{R}_2 \rightarrow \mathbf{R}_1$  well-defined.

Note that  $\mathcal{R}_2(\psi_1(a)) = \{q \in \mathbf{Q} : \varphi_2(q) \leq \sup_{\mathcal{R}_1(a)} \varphi_2\} = \mathcal{R}_1(a)$ .

Now  $\psi_2(\psi_1(a)) = \sup_{\mathcal{R}_1(a)} \varphi_1 = a$ . Rev the roles of  $\mathbf{R}_1, \mathbf{R}_2$ .

(b) Note that  $\varphi(q) < 0 \iff q < 0$ , and  $\varphi(0) = 0$ .

$\forall q \in \mathcal{R}_1(1) = \{1/m \in \mathbf{Q} : m \in \mathbf{N}^+, \varphi_1(1/m) = (1 + \dots + 1)^{-1} \leq 1\} \cup (\mathbf{Q} \setminus \mathbf{Q}^+), \varphi_2(q) \leq 1$ .

(c)  $\mathcal{R}_1(a \pm b) = \{p \pm q \in \mathbf{Q} : \varphi_1(p) \pm \varphi_1(q) \leq a \pm b\}$ .

$\mathcal{R}_1(ab^{-1}) = \{pq^{-1} \in \mathbf{Q} : \varphi_1(q) \cdot \varphi_1(q)^{-1} \leq ab^{-1}\}$ .

(d)  $a > 0 \iff \exists n \in \mathbf{N}^+, 1/n < a \iff \psi_1(a) > 0$ .  $\square$