



This work is licensed under the terms of the CC BY-NC-SA 4.0 International License (<https://creativecommons.org/licenses/by-nc-sa/4.0>). This license requires that reusers give credit to the creator. It allows reusers to distribute, remix, adapt, and build upon the material in any medium or format, for noncommercial purposes only. If others modify or adapt the material, they must license the modified material under identical terms. *All images except for 'by-nc-sa.png' in this manual are licensed under CC0.*

这是我个人挑战「*Measure, Integration & Real Analysis, by Sheldon Axler*」的学习笔记，包括课文补注和部分习题。我先从 *Supplement* 即第 0 章开始。我当时并没有学过数学分析，以为学完 Axler 的这个 Supplement 就能具备所有必要的知识基础。

0.B 节本来不太要命，但超出课文的补助却让我折戟沉沙——的确，它们不需要硬性知识门槛，可以用集合和数理逻辑来推导 \mathcal{D} 的一切。或许是缺乏 Dedekind cut 的系统学习，我推导这一切时感到我在亲手缔造一个数学分支；我不是自傲的意思，只是说，**这非常艰难**。但我还是坚持下来了；在此过程中我肉眼可见我在数理逻辑上的提升。

ABBREVIATION TABLE

A B

| | |
|-------|-----------------------|
| abs | absolute |
| add | addi(tion)(tive) |
| adj | adjoint |
| algo | algorithm |
| arb | arbitrary |
| assoc | associa(tive)(tivity) |
| asum | assum(e)(ption) |
| becs | because |

E

| | |
|---------|-------------------------|
| -ec | -ec(t)(tor)(tion)(tive) |
| elem | element(s) |
| ent | entr(y)(ies) |
| equa | equality |
| equiv | equivalen(t)(ce) |
| exa | example |
| exe | exercise |
| exis | exist(s)(ing) |
| existns | existence |
| expo | exponent |
| expr | expression |

L

| | |
|--------|------------|
| liney | linear(ly) |
| linity | linearity |
| len | length |
| low- | lower- |

R

| | |
|-----------|-------------------------|
| recurly | recursively |
| repeti | repetition(s) |
| repres | represent(s)(ation(s)) |
| req | require(s)(d)/requiring |
| respectly | respectively |
| restr | restrict(ion)(ive)(ing) |
| rev | revers(e(s))(ed)(ing) |

C

| | |
|---------|--------------------------|
| closd | closed under |
| coeff | coefficient |
| combina | combination |
| commu | commut(es)(ing)(ativity) |
| cond | condition |
| corres | correspond(s)(ing) |
| conveni | convenience |
| convly | conversely |
| count- | counter- |
| ctradic | contradict(s)(ion) |
| ctrapos | constrapositive |

F G H

| | |
|----------|--------------------|
| factoriz | factorizaion |
| fini | finite |
| finide | finite-dimensional |
| homo | homogeneity |
| hypo | hypothesis |

M N

| | |
|-------|-------------------------|
| max | maxi(mal(ity))(mum) |
| min | mini(mal(ity))(mum) |
| multi | multipl(e)(icati-on/ve) |
| non0 | nonzero |
| nonC | nonconst |
| notat | notation(al) |

S

| | |
|-------|-------------------|
| seq | sequence |
| simlr | similar(ly) |
| solus | solution |
| sp | space |
| stmt | statement |
| std | standard |
| supp | suppose |
| surj | surjectiv(e)(ity) |
| suth | such that |

D

| | |
|-------|-------------------------------|
| Ddkd | Dedekind |
| def | definition |
| deg | degree |
| deri | derivative(s) |
| diff | differentia(l)(ting)(tion) |
| dim | dimension(al) |
| disti | distinct |
| distr | distributive propert(ies)(ty) |
| div | div(ide)(ision) |

I

| | |
|--------|-------------------------|
| id | identity |
| immed | immediately |
| induc | induct(ion)(ive) |
| infily | infinitely |
| inje | injectiv(e)(ity) |
| inv | inver(se)(tib-le/ility) |
| iso | isomorph(ism)(ic) |

O P Q

| | |
|----------|-------------|
| othws | otherwise |
| orthog | orthogonal |
| orthon | orthonormal |
| poly | polynomial |
| posi | positive |
| prod | product |
| quad | quadratic |
| quotient | quot |

T U V W X Y Z

| | |
|---------|-----------------|
| uniq | unique |
| uniques | uniqueness |
| val | value |
| -wd | -ward |
| -ws | -wise |
| wrto | with respect to |

0.B NOTE: C, D are Dedekind cuts. Numbers used here are always rational.

• Define $\tilde{q} = \{a : a < q\}$, and $-\tilde{q} = \widetilde{-q} = \{a : a < -q\}$.

Then $\tilde{0} = \{a : a < 0\} = \mathbf{Q} \setminus \mathbf{Q}^* \Rightarrow -\tilde{0} = \{a : a < -0 \leq 0\} = \tilde{0}$.

• Define $-D = \{a : a < -b, b \notin D\} = \{-a : -a < -b \Leftrightarrow a > b, b \notin D\}$.

$-(-D) = -\{a : a < -b, b \notin D\} = \{c : c < -a, a \geq -b, \forall b \notin D\} = \{c : c < b, \forall b \notin D\} = D$.

The last equa is becs (a) $d \notin D \Rightarrow \exists b \notin D, d \geq b$, and (b) $d \in D \Rightarrow$ if $\exists b \notin D$ suth $d \geq b$, then $b \in D$, ctradic.

• **TIPS:** Prove $\forall \varepsilon > 0, \exists b \notin D$ suth $b - \varepsilon \in D$.

SOLUS: Asum $\exists \varepsilon > 0$ suth $\nexists b \notin D, b - \varepsilon \in D \Leftrightarrow \forall b \notin D, b - \varepsilon \notin D$.

Then $(b - \varepsilon) - \dots - \varepsilon = b - n \cdot \varepsilon \notin D$ for any $n \in \mathbf{N}^+$.

Now $\forall d \in D, \exists n \in \mathbf{N}^+$ suth $b - n \cdot \varepsilon < d \Rightarrow b - n \cdot \varepsilon \in D$, ctradic. □

1 Prove (a) $D + \tilde{0} = D$, (b) $-D$ is Dedekind cut, and $D + (-D) = \tilde{0}$.

SOLUS: (a) $\forall d \in D, \exists \varepsilon > 0, d + \varepsilon \in D \Rightarrow (d + \varepsilon) + (-\varepsilon) \in D + \tilde{0}$.

(b) Asum $x \in -D$ is the largest elem of $-D \Rightarrow \exists b \notin D, x < -b \Rightarrow 0 < -b - x$.

Let $\delta = (-b - x)/2 \Rightarrow 0 < \delta < -b - x \Rightarrow x < x + \delta < -b$.

Thus by def, $x + \delta \in -D$, ctradic the max of $x \in -D$. Hence $-D$ is Ddkd cut.

$D + (-D) = \{x + y : x + y < x - b, x \in D, b \notin D\}$.

Supp $a \in \tilde{0} \Rightarrow -a > 0$. By TIPS, $\exists b \notin D$ suth $b + a \in D$.

Note that $b < b - a \notin D \Rightarrow -b > -b + a \in -D$. Then $(-b + a) + (b + a) = 2a < 0$.

Thus $\forall a \in \tilde{0}, \exists b \notin D, d = b + \frac{1}{2}a \in D, c = -b + \frac{1}{2}a \in -D \Rightarrow c + d = a \in D + (-D)$. □

3 Show $C \subsetneq D \Leftrightarrow D - C = \{d - y : d \in D, y > x, x \notin C\}$ posi.

SOLUS: (a) $C \subsetneq D \Rightarrow \exists x \in D \setminus C \Rightarrow \exists y \in D, y > x \Rightarrow \exists d \in D, d > y \Leftrightarrow 0 < d - y \in D - C$.

(b) $0 \in D - C \Rightarrow \exists y > x \notin C, y \in D \Rightarrow \forall c \in C, c < x < y \in D \Rightarrow C \subseteq D$. 又 $D - C \neq \tilde{0}$. □

5 Prove (a) D posi $\Rightarrow -D$ not posi, (b) non0 $-D$ not posi $\Rightarrow D$ posi.

SOLUS: (a) $0 \notin \{a : a < -b, b \notin D\} \Leftrightarrow \nexists b \notin D, 0 < -b \Leftrightarrow \forall b \notin D, b \geq 0 \Leftrightarrow 0 \in D$.

(b) Becs $\tilde{0}$ is the largest non posi cuts. Thus $-D \neq \tilde{0} \Rightarrow -D \subsetneq \tilde{0} \Rightarrow \tilde{0} - (-D) = D$ posi.

OR. $\exists a < 0, a \notin -D = \{a : -a > b, b \notin D\} \Leftrightarrow \nexists b \notin D, -a > b \Leftrightarrow \forall b \notin D, 0 < -a \leq b$. □

• Define $D^+ = \{d \in D : d > 0\} = D \cap \mathbf{Q}^+$. Then $D^+ \neq \emptyset \Leftrightarrow \mathbf{Q} \setminus \mathbf{Q}^+ \subsetneq D \Leftrightarrow 0 \in D \Leftrightarrow D$ posi.

Define $D^- = \{r \notin D : r \leq 0\} = (\mathbf{Q} \setminus D) \cap (\mathbf{Q} \setminus \mathbf{Q}^+) = \mathbf{Q} \setminus (D \cup \mathbf{Q}^+)$.

(a) $D^- = \{0\} \Leftrightarrow D = \tilde{0}$. Convly, $\{r \notin D : r \leq 0\} = \{0\} \Rightarrow \mathbf{Q} \setminus D = \mathbf{Q}^*$.

(b) $D^- = \emptyset \Leftrightarrow D \cup \mathbf{Q}^+ = \mathbf{Q} \Leftrightarrow \mathbf{Q} \setminus \mathbf{Q}^+ \subseteq D \Leftrightarrow 0 \in D \Leftrightarrow D$ posi. **CORO:** D not posi $\Leftrightarrow 0 \in D^-$.

(c) $(D^-)^- = \{r \in D : r \leq 0\} = \mathbf{Q} \setminus D^+$. **CORO:** D not posi $\Leftrightarrow (D^-)^- = D$.

• $(-D)^+ = (-D) \cap \mathbf{Q}^+ = \{a : 0 < a < -b, b \notin D \Leftrightarrow b \in D^- \setminus \{0\}\}$.

$(-D)^- = (\mathbf{Q} \setminus -D) \cap (\mathbf{Q} \setminus \mathbf{Q}^+) = \{a : 0 \geq a \geq -b, \forall b \notin D\}$.

• For C, D posi, define $CD = \{a : a \leq cd, c \in C^+, d \in D^+\} = \{cd : c \in C^+, d \in D^+\} \cup (\mathbf{Q} \setminus \mathbf{Q}^+)$.

$\{cd : c \in C^+, d \in D^+\} = CD \cap \mathbf{Q}^+ = (CD)^+$. Note that ' $a \leq cd$ ' here is equiv to ' $a < cd$ '.

- For $-C, -D$ posi, define $CD = (-C)(-D) = \{cd : c \in (-C)^+, d \in (-D)^+\} \cup (\mathbf{Q} \setminus \mathbf{Q}^+)$.
 $CD = \{0 < cd < (-r)(-s) : r \in C^- \setminus \{0\}, s \in D^- \setminus \{0\}\} \cup (\mathbf{Q} \setminus \mathbf{Q}^+) = \{a : a < rs, r \in C^-, s \in D^-\}$.
 If $C, -C$ not posi $\Rightarrow C = \tilde{0}$, then with the asum $\tilde{0}D = \tilde{0}$, it still holds. Simlr for $D, -D$ not posi.
 - The intuitive key point is that the prod of cuts is the cut with the endpoint being the prod of endpoints of cuts.
-

- For D posi, define $D^{-1} = \{a : a < 1/b, b \notin D\} \Rightarrow DD^{-1} = \{a : a \leq d/b < 1, b \notin D, d \in D^+\} = \tilde{1}$.
 The last equa holds becs $\forall a \in \tilde{1} \cap \mathbf{Q}^+, \exists d \in D^+$ suth $b = d/a \notin D \Rightarrow d/b = a \in DD^{-1}$.
 - For non0 D not posi, define $D^{-1} = \{a : a < 1/b, b \in D^-\} \Rightarrow (D^{-1})^- = \{s : 0 \geq s \geq 1/b, \forall b \in D^-\}$.
 Then $DD^{-1} = \{a : a < rs, r \in D^-, s \in (D^{-1})^-\} = \{a : a < rs, \exists r \in D^-, 0 \leq rs \leq r/b, \forall b \in D^-\}$.
 Asum $\exists a \in \tilde{1} \cap \mathbf{Q}^+, \forall r \in D^-, s \in (D^{-1})^-, a \geq rs \Rightarrow a \geq r/b \Leftrightarrow r/a \leq b, \forall b \in D^-$. Let $r = 0$.
-

- For C not posi and D posi, we expect that CD not posi. Consider C and $-D$ both not posi.
 $CD = -C(-D) = -\{a : a < rt, r \in C^-, t \in (-D)^-\} = \{-a : a > b, b \geq rt, \forall r \in C^-, t \in (-D)^-\}$
 $= \{-a : a > rt, \forall r \in C^-, 0 \geq t \geq -s, \forall s \notin D\} = \{a : a < ru, \forall r \in C^-, 0 \leq u \leq s, \forall s \notin D\}$.
 $(r \leq 0 < s, rs \leq ru = -rt \leq 0 \leq rt \leq -rs.)$
 - Note the ' $0 \leq u$ '. Becs $C^- \neq \emptyset \Rightarrow 0 \in C^-$. If it is to be exactly $CD = \{a : a < 0\}$, then $C^- = \{0\}$,
 for if not, $\exists u > 0$, and $\exists r \in C^- \setminus \{0\}$, such that $\exists a < ru < 0$. Hence ' $0 \leq u$ ' is actually ' $0 < u$ '.
 - ' $u \leq s$ ' cannot be abbreviated as in $\{-a : a > -rs, \forall s \notin D, r \in C^-\} = \{a : a < rs, \forall s \notin D, r \in C^-\}$.
 ' $u \leq s$ ' cannot be ' $u < s$ ', becs here $rs < ru \Rightarrow \exists a = rs$. Simlr for ' $a < ru$ ' to be ' $a \leq ru$ ' with ' $u < s$ '.
 - Note that $\{u : 0 < u \leq s, \forall s \notin D\} \supsetneq D^+$. We show " $0 < u \leq s, \forall s \notin D$ " is equiv to " $\forall u \in D^+$ ".
 $LHS = \{a : a < ru, \forall r \in C^-, 0 < u \leq s, \forall s \notin D\}, \{a : a < rd, \forall r \in C^-, \forall d \in D^+\} = RHS$.
 Immed, $LHS \subseteq RHS$. For $RHS \subseteq LHS$:
 (I) If we cannot fix " $\min \mathbf{Q} \setminus D$ " by a certain scalar s_M , then becs ' $u \leq s$ ' is actually ' $u < s$ ',
 $u \in \mathbf{Q}^+, u \notin D^+ \Leftrightarrow u \geq s, \exists s \notin D \Leftrightarrow u \notin \{u : 0 < u < s, \forall s \notin D\}$.
 (II) Othws, $s_M = \max D$. Now in LHS , " $a < rs_M, \forall r \in C^-$ ".
 Supp $0 < d_1 < \dots < d_n < \dots \in D^+ [d_n < s_M]$ and $0 > r_1 > \dots > r_m > \dots \in C^-$.
 Becs $r_{m+1}s_M < r_ms_M < r_md_{n+1} < r_md_n > r_{m+1}d_{n+1}$. Now $r_{m+1}s_M < r_ms_M \notin CD = RHS$.
 Asum $\exists t \in RHS \setminus LHS \Rightarrow \underline{r_ms_M < t} < r_md_n, \forall m, n \in \mathbf{N}^+$. Let $u = t/r_m \Rightarrow s_M > u > d_n$. Ctradic. \square
-

- Consider $-C$ and D both posi. Omitting $C = \tilde{0}$.
 $CD = -[(-C)D] = -\{a : a \leq cd, c \in (-C)^+, d \in D^+\} = \{-a : a > b, b > cd, \forall c \in (-C)^+, d \in D^+\}$
 $= \{a : a < -cd, \forall d \in D^+, \forall c \text{ suth } \underline{0 < c < -r}, \exists r \in C^-\} \quad (rd < -cd < 0)$
 $= \{a : a < td, \forall d \in D^+, \forall t \text{ suth } r < t < 0, \exists r \in C^-\}$
 $= \{a : a < rd, \forall d \in D^+, \forall r \in C^- \setminus \{0\}\}.$
-

- For C posi and D not posi, to make $CD = DC$, we define $CD = \{a : a < cs, \forall c \in C^+, s \in D^-\}$.
-

4 Supp B, C, D non0 Dedekind cuts. Show $(BC)D = B(CD), B(C + D) = BC + BD$.

SOLUS: We discuss in cases.

| \backslash | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
|--------------|---|---|---|---|---|---|---|---|
| B | + | + | + | + | - | - | - | - |
| C | + | + | - | - | - | + | - | + |
| D | + | - | + | - | - | - | + | + |

$$(1) ((BC)D)^+ = \{(bc)d : bc \in (BC)^+, d \in D^+\} = \{b(cd) : b \in B^+, cd \in (CD)^+\} = (B(CD))^+.$$

$$(4) (BC)D = \{a : a < rs, r \in (BC)^-, s \in D^-\}.$$

$$(B(C + D))^+ = \{bs : b \in B^+, s \in (C + D)^+\} = \{bc + bd : b \in B^+, -bc < bd, c \in C, d \in D\} = P.$$

$$\{bc : b \in B^+, c \in C^+\} + \{b'd : b' \in B^+, d \in D^+\} = \{bc + b'd : b, b' \in B^+, c \in C^+, d \in D^+\} = Q.$$

ENDED

0.C

5 Supp a_1, a_2, \dots is a seq in \mathbf{Q} , and $\sup\{a_1, a_2, \dots\} = \sqrt{2}$.

Prove $\sup\{a_n, a_{n+1}, \dots\} = \sqrt{2}$ for all $n \in \mathbf{N}^+$.

SOLUS: Becs the sup not in seq \Rightarrow infily many disti elem.

$\forall a_i, \exists a_j, a_i < a_j < \sqrt{2}$. For a_{n+k} , choose $a_i > a_{n+k}$. If $i \in \{1, \dots, n\}$, then choose $a_j > a_i$.

After at most $(n+1)$ steps, we have a_m with $m > n$. Thus $\forall a_{n+i}, \exists a_{n+j}, a_{n+i} < a_{n+j} < \sqrt{2}$. \square

• Supp nonempty $A \subseteq \mathbf{R}$.

• **TIPS 1:** Define $-A = \{-a : a \in A\} \Rightarrow -(-A) = A$. Prove $\sup(-A) = -\inf A$.

SOLUS: $-b$ is an upper bound of $-A \iff \forall a \in A, -a \leq -b \iff a \geq b \iff b$ is a lower bound of A .

Thus $-b_M = \sup(-A) \iff -b_M \leq -b \iff b_M \geq b \iff b_M = \inf A$. \square

• **TIPS 2:** Show if $x \in \mathbf{R}$, (a) $\sup A > x \Rightarrow \exists a \in A, a > x$, (b) $\inf A < x \Rightarrow \exists a \in A, a < x$.

SOLUS: (a) $\nexists a > x \iff \forall a \in A, a \leq x$. Then by def of sup.

OR. By (b), $\inf(-A) = -\sup A < -x \Rightarrow \exists -a \in A, -a < -x$. Simlr for (b). \square

6 Supp $A, B \subseteq \mathbf{R}$ has infily many disti elem, so has $A + B = \{a + b : a \in A, b \in B\}$.

Prove $\sup(A + B) = \sup A + \sup B$, and $\inf(A + B) = \inf A + \inf B$.

SOLUS: $\inf A + \inf B \leq a + b \leq \sup A + \sup B \Rightarrow \sup(A + B) \leq \sup A + \sup B, \inf A + \inf B \leq \inf(A + B)$.

$\sup A + \sup B > \sup(A + B) \iff \sup A > \sup(A + B) - \sup B$

$\iff \exists a + \sup B > \sup(A + B) \iff \sup B > \sup(A + B) - a \iff \exists a + b > \sup(A + B)$. Ctradic.

Simlr for $\inf(A + B) \in A + B$. OR. Apply to $-A - B$, becs $\sup(-A) = -\inf A$. \square

16 Supp $\mathbf{R}_1, \mathbf{R}_2$ are complete ordered fields. Define φ_1, φ_2 as in [0.11].

Define $\mathcal{R}_1(a) = \{q \in \mathbf{Q} : \varphi_1(q) \leq a\} \Rightarrow \sup_{\mathcal{R}_1(a)} \varphi_1 = a$. Define $\psi_1(a) = \sup_{\mathcal{R}_1(a)} \varphi_2$.

(a) Show $\psi = \psi_1 : \mathbf{R}_1 \rightarrow \mathbf{R}_2$ is one-to-one. (b) Show $\psi(0) = 0, \psi(1) = 1$.

(c) Show $\psi(a \pm b) = \psi(a) \pm \psi(b)$, and $\psi(ab^{-1}) = \psi(a)\psi(b)^{-1}$.

(d) Supp $a \in \mathbf{R}_1$. Show $a > 0 \iff \psi(a) > 0$.

SOLUS: (a) Define $\mathcal{R}_2(c) = \{q \in \mathbf{Q} : \varphi_2(q) \leq c\} \Rightarrow \sup_{\mathcal{R}_2(c)} \varphi_2 = c$.

Define $\psi_2(c) = \sup_{\mathcal{R}_2(c)} \varphi_1$. Then $\psi_2 : \mathbf{R}_2 \rightarrow \mathbf{R}_1$ well-defined.

Note that $\mathcal{R}_2(\psi_1(a)) = \{q \in \mathbf{Q} : \varphi_2(q) \leq \sup_{\mathcal{R}_1(a)} \varphi_2\} = \mathcal{R}_1(a)$.

Now $\psi_2(\psi_1(a)) = \sup_{\mathcal{R}_1(a)} \varphi_1 = a$. Rev the roles of $\mathbf{R}_1, \mathbf{R}_2$.

(b) Note that $\varphi(q) < 0 \iff q < 0$, and $\varphi(0) = 0$.

$\forall q \in \mathcal{R}_1(1) = \{1/m \in \mathbf{Q} : m \in \mathbf{N}^+, \varphi_1(1/m) = (1 + \dots + 1)^{-1} \leq 1\} \cup (\mathbf{Q} \setminus \mathbf{Q}^+), \varphi_2(q) \leq 1$.

(c) $\mathcal{R}_1(a \pm b) = \{p \pm q \in \mathbf{Q} : \varphi_1(p) \pm \varphi_1(q) \leq a \pm b\}$.

$\mathcal{R}_1(ab^{-1}) = \{pq^{-1} \in \mathbf{Q} : \varphi_1(q) \cdot \varphi_1(q)^{-1} \leq ab^{-1}\}$.

(d) $a > 0 \iff \exists n \in \mathbf{N}^+, 1/n < a \iff \psi_1(a) > 0$. \square