

8D latent | Command-conditioned orbit families

Frozen offline dynamics | 8 initial seeds | 100 burn-in cycles | 128 phase samples

One PCA basis per panel, fitted to all command values and all final seed curves.


— Actual representative seed — Other converged seeds ---- Cycle residual > 0.01

(a) Forward velocity (m/s)

Seed spread max $1.8e-07$ | Cycle residual max $3.72e-07$

Numerical groups 1-1 | Unresolved seeds 0

— -0.8 — 0 — 0.8 — 1.6
— -0.4 — 0.4 — 1.2 — 2




(b) Lateral velocity (m/s)

Seed spread max $1.79e-07$ | Cycle residual max $3.01e-07$

Numerical groups 1-1 | Unresolved seeds 0

— -1 — -0.33 — 0.33 — 1
— -0.67 — 0 — 0.67




(c) Yaw rate (rad/s)

Seed spread max $1.79e-07$ | Cycle residual max $3.1e-07$

Numerical groups 1-1 | Unresolved seeds 0

— -1 — -0.33 — 0.33 — 1
— -0.67 — 0 — 0.67




(d) Gait period (s)

Seed spread max $2.18e-07$ | Cycle residual max $4.84e-07$

Numerical groups 1-1 | Unresolved seeds 0

— 0.5 — 0.7 — 0.9 — 1
— 0.6 — 0.8




(e) Swing height (m)

Seed spread max $2e-07$ | Cycle residual max $3.38e-07$

Numerical groups 1-1 | Unresolved seeds 0

— 0.06 — 0.1 — 0.14 — 0.16
— 0.08 — 0.12




(f) Torso height (m)

Seed spread max $3.35e-07$ | Cycle residual max $6.62e-07$

Numerical groups 1-1 | Unresolved seeds 0

— 0.6 — 0.7 — 0.8 — 0.85
— 0.64 — 0.75



Dots: phase zero. Groups use complete-link RMS diameter ≤ 0.01 among cycle-resolved seeds; not a proof of multiple attractors.