

Convex Optimization Note

Garrick Lin

March 12, 2019

1 Affine Set

1.1 Definition

A set \mathcal{A} is called affine set when it satisfied:

If $\mathbf{x}_1 \in \mathcal{A}$ and $\mathbf{x}_2 \in \mathcal{A}$, then, $\forall \theta \in \mathcal{R}$, $x = \theta\mathbf{x}_1 + (1 - \theta)\mathbf{x}_2$ also belong to \mathcal{A} .

The expression $\theta\mathbf{x}_1 + (1 - \theta)\mathbf{x}_2$ represent a line that cross through \mathbf{x}_1 and \mathbf{x}_2 . Hence, affine set can be explained intuitively as:

Affine set is a set that contain the line which cross through any two point within this set.

1.2 Properties

Assume a affine set $\mathcal{A} \subseteq \mathcal{R}^n$. If \mathcal{A} contain original point($\mathbf{0}$), then, \mathcal{A} is a subspace. In order to prove this, \mathcal{A} must satisfied following three rules:

1. \mathcal{A} contain original point.
2. If $\mathbf{v} \in \mathcal{A}$, then, $\forall \theta \in \mathcal{R}$, $\theta\mathbf{v} \in \mathcal{A}$.
3. If $\mathbf{v}_1, \mathbf{v}_2 \in \mathcal{A}$, then, $\mathbf{v}_1 + \mathbf{v}_2 \in \mathcal{A}$.

The first rule is already satisfied. For the second rule, $\forall \theta \in \mathcal{R}$, $\theta\mathbf{v}$ satisfied:

$$\theta\mathbf{v} = \theta\mathbf{v} + (1 - \theta)\mathbf{0}$$

Since \mathcal{A} is affine and $\mathbf{v}, \mathbf{0} \in \mathcal{A}$, $\theta\mathbf{v} \in \mathcal{A}$ always true. For the third rule, there is:

$$\mathbf{v}_1 + \mathbf{v}_2 = 2(\frac{1}{2}\mathbf{v}_1 + \frac{1}{2}\mathbf{v}_2)$$

$\frac{1}{2}\mathbf{v}_1 + \frac{1}{2}\mathbf{v}_2$ is in \mathcal{A} as \mathcal{A} is affine and $\frac{1}{2}\mathbf{v}_1, \frac{1}{2}\mathbf{v}_2 \in \mathcal{A}$ (According to the second rule, $\mathbf{v}_1, \mathbf{v}_2 \in \mathcal{A}$, then, $\frac{1}{2}\mathbf{v}_1, \frac{1}{2}\mathbf{v}_2 \in \mathcal{A}$). According to the second rule, $2(\frac{1}{2}\mathbf{v}_1 + \frac{1}{2}\mathbf{v}_2) \in \mathcal{A}$

Therefore, any affine set \mathcal{A}' can be thought as a subspace \mathcal{V} with a transportation \mathbf{v} , which means, $\mathcal{A}' = \mathcal{V} + \mathbf{v}$. On the other hand, an affine set subtract some constant vector ($\mathcal{A}' - \mathbf{v}$), which make the new set contain original point, can form a subspace. Formally, for a affine set $\mathcal{A} = \{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n\}$ and $\mathbf{a}_i \in \mathcal{A}$, the set:

$$\mathcal{A} - \mathbf{a}_i = \{\mathbf{a}_1 - \mathbf{a}_i, \mathbf{a}_2 - \mathbf{a}_i, \dots, \mathbf{a}_{i-1} - \mathbf{a}_i, \mathbf{0}, \mathbf{a}_{i+1} - \mathbf{a}_i, \dots, \mathbf{a}_n - \mathbf{a}_i\}$$

form a subspace.