

Math 440 – Real Analysis II

Homework 6

Amandeep Gill

March 27, 2015

Problem 36 Let X be a vector space over \mathbb{R} with inner product $\langle \cdot, \cdot \rangle$, where $\|\cdot\|$ is defined for all $\vec{x} \in X$ as $\|\vec{x}\| = \sqrt{\langle \vec{x}, \vec{x} \rangle}$. Then for all $\vec{x}, \vec{y} \in X$, $\|\vec{x} + \vec{y}\|^2 = \|\vec{x}\|^2 + \|\vec{y}\|^2$ if and only if $\langle \vec{x}, \vec{y} \rangle = 0$.

Proof: For all $\vec{x}, \vec{y} \in X$, using the definition of $\|\cdot\|$ on X ,

$$\begin{aligned} \|\vec{x} + \vec{y}\|^2 &= \langle \vec{x} + \vec{y}, \vec{x} + \vec{y} \rangle \\ &= \langle \vec{x}, \vec{x} + \vec{y} \rangle + \langle \vec{y}, \vec{x} + \vec{y} \rangle \\ &= \langle \vec{x}, \vec{x} \rangle + \langle \vec{x}, \vec{y} \rangle + \langle \vec{y}, \vec{x} \rangle + \langle \vec{y}, \vec{y} \rangle \\ &= \|\vec{x}\|^2 + 2\langle \vec{x}, \vec{y} \rangle + \|\vec{y}\|^2 \end{aligned}$$

As this is a chain of strict equality, and since the inner product is a non-negative function, $\|\vec{x} + \vec{y}\|^2 = \|\vec{x}\|^2 + \|\vec{y}\|^2$ only if $\langle \vec{x}, \vec{y} \rangle = 0$. ■

Problem 37 Let X be a vector space over \mathbb{R} with inner product $\langle \cdot, \cdot \rangle$, and define $\|\cdot\|$ as $\|\vec{x}\| = \sqrt{\langle \vec{x}, \vec{x} \rangle}$. $\|\cdot\|$ is a norm on X .

(a) $\|\vec{x}\| \geq 0$ for all $\vec{x} \in X$

Proof: This follows from the fact that $0 \leq \langle \vec{x}, \vec{x} \rangle$ for all $\vec{x} \in X$, and that the square root function has no negative outputs on the non-negative reals. ■

(b) $\|\vec{x}\| = 0$ if and only if $\vec{x} = \vec{0}$

Proof: This follows from the fact that $\langle \vec{x}, \vec{x} \rangle = 0$ if and only if $\vec{x} = \vec{0}$, and that the square root function has only one zero, which is at 0. ■

(c) $\|c \cdot \vec{x}\| = |c| \cdot \|\vec{x}\|$ for all $c \in \mathbb{R}$ and all $\vec{x} \in X$

Proof: Let $c \in \mathbb{R}$ and $\vec{x} \in X$, then $\|c \cdot \vec{x}\| = \sqrt{\langle c\vec{x}, c\vec{x} \rangle} = \sqrt{c^2 \langle \vec{x}, \vec{x} \rangle}$. This is acceptable because it is a double application of property (d) of Orthogonal Functions twice. Since $\sqrt{c^2} = |c|$, $\|c \cdot \vec{x}\| = |c| \cdot \|\vec{x}\|$. ■

Problem 37 continued

(d) $\|\vec{x} + \vec{y}\| \leq \|\vec{x}\| + \|\vec{y}\|$ for all $\vec{x}, \vec{y} \in X$

Proof: Let $\vec{x}, \vec{y} \in X$. By previous work, $\|\vec{x} + \vec{y}\|^2 = \|\vec{x}\|^2 + 2\langle \vec{x}, \vec{y} \rangle + \|\vec{y}\|^2$. Because $\langle \vec{x}, \vec{y} \rangle \leq \|\vec{x}\| \|\vec{y}\|$ by the Cauchy-Schwartz Inequality, so this gives $\|\vec{x} + \vec{y}\|^2 \leq \|\vec{x}\|^2 + 2\|\vec{x}\| \|\vec{y}\| + \|\vec{y}\|^2 = (\|\vec{x}\| + \|\vec{y}\|)^2$. Thus, for all $\vec{x}, \vec{y} \in X$, $\|\vec{x} + \vec{y}\| \leq \|\vec{x}\| + \|\vec{y}\|$. ■

Problem 38 Let $f(x) = \sin(\pi x)$, $\phi_1(x) = 1$, and $\phi_2(x) = x$. Find c_1 and c_2 such that $S_2(x) = c_1\phi_1(x) + c_2\phi_2(x)$ gives the best approximation of f on $[-1, 1]$.

Solution: Need to choose c_1 and c_2 such that

$$\int_{-1}^1 \sin^2(\pi x) dx - c_1^2 \left(\int_{-1}^1 f_1^2(x) \right) dx - c_2^2 \left(\int_{-1}^1 f_2^2(x) \right) dx \text{ is minimized.}$$

$$\text{Since } \int_{-1}^1 \sin^2(\pi x) dx = 1$$

Problem 40 If f and f_n for $n \in \mathbb{N}$ are Riemann integrable functions on $[a, b]$, and $\{f_n\}$ converges uniformly to f on $[a, b]$ then $\{f_n\}$ converges in the mean to f on $[a, b]$.

Proof: Let f and f_n for $n \in \mathbb{N}$ be Riemann integrable functions on $[a, b]$, where $\{f_n\}$ converges uniformly to f on $[a, b]$, then there exists a number n_o such that for all $\epsilon > 0$ and $n \in \mathbb{N}$, if $n > n_o$ then $|f - f_n| < \frac{\sqrt{\epsilon}}{\sqrt{b-a}}$.

From this, we get $\int_a^b (f - f_n)^2 dx < \int_a^b \left(\frac{\sqrt{\epsilon}}{\sqrt{b-a}} \right)^2 dx = \frac{\epsilon x}{b-a} \Big|_a^b = \epsilon$. Therefore,

$$\lim_{n \rightarrow \infty} \int_a^b (f - f_n)^2 dx = 0 \text{ and } \{f_n\} \text{ converges in the mean to } f \text{ on } [a, b].$$
■