

A deep-learning algorithm using real-time collected intraoperative vital sign signals for predicting acute kidney injury after major non-cardiac surgeries: A modelling study

Soomin Chung¹, Sehoon Park², Soie Kwon³, Jeong Min Cho², Jiwon Ryu⁴, Sejoong Kim⁴, Jeonghwan Lee⁵, Kwangsoo Kim⁶, Hajeong Lee¹



1 Department of Interdisciplinary Program in Bioengineering, Seoul National University, Korea, Republic of
2 Department of Internal Medicine-Nephrology, Seoul National University Hospital, Korea, Republic of
3 Department of Internal Medicine-Nephrology, Chung-Ang University Hospital, Korea, Republic of
4 Department of Internal Medicine-Nephrology, Seoul National University Bundang Hospital, Korea, Republic of
5 Department of Internal Medicine-Nephrology, Seoul National University College of Medicine, Korea, Republic of
6 Department of Transdisciplinary Department of Medicine & Advanced Technology, Seoul National University Hospital, Korea, Republic of


INTRODUCTION

- PO-AKI affects 5-7% of non-cardiac surgeries with high morbidity/mortality [1,2]
- Current models use only preoperative data (demographics, labs)**, missing critical intraoperative hemodynamics [3,4]
- Kidney susceptibility to BP/HR fluctuations makes **real-time monitoring essential for AKI prediction** [5]
- Deep learning enables analysis of complex minute-scale vital sign patterns [6]

OBJECTIVES

- Develop a **deep-learning algorithm** using **intraoperative vital sign signals** (DL-IVSS) to predict PO-AKI
- Compare **DL-IVSS** with traditional preoperative-only models
- Evaluate **Ensemble approach** combining clinical variables with DL-IVSS
- External validation** across multiple hospital cohorts

METHODS



CONCLUSION

- First study** to leverage **minute-scale intraoperative vital signs** (~9.3M data points) for PO-AKI prediction
- Ensemble model achieved superior performance** (AUROC 0.795) vs. conventional SPARK model (0.724)
- Intraoperative hemodynamics outperformed preoperative factors** in feature importance analysis
- Clinical utility**: Model maintains **>95% sensitivity and >95% specificity** at preset thresholds across validation cohorts
- Robust external validation**: Consistent performance across 3 independent hospital datasets

REFERENCES

- Grams ME, et al. Acute kidney injury after major surgery: a retrospective analysis of veterans health administration data. *Am J Kidney Dis.* 2016;67(6):872-80.
- Biteker M, et al. Incidence, risk factors, and outcomes of perioperative acute kidney injury in noncardiac and nonvascular surgery. *Am J Surg.* 2014;207(1):53-9.
- Bell S, et al. Risk of postoperative acute kidney injury in patients undergoing orthopaedic surgery. *BMJ.* 2015;351:h5639.
- Park S, et al. Simple postoperative AKI risk (SPARK) classification before noncardiac surgery: a prediction index development study with external validation. *J Am Soc Nephrol.* 2019;30(1):170-81.
- Park S, et al. Intraoperative arterial pressure variability and postoperative acute kidney injury. *Clin J Am Soc Nephrol.* 2020;15(1):35-46.
- Adhikari L, et al. Improved predictive models for acute kidney injury with IDEA: intraoperative data embedded analytics. *PLoS One.* 2019;14(4):e0214904.
- Walsh M, et al. Relationship between intraoperative mean arterial pressure and clinical outcomes after noncardiac surgery. *Anesthesiology.* 2013;119(3):507-15.
- Prowle JR, et al. Postoperative acute kidney injury in adult non-cardiac surgery: joint consensus report of the acute disease quality initiative and perioperative quality initiative. *Nat Rev Nephrol.* 2021;17(9):605-18.


RESULTS

Performance Comparison


Hospital (Cases N (%))	Method	AUROC	Balanced Accuracy
Developmental cohort (3,188 (6.2%))	SPARK	0.724	0.664
	DL-IVSS_only	0.707	0.666
	DL-IVSS_PCFs 11	0.765	0.708
	Ensemble_PCFs 11	0.795	0.732
EVC 1 (2,519 (5.3%))	SPARK	0.697	0.646
	DL-IVSS_only	0.637	0.601
	DL-IVSS_PCFs 11	0.716	0.655
	Ensemble_PCFs 11	0.762	0.696
EVC 2 (579 (4.7%))	SPARK	0.745	0.689
	DL-IVSS_only	0.607	0.588
	DL-IVSS_PCFs 11	0.761	0.701
	Ensemble_PCFs 11	0.786	0.715

DL-IVSS_only; only use Intra operative signals
DL-IVSS_PCFs 11; add Preoperative Clinical Features 11 on DL-IVSS_only
Ensemble_PCFs 11; CatBoost + DL-IVSS_PCFs 11


Feature importance



Calibration plot



Clinical threshold



CONTACT

Soomin Chung
soomin.chung.9910@gmail.com

Hajeong Lee
mdhjlee@gmail.com