



A**P = 0.779***Mann–Whitney U, two-sided*


Full v2 cohort (good n=18 vs bad n=17). Each dot = one pre-CRT tumor. Box = Q1/median/Q3, whiskers = 1.5×IQR. Black diamond = median.

B**P = 0.961***Mann–Whitney U, two-sided**IC50 < 500 nM across all HLA–peptide predictions (v2 pvacseq re-run, MHCflurry).*


C

P = 0.609


Mann–Whitney U, two-sided



$$PCN = \sum (2 \cdot VAF \times \text{binder count}) \text{ across mutation sites.}$$

D**P = 0.429***good vs bad (Mann-Whitney, two-sided)**within-good Wilcoxon: P = 0.125**within-bad Wilcoxon: P = 0.062*

N = 11 paired subjects. Lines go from pre-baseline (0) to post- Δ . Black diamond = group median.

E**P = 0.247***good vs bad (Mann-Whitney, two-sided)**within-good Wilcoxon: P = 0.125**within-bad Wilcoxon: P = 0.031**Mutation sites with ≥ 1 binder after CRT minus those before CRT.*


F

P = 0.537

good vs bad (Mann-Whitney, two-sided)

within-good Wilcoxon: P = 0.125


within-bad Wilcoxon: P = 0.031



PCN integrates allele-frequency-weighted binder count across sites.


G

• median Δ with 95% BCa CI (bootstrap $n=2000$)



Zero-crossing CI indicates non-significance. Normalization: each metric scaled to its own max $|CI \text{ bound}|$; raw values shown on right.

H



N = 11 paired subjects (subject 13 excluded per manuscript tumor-only rule). Negative bars = post-CRT neoantigen clearance.