

My Lower-Risk Plan

Choose the parts that fit your actual pattern. This is not a promise to be perfect.

The substance or situation I most want to make safer:

What raises my risk?

- | | |
|--|--|
| ☐ Using alone | ☐ Unknown or changing supply |
| ☐ Opioids may be involved | ☐ Mixing substances |
| ☐ Alcohol with other drugs | ☐ Sedating medications |
| ☐ Recent period of not using | ☐ Using more than intended |
| ☐ Strong cravings or withdrawal | ☐ Housing or privacy problems |
| ☐ Using when very distressed | ☐ Pain or medical problems |
| ☐ I am not sure what is in the drug | ☐ Other |

The one risk I most want to lower:

Safety steps I am willing to use

- | | |
|---|---|
| ☐ Carry naloxone | ☐ Tell someone where naloxone is |
| ☐ Avoid using alone | ☐ Have someone check on me |
| ☐ Use drug checking / fentanyl test strips | ☐ Avoid mixing sedating substances |
| ☐ Use less when supply or tolerance is uncertain | ☐ Keep emergency information visible |
| ☐ Use sterile equipment / safer supplies | ☐ Get wound or medical care earlier |
| ☐ Talk with a prescriber or pharmacist | ☐ Ask about medication treatment |

Plan for High-Risk Moments

A situation where my risk usually goes up:

A person or service I can contact before or during that situation:

If I have been abstinent or using less for a while, how will I account for lower tolerance?

Where I will keep naloxone or other emergency supplies:

What I want people around me to do if I become unresponsive or stop breathing normally:

Change goal, if I want one

- ☐ Make one part safer
- ☐ Use less often
- ☐ Use a smaller amount
- ☐ Avoid one high-risk combination
- ☐ Try medication treatment
- ☐ Try counseling or peer support
- ☐ Take a break from use
- ☐ Work toward abstinence
- ☐ I am not choosing a change goal yet

My next step:

Medical caution. If you may be physically dependent on alcohol or benzodiazepines, do not assume abrupt stopping at home is safe. Seek medical guidance about withdrawal risk.