



Rainfall and Water Quality

Benoit Roig, Estelle Baurès, Aude Valerie Jung, Ianis Delpla, Olivier Thomas

► To cite this version:

Benoit Roig, Estelle Baurès, Aude Valerie Jung, Ianis Delpla, Olivier Thomas. Rainfall and Water Quality. Olga E. Martín and Tricia M. Roberts Eds. Rainfall: Behavior, Forecasting and Distribution, Nova Science Publisher, Inc, pp.91-104, 2012, 978-1620815519. hal-01664965

HAL Id: hal-01664965

<https://hal.science/hal-01664965>

Submitted on 22 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RAINFALL AND WATER QUALITY

Benoit Roig^{1,2,*}, Estelle Baures^{1,2}, Aude Valérie Jung³, Ianis Delpla^{1,2} and Olivier Thomas^{1,2}

¹Advanced School of Public Health (EHESP), LERES, Avenue du Professeur Léon Bernard - CS 74312, Rennes Cedex, France

²INSERM U1085, LERES, Rennes Cedex, France

³School of Environmental Engineering (EME), Campus de Ker Lann, Avenue Robert Schumann, Bruz, France

1. INTRODUCTION

The consideration of rainfalls impact on water management is not new and started with the history of sanitation. At the beginning of the 20th century sewer networks in urban area were designed both for sewage and for stormwater management. In the 70's the impact of urban runoff on water quality was already studied (Pitt et al. 1977) and more recently the question of climate change impacts on water quality was considered with the effect of heavier rainy events. Commonly, categories of rain are defined by their intensity: heavy, moderate and light rain being characterised by the amount of mm of water/hour (respectively 2.5, 2.5 to 7.5, >7.5). According to the duration of a given rainfall, the same intensity can be observed during a period (shower) or a longer one (storm). Consequences of these precipitations can result in so called extreme rain or flood respectively at local or regional scale. As reported by the 2007 IPCC report, it is likely that the frequency of heavy precipitation events has increased over most areas, particularly in Europe and North America, and that available research suggests a significant future increase in heavy rainfall events in many regions. Such phenomena, increasing in frequency and intensity, may have a strong impact on the quality of the waters. Indeed, resulting changes in flow regimes will influence the chemistry, hydromorphology and ecology of water bodies at the environmental level as well as, in urban area, the wastewater quality, the sewage treatment efficiency and the corresponding pollutants load. The effect of rainy events can also reach drinking water production system, impairing the quality of the distributed water (Figure 1)

This chapter aims to make an overview of the impact of rainfall on water quality, by considering, within the water life cycle, chemical, biological and physico-chemical parameters of water quality.




Figure 1. Impact of heavy events on water life cycle.

* E-mail: benoit.roig@ehesp.fr.

2. WATER QUALITY

Water bodies are characterised regarding their chemical and biological status. A number of parameters are used to analyse water quality. European water regulations (Drinking-water Directive (DWD) and the Water Framework Directive (WFD)) have been established, since the end of the 1990s, to ensure public health security for water intended for human consumption by promoting a policy for the management and protection of water resources at European level. More recently, a new preventive risk management approach was initiated by WHO and water suppliers, via the implementation of Water Safety Plans (Bartram et al., 2009), the aim of which is to guarantee the safety of a drinking-water supply through the use of a comprehensive risk assessment and risk management approach. Despite this regulatory framework, numerous water bodies are often of impaired quality and non-compliant with European regulations (Bloech, 2011).

In general, raw water quality is influenced by both 'natural' and anthropogenic driving forces. The former include spatial and temporal variation in water availability, wildlife, topography, geology, vegetation and the future impact of climate change, particularly with respect to the frequency and severity of droughts or rainfalls. Anthropogenic driving forces come from each of the key sectors that use water (industry, energy production, public water supply and agriculture) and include point sources (e.g. wastewater discharges) and non-point sources (e.g. surface runoff). In general, water quality degradation is directly linked to anthropogenic pressure, and climate change acts as an indirect aggravating factor for these pressures on the environment. In particular, the main determinants of climate change having a direct or indirect impact on water quality are air temperature and extreme water events (flood, drought) (Delpla et al., 2009). Resource availability is linked to these parameters, and it becomes important to consider extreme events in assessing both evolution of water stress and efficiency of the treatment processes (sanitation and sewage treatment).

3. IMPACT OF RAINFALL ON URBAN WATER CYCLE (FROM WASTEWATER TO THE ENVIRONMENT)

The historical question of rainfall impact on urban water cycle is linked to urban runoff, well studied since the 70's. For instance, Pitt et al. (1977) studied the water-quality effects from urban runoff for a hypothetical city of 100,000 people. Total solids concentration and oxygen consumption were calculated for receiving waters using various storms simulated in the study. and more research on the long-term toxic effects of storm flow were suggested to help solve urban runoff problems.

Cities are becoming increasingly vulnerable to flooding because of rapid urbanization, installation of complex infrastructure and changes in the precipitation patterns caused by anthropogenic climate change (Willems et al; 2012). For more than a century, sewer systems have been constructed at large scale across cities worldwide. These sewer systems could make the cities more vulnerable to extreme rainfall, partly due to the lack of consideration to what occurs when the design criteria are exceeded. Next to this increase in the vulnerability, there is strong evidence that, due to global warming, the probabilities and risk to sewer surcharge and flooding are under discussion. But impacts of climate change on urban drainage are still weakly investigated (Willems et al., 2012).

In AMWA report (2007), the different impacts of climate changes on water suppliers are identified and described as follow:

- i. direct impacts resulting from the effects of climate change on water utility functions and operations,
- ii. indirect impact resulting from the effects of climate change on the baseline environment in which water utility functions and operations are carried out and
- iii. compound impact resulting from the cumulative effect of the direct and indirect impacts which affect the same natural systems or utility systems.

Among the direct impacts, more frequent and intense rainfall events are intended to increase turbidity, sedimentation, risk of direct flood and consequently damage water utility facilities: for example, loss of reservoir capacity storage (shallower, warmer water, increased evaporation and eutrophication and potential conflicts with flood control objectives) with consequences on water treatment (AMWA, 2007). However for urban watercourses, there is often "flashy" hydraulic regimes and poor water quality, making them potential hazards.

Furthermore, the knowledge of impact of intense rainfall events on urban river water discharge and, in particular, fine particle transport is limited (Old et al., 2003). The washed off pollutants such as suspended solids increases the turbidity of the receiving water, thereby reducing the penetration of light, resulting in decreased activity and growth of photosynthetic

organisms. Clogging of fish gills has also been attributed to the presence of suspended solids. These ones settle in receiving water pose long-term threats resulting from increased oxygen demand and gradual accumulation of toxic pollutants. Such detrimental effects have made suspended solid concentration one of the pollutant measurement indices (Aryal, 2009).

Moreover, Old et al. (2003) demonstrated that this extremely large and urban flux of fine sediment is likely to have a major downstream impact on water quality in the river.

Brodie and Egodawatta (2011) have shown that suspended particles in stormwater runoff generated from road surfaces loads increase linearly with average rainfall intensity. Above a threshold intensity, there is evidence to suggest a constant or plateau particle load is reached.

Cheng et al (2009) have studied the impact of four different intensity rainfall (light rain, moderate rain, heavy rain and storm) on water quality of urban combined sewerage system (CSS) in highly urbanized region and the correlations of parameters with both rainfall intensity and runoff graph. Basic parameters and metals concentrations were monitored and shown good relation between them and runoff.

Temporal evolution of microbiological, physical, and chemical quality of stormwater runoff from a stormwater drain was studied by He et al (2010) in an urban residential area in Calgary, Canada. Intensive rainfall events produced highly polluted stormwater runoff when pollutant source limitation did not occur. Inconsistent event-based correlations between total suspended solids (TSS) concentrations and water quality parameters were observed. During storms, the loading of TSS exhibited a flow-dependent nature, whereas microorganism discharge appeared to be governed by a flow-independent mechanism. No strong first-flush effect was observed in either TSS or microorganisms, on average. No correlations of first-flush loads of TSS with rainfall characteristics were identified.

Rainfall events can also play an important role regarding micropollutants load at sewage treatment works (STWs). In general, urban runoff tends to originate from five main categories: transportation (i.e. roads and vehicles), roofing materials, construction activities, vegetation and soil. There may be many variables within a given source, e.g. runoff from roads and associated traffic emissions may contain vehicle lubricants, exhaust emissions, degradation from tyres and brake linings, road maintenance and surface degradation, accidental spillages and road salting in winter. The extent of urban runoff is dependent on levels of contaminants deposited on urban surfaces through wet and dry deposition, the length of the preceding dry spell and the characteristics and size of the urban catchment. Assessing the effects of rainfall STW influent has produced results that suggest that with short rainfall incidences the level of most micropollutants will increase due to 'first flushing' events, whereas continuous high levels of rainfall will result in a dilution effect on micropollutant levels (Rowse et al., 2010).

Another consequence of runoff on water quality may be ecotoxicological impact linked to the presence of contaminants. Meland et al. (2010), exposed brown trout (*Salmo trutta* L.) to highway runoff samples during 24 h. The metabolism of the fish through reduced condition factor was affected and the observed effects were likely caused by multiple stressors and not by a single contaminant. A sedimentation pond clearly reduced the toxicity of the highway runoff. But even in the least polluted exposure condition signs of physiological disturbances were evident.

Considering the threats originating from wastewater utilities, the type of wastewater system is of crucial importance. More often a combined wastewater system is prevalent in urban area, with domestic wastewater and storm water runoff from impervious surfaces (streets, roofs etc.) collected in the same sewer system.

During rain events, the wastewater treatment plants as well as the sewer systems have to cope with a high load of combined wastewater that may exceed the systems' capacity. Thus, certain volumes of wastewater must be stored or discharged. This may be achieved by storm water overflow basins or storm water tanks (SWT), with both retention and treatment functions (Heins, 2009). Combined sewage overflows which were previously designed to directly discharge the exceeding flow tend to be removed from wastewater networks.

In case of heavy rainfall, the wastewater treatment plant temporarily receives low concentrated sewage and high hydraulic loading. Prolonged exposure to storm water conditions will negatively affect all treatment steps by decreasing the residence time and particularly the settling processes (primary and secondary) as well as the biological part of the treatment with activated sludge, and possibly denitrification and phosphorus removal processes. For this last step, it is still not clear whether this is due to inhibition of phosphorus-removing bacteria (PRB) or that the low COD concentrations in the influent are causing the problem. It has been reported that, for example, the deterioration of PRB efficiency regularly occurred at some wastewater treatment plants (WWTPs) after heavy rainfall. The phenomenon was attributed to low plant loading that took place during such events (Brdjanovic et al., 1998).

An observatory of urban pollutants was created in France for the purpose of assessing the dynamic of wastewater and wet weather flow (WWF). The magnitude of WWF pollutant loads characterized by physical-chemical parameters: total suspended solid, DOC, heavy metals (Cu, Zn), polycyclic aromatic hydrocarbons, and their subsequent acute impact on receiving waters have been extensively studied since 1970 (Suarez and Puertas, 2005; Gasperi et al., 2010). Some studies have observed the becoming of specific pollutants concentrations during wet weather flows in wastewater. Gilbert et al.,

(2011) for instance showed that alkylphenols (AP) and polybromodiphenylether (PBDE) concentrations are 1.5 to 5 fold higher under wet weather flows.

The removal rates of several antiphlogistics and lipid regulating agents was investigated during a rainfall event (Ternes, 1998). The elimination rates of drugs like bezafibrate, diclofenac, naproxen and clofibric acid were significantly reduced on the rainfall day. For bezafibrate a reduced elimination rate (<5%) was observed the last day of the investigation period. These results indicate that the rainfall event may be presumably responsible for the decreased drug elimination rates by the sewage treatment plant probably due to a reduced microbial activity and/or altered sorption and flocculation conditions within this rain period (Ternes, 1998).

4. RAINFALL AND WATER RESOURCE (QUALITY MODIFICATION OF NATURAL WATER)

Water quality depends on stressors among which climate-related parameters such as water temperature and river flows (Murdoch et al., 2000). According to these authors, changes in water quality is related to hydrological factors (e.g. limited dilution of point sources during low river flows, or increased runoff from agricultural area during rainstorms), terrestrial factors (e.g. changes in vegetation and soil structure) and resource-use factors (e.g. increased water use, increased demand for cooling capacity of industrial processes). More recently, bibliographic reviews and case studies (Delpla et al., 2009, Zwolsman and van Bokhoven, 2007, Van Vliet and Zwolsman, 2008) concluded that climate change has a negative impact on water quality. Depending on the season, the same impact (increase of water pollution) may be related to different conditions. During winter rainfall, water and air temperatures are low, river flows generally high and of soil leaching and runoff increase), contrary to summertime where very high air temperature may lead to drought (e.g. Krysanova et al., 2005). Both cases may give a high pollution level either by runoff (heavy rain in winter) or by physical concentration (drought in summer).

According to Bates et al. (2008) and Brunetti et al. (2001), climate change in temperate area will lead to a decrease of the number of rainy days, with however a significant increase of the mean volume for each event. Consequently, the drought and re-humidification cycles for the soils will allow the decomposition and leaching of the organic matter to the surface waters and thus impact water quality as a whole (Evans et al., 2005). Heavy rainfall events could lead to an increase in dissolved organic carbon concentration (Evans et al., 2005 ; Fellman et al., 2009) which is known to play an important role concerning the transport and release of pollutants (Lennartz and Louchart, 2007; Pédrot et al., 2008).

Nutrients concentrations (nitrates, phosphates) and obviously particulate and colloidal suspended matter are directly correlated with rainy events (Prathumratana et al., 2008; Bhat et al., 2007; Drewry et al., 2009) and could consequently exceed water quality regulated limits.

For heavy rainy events leading to an increase of river flows, the major risk is also certainly the transport of pollutants by run-off (such as pesticides and pharmaceuticals). It was often acknowledged that the timing and magnitude of rainfall events are significant on pesticide transport through rapid flow processes (Leonard, 1990, Flury, 1996; Kladvik et al., 2001, McGrath et al., 2010) with in general an increase of pesticide leaching with increasing rainfall amount (Tiktak et al. 2004, Jiang et al., 2012). Pharmaceuticals considered as « emerging substances » are of a special importance because they are not yet regulated. Some studies specifically show the variations of some of these compounds (carbamazepine, monensin, clofibric acid, iopromide, sulfamethoxazole, trimethoprim, flumequine) with flow rates variation during rainfall events (Oppel et al., 2004; Lissemore et al., 2006; Tamtam et al., 2008).

Variations of temperature, pH and water composition occurring during rainfall events could also have an influence on contaminants by impacting their sorption on mineral phases. Moreover, the influence of soils properties has to be taken into account for the leaching behaviour assessment of anthropogenic compounds (Oppel et al., 2004; Yu et al., 2009).

Neal et al (2004), studied the water quality of rainfall and runoff for two catchments of different characteristics. Average water quality concentrations and fluxes vary across the sites, typically by about 30% but stream chemistry is much less variable due to water coming from aquifer sources of high storage. Moreover, the catchments appear to be retain both P and N coming from agricultural practices and sewage inputs to the streams.

Some studies on coastal water quality impacts of rainfall like Coulliette et al. (2008) have shown the effect of non-point source contamination linked to stormwater. Fecal coliforms concentrations increased with rainfall intensity but higher than expected values existed during conditions of negligible rainfall (< 0.25 cm), indicating a possible reservoir population in the sediment.

Liu et al (2010), worked on the occurrence and distribution of three typical endocrine-disrupting chemicals (EDCs), nonylphenol mixture (NPs), bisphenol A (BPA), and 17-ethynilestrdiol (EE2), in the coastal water, suspended solid, and

sediment. With the increasing rainfall, the concentrations of target EDCs decreased in water and sediment and increased in suspended solid.

The genotoxic impact on fish of heavy rainfall following herbicide application has been investigated by Polard et al. (2011). For three hydrological conditions (basal flow, winter flood, and spring flood), chemical analysis of the water samples confirmed the higher contamination of the spring flood water, mainly explained by a peak of metolachlor. Genotoxicity was evaluated on Crucian carp and exposure to spring flood water resulted in the highest damage induction. Further experiments on pesticides mixture added on water flood revealed a mutagenic impact of water contamination during the spring flood, emphasizing the need to consider these transient events in water quality monitoring programs.

Concerning pathogens, floods frequently lead to a faecal contamination of surface and deep waters and also to waterborne diseases (Beaudeau et al., 2010). As an example, very elevated levels of bacteria *E. Coli* (2000 – 7000 CFU/mL) have been measured downstream Milwaukee river basin after rainfall events (Patz et al., 2008). Curriero et al., (2001) have shown that half of the waterborne diseases outbreaks in United States during the second half of the 20th century followed a period of extreme rainfall (defined as the excess of the 90th percentile based on 50-year distribution).

Finally changes in rainfalls patterns and temperature could also have an effect on blooms occurrence of cyanobacteria. Intense rainfalls could interrupt bloom development in a water body by flushing and mixing at the event scale but increase the nutrient load and the rate of eutrophication which favour cyanobacteria at a longer time scale (Reichwaldt and Ghadouani, 2011).

5. IMPACT ON DRINKING WATER PRODUCTION

Rainfalls events could lead to elevated levels of turbidity and organic matter in river waters which in turn could cause a decline in treatment performance. Mineral particles leaching could lead to high concentrations in natural waters, having a direct impact on coagulant demand during water treatment (Shin et al., 2008) and on Disinfection By-Products (DBPs) formation. However, this decrease in efficiency could be also due to a combination of a change in the nature and increased concentrations of Natural Organic Matter (NOM) and lower water temperatures in the natural water (Hurst et al., 2004). This could explain why seasonal differences have a significant impact on process robustness which is independent of the raw water turbidity. Moreover, in the distribution system, the highest spatial variation in Trihalomethans concentration (from 2 to 4 times) is measured for elevated water temperatures (>18°C) typically during the summer season, whereas this variation is small for water temperature lower than 15°C during the spring or fall seasons (Rodriguez and Serodes, 2001).

Although very few studies reported the relationship between algae and DBPs precursors, some authors pointed out the algal contribution to some DBPs formation such as Haloacetic Acids (Chen and Zhang, 2008).

Heavy rainfall and associated floods could flush biological contaminants into waterways and aquifers, for example with combined sewer systems overflowing and agricultural lands leaching. Greater water flow contributes to increase the pathogen load and the penetration speed of pathogens into resource waters and the drinking-water supply (Unc and Goss, 2003). A significant number of waterborne disease outbreaks have involved the transport of bacteria, viruses, or small parasites into water systems or well-heads during an heavy rainfall event (Auld et al., 2004; Thomas et al., 2006). As an example, heavy rainfalls contributed to the Milwaukee (United States) outbreak of *Cryptosporidium* in 1993 (54 deaths, 403000 cases) and the Walkerton (Canada) *E. Coli* O157:H7 outbreak in 2000 (7 deaths, 4300 cases). Monitoring the change of the biological charge with adequate and rapid tools is very important to ensure maximum efficiency of the treatment and protection of the population after rainfall events. This is of primary importance regarding the fact that routine monitoring programs mainly use techniques that are time consuming and not allow a representative picture of the wide range of biological contaminants that could be present in water.

The main recent studies dealing with the occurrence and fate of micropollutants with respect to drinking water treatment are related to pharmaceuticals. Actually, the majority of studies on pharmaceuticals focused on wastewater treatment, for which the insufficient removal efficiency may affect the quality of water resources downstream a treated effluent discharge. Although they are partially removed, residual quantities may remain in treated water and contaminate drinking water resources (Al-Ahmad et al., 1999; Hernando et al., 2006). Actually, more than 20 substances (parent compound and residues) have been found in drinking water (Mompelat et al., 2009).

The efficiency of pharmaceuticals removal varies with treatment processes and also with temperature and weather (Choi et al., 2008). For instance, diclofenac showed largely different elimination rates whose range, comprised between 17% (Heberer et al., 2002), 69% (Ternes et al., 1998), and 100% (Thomas and Foster, 2004) depends on these two last parameters. Pharmaceuticals removal could only be achieved with advanced processes as ozonation, activated carbon adsorption, membrane filtration (Ternes et al., 2002) or UV treatment (Canonica et al., 2008). Finally, only some lipophilic

pesticides are removed adequately in conventional physico-chemical drinking water treatment processes, such as flocculation, sedimentation, filtration, or lime softening (Baldauf, 2006).

Concerning natural micropollutants, mainly represented by cyanotoxins, chlorination, micro/ultra-filtration and especially ozonation are the most effective water treatment procedures in destroying cyanobacteria and in removing microcystins (Hitzfeld et al., 2000). Lots of studies concerning removal of cyanobacterial toxins from water showed that the effectiveness of the oxidation process is not only dependant on the reactant concentration, but also on temperature, pH, ionic composition (Rositano et al., 1998; Shawwa and Smith, 2001) and NOM concentration (Al Momani et al., 2008). Moreover, some studies tried to improve by-products identification and toxicity assessment for cyanotoxins (Rodriguez et al., 2007; Mérel et al., 2009). Although DBPs formation has largely not been investigated, the few studies conducted have concluded to a decrease in cyanotoxins toxicity (microcystin, cylindrospermopsin, saxitoxin) after chlorination (Mérel et al., 2010).

Extreme weather events could affect the efficiency of drinking water treatment processes, leading to a possible degradation of quality in distributed drinking water (Sinisi and Aertgeerts, 2010). These concerns are of primary importance for small-scale water suppliers (SSWSs), private, community-based or public, which usually prevail in rural areas. A significant part of the population living in these areas (16 million people in Europe (WHO, 2010) do not have access to improved drinking water resources. This vulnerability of SSWSs is due to an insufficient sanitary protection of resources, agricultural pressures, limited water treatment technologies, relatively greater capital costs of technical installations and larger per unit costs of materials and construction. By considering these existing pressures, and the indirect risks posed by climate change, SSWSs are facing several challenges, including their regulatory environment, administration, management, operation, and their available technical, human and financial resources (Sinisi and Aertgeerts, 2010).

CONCLUSION

Rainfall can be a significant source of variation in water quality. In urban area, depending on its intensity, rainfall can have an effect on utility functions and operation as well as water discharge due to runoff (road, roof, ...). The relation between water quality and runoff is in particular well known since the 70's as an increase of turbidity, sedimentation, transport of more or less fine particles and pollutants (conventional and/or emerging). The main outcomes were the improvement of sewer and rainstorm networks design with in situ settling systems for TSS removal and flow storage. More recently the effect of rainfall on soil erosion has been also highlighted, in particular the suspended solid exportation and increase of nutrient in surface water.

Water resource is also stressed by rainfall. Changes in water quality rely in particular to hydrological, terrestrial and resource-use factors as well as climate. Climate changes are more and more studied with consequences on micropollutants transportation and cyanobacteria crisis

One important point of the relation between water quality and rainfall is the evolution of transient effect (in particular toxicity and ecotoxicity) and the effect on sanitary water.

REFERENCES

- Al-Ahmad, A; Daschner, FD; Kummerer, K. Biodegradability of cefotiam, ciprofloxacin, meropenem, penicillin G, and sulfamethoxazole and inhibition of waste water bacteria. *Arch Environ Contam Toxicol* 1999, 37,158–63.
- Al Momani, F; Smith, DW; El-Din, MG. Degradation of cyanobacteria toxin by advanced oxidation process. *J Hazard Mater* 2008, 150, 238–49.
- AMWA (Association of metropolitan water agencies), *Implications of climate change for urban water utilities*, Washington DC, December 2007, 18 p.
- Auld, H; MacIver, D; Klaassen, J. Heavy rainfall and waterborne disease outbreaks: the Walkerton example. *J. Toxicol. Environ. Health A* 2004, 67, 1879-1887.
- Aryal, RK; Lee, B.-K. Characteristics of Suspended Solids and Micropollutants in First-Flush Highway Runoff. *Water Air Soil Pollut: Focus* 2009, 9, 339–346.
- Baldauf, G. Removal of pesticides in drinking water treatment. *Acta Hydrochim Hydrobiol* 2006, 21, 203–8.
- Bartram, J; Corrales, L; Davison, A; Deere, D; Drury, D; Gordon, B; Howard, G; Rinehold, A; Stevens, M; Water Safety Plan Manual: Step-By-Step Risk Management for Drinking-Water Suppliers, *World Health Organization*, Geneva, Switzerland, 108p, 2009.

- Bates, B; Kundzewickz, ZW; Wu, S; Palutikof, J. (Editors) Climate Change and Water, *Technical Paper of the Intergovernmental Panel on Climate Change, IPPC Secretariat*, Geneva, Switzerland, 210 p., 2008.
- Beaudeau, P; Valdes, D; Mouly, D; Stempfelet, M; Seux, R. Natural and technical factors in faecal contamination incidents of drinking water in small distribution networks, France, 2003-2004: a geographical study. *J. Water Health* 2010, 08.1, 20-34.
- Bhat, S; Hatfield, K; Jacobs, JM; Lowrance, R; Williams, R. Surface runoff contribution of nitrogen during storm events in a forested watershed. *Biogeochemistry* 2007, 85, 253-262.
- Bloech, H. *Summary Record of the Meeting of the Committee under Article 12 of Directive 98/83/EC*, DG ENV, 22/02/2011.
- Brdjanovic, D; Slamet, A; Van Loosdercht, MCM; Hooijmans, CM; Alaerts, GJ; Heijnen, JJ. Impact of excessive aeration on biological phosphorus removal from wastewater, *Water Research* 1998, 32, 200-208.
- Brodie, IM; Egodawatta, P. Relationships between rainfall intensity, duration and suspended particle washoff from an urban road surface, *Hydrology Research* 2011, 42, 239-249.
- Brunetti, M; Maugeri, M; Nanni, T. Changes in total precipitation rainy days and extreme events in Northeastern Italy. *Int. J. Climatol.* 2001, 21, 861-871.
- Canonica, S; Meunier, L; von Gunten, V. Phototransformation of selected pharmaceuticals during UV treatment of drinking water. *Water Res.* 2008, 42, 121-128.
- Chen, C; Zhang, X-J; Zhu, L-x; Liu, J; He, W-j; Han, H-d. Disinfection by-products and their precursors in a water treatment plant in North China: seasonal changes and fraction analysis. *Sci Total Environ* 2008, 397, 140-7.
- Cheng, J; Yang, K; Huang, X-F; Lue, Y-P. Hydrology and Water Quality of Rainfall-Runoff in Combined Sewerage System Along Suzhou Creek in Central Shanghai. *Environmental Science* 2009, 30, 1893-1900.
- Choi, K; Kima, Y; Park, J; Park, CK; Kim, M; Kim, HS; Kim, P. Seasonal variations of several pharmaceutical residues in surface water and sewage treatment plants of Han River. *Sci Total Environ* 2008, 405, 102-28.
- Coulliette, AD; Noble, R T. Impacts of rainfall on the water quality of the Newport River Estuary (Eastern North Carolina, USA). *Journal of Water and Health* 2008, 6, 473-482.
- Curriero, FC; Patz, JA; Rose, JB; Lele, S. The association between extreme precipitation and waterborne disease outbreaks in the United States, 1948-1994. *Am. J. Public Health*, 2001, 91, 1194-1199.
- Delpla, I; Jung, A-V; Baures, E.; Clement, M; Thomas, O. Impacts of climate change on surface water quality in relation to drinking water production. *Environment International* 2009, 35, 1225-1233.
- Drewry, JJ; Newham, LTH; Croke, BFW. Suspended sediment, nitrogen and phosphorus concentrations and exports during storm-events to the Tuross estuary, Australia. *J. Environ. Manag.* 2009, 90, 879-887.
- Evans, CD; Monteith, DT; Cooper, DM. Long-term increases in surface water dissolved organic carbon: observations, possible causes and environmental impacts. *Environment Pollution* 2005, 137, 55-71.
- Fellman, B; Hood, E; Edwards, RT; D'Amore, DV. Changes in the concentration, biodegradability, and fluorescent properties of dissolved organic matter during stormflows in coastal temperate watersheds. *Journal of Geophysical Research*, 2009, 114, G01021, doi:10.1029/2008JG000790.
- Flury, M. Experimental evidence of transport of pesticides through field soils — a review. *J. Environ. Qual.* 1996, 25, 25-45.
- Gasperi, J; Gromaire, MC; Kafi, M; Moilleron, R; Chebbo, G. Contributions of wastewater, runoff and sewer deposit erosion to wet weather pollutant load in combined sewer systems, *Water Research* 2010, 44, 5875-5886.
- Gilbert, DS; Gasperi, J; Rocher, V; Lorgeoux, C; Chebbo, G. Efficiency of the seine centre wastewater treatment plant during wet weather periods regarding conventional wastewater parameters, alkylphenols and polybromodiphenylethers, *Techniques Sciences et Méthodes* 2011, 4, 63-72.
- He, J.; Valeo, C.; Chu, A.; Neumann, N. F. Characteristics of Suspended Solids, Microorganisms, and Chemical Water Quality in Event-Based Stormwater Runoff from an Urban Residential Area. *Water Environment Research* 82. 12 (Dec 2010): 2333-2345.
- Heberer, T; Reddersen, K; Mechlinski, A. From municipal sewage to drinking water: fate and removal of pharmaceutical residues in the aquatic environment in urban areas. *Water Sci Technol* 2002, 46, 81-88.
- Heinz, B; Birk, S; Liedl, R; Geyer, T; Straub, KL; Andresen, J; Bester, K; Kappler, A. Water quality deterioration at a karst spring (Gallusquelle, Germany) due to combined sewer overflow: evidence of bacterial and micro-pollutant contamination. *Environmental Geology* 2009, 57, 797-808.
- Hernando, MD; Mezcuca, M; Fernandez-Alba, AR.; Barceló, D. Environmental risk assessment of pharmaceutical residues in wastewater effluents, surface waters and sediments. *Talanta* 2006, 69, 334-342.
- Hitzfeld, BC; Höger, SJ; Dietrich, DR. Cyanobacterial toxins removal during drinking water treatment and human risk assessment. *Environ Health Perspect* 2000, 108, 113-122.

- Hurst, AM; Edwards, MJ; Chipps, M; Jefferson, B; Parsons, SA. The impact of rainstorm events on coagulation and clarifier performance in potable water treatment. *Sci Total Environ* 2004, 321, 219–230.
- Jiang, W; Haver, D; Rust, M; Gan, J. Runoff of pyrethroid insecticides from concrete surfaces following simulated and natural rainfalls. *Water Res.* 2012, 46, 645–52.
- Kladivko, EJ; Brown, LC; Baker, JL. Pesticide transport to subsurface tile drains in humid regions of north America. *Crit. Rev. Environ. Sci. Tech.* 2001, 31, 1–62.
- Krysanova, V; Hattermann, F., Habeck, A. Expected changes in water resources availability and water quality with respect to climate change in the Elbe River basin (Germany). *Nordic Hydrology* 2005, 36, 321–333.
- Lennartz, B; Louchart, X. Effect of drying on the desorption of diuron and terbuthylazine from natural soils. *Environmental Pollution* 2007, 146, 180–187.
- Leonard, R., 1990. Movement of pesticides into surface waters. In: Cheng, H.H. (Ed.), *Pesticides in the Soil Environment. Soil Science Society of America*, Madison, USA, pp. 303–349.
- Lissemore, L; Hao, C; Yang, P; Sibley, PK; Mabury, S; Solomon, KR. An exposure assessment for selected pharmaceuticals within a watershed in Southern Ontario. *Chemosphere* 2006, 64, 717–729.
- Liu, Y.; Guan, Y.; Tam, NFY; Mizuno, T.; Tsuno, H.; Zhu, W. *Influence of Rainfall and Basic Water Quality Parameters on the Distribution of Endocrine-Disrupting Chemicals in Coastal Area. Water, Air, & Soil Pollution* 2010, 209 333–343.
- Meland, S; Salbu, B; Rosseland, BO. Ecotoxicological impact of highway runoff using brown trout (*Salmo trutta* L.) as an indicator model. *Journal of Environmental Monitoring* 2010, 12, 654–664.
- Mérel, S; Lebot, B; Clement, M; Seux, R; Thomas, O. MS identification of microcystin-LR chlorination by-products. *Chemosphere* 2009, 74, 832–9.
- McGrath, G; Hinz, C; Sivapalan, M. Assessing the impact of regional rainfall variability on rapid pesticide leaching potential. *J Contam Hydrol.* 2010, 113, 56–65.
- Mérel, S; Clément, M; Thomas, O. State of the art on cyanotoxins in water and their behaviour towards chlorine. *Toxicon* 2010, 55, 677–691.
- Mompelat, S; Le Bot, B; Thomas, O. Occurrence and fate of pharmaceutical products and by-products, from resource to drinking water. *Environ Int.* 2009, 35, 803–814.
- Murdoch, PS; Baron, JS; Miller, TL. Potential effects of climate change on surface water quality in North America. *Journal of the American Water Resources Association* 2000, 36, 347–366.
- Neal, C; Skeffington, R; Neal, M; Wyatt, R; Wickham, H; et al. Rainfall and runoff water quality of the Pang and Lambourn, tributaries of the River Thames, south-eastern England. *Hydrology and Earth System Sciences* 2004, 8, 601–613.
- Old, GH; Leeks, GJL; Packman, JC; Smith, BPG; Lewis, S; Hewitt, EJ; Homles, M; Young, A. The impact of convectional summer rainfall event on river flow on fine sediment transport in a highly urbanized catchment: Bradford, West Yorkshire, *The Science of the Total Environment* 2003, 314–316, 495–512.
- Oppel, J; Broll, G; Löffler, D; Meller, M; Römbke, J; Ternes, T. Leaching behaviour of pharmaceuticals in soil testing systems: a part of an environmental risk assessment for groundwater contamination. *Sci Total Environ* 2004, 328, 265–273.
- Patz, JA; Vavrus, SJ; Uejio, CK; McLellan, SL. Climate change and waterborne disease risk in the Great Lakes region of the US. *Am J Prev Med* 2008, 35, 451–458.
- Pédrot, M; Dia, A; Davranche, M; Bouhnik-Le Coz, M; Henin, O; Gruau, G. Insights into colloid-mediated trace element release at the soil/water interface. *J Colloid Interface Science* 2008, 325, 187–197.
- Pitt, R; Field, R. Water-Quality Effects from Urban Runoff. *J. Amer Water Works Assoc.* 1977, 69, 432–436.
- Polard, T; Jean, S; Gauthier, L; Laplanche, C; Merlina, G; et al. Mutagenic impact on fish of runoff events in agricultural areas in south-west France. *Aquatic Toxicology* 2011, 101, 126–134.
- Prathumratana, L; Sthiannopkao, S; Kim, KW. The relationship of climatic and hydrological parameters to surface water quality in the lower Mekong River. *Environ. Int.* 2008, 34, 860–866.
- Reichwaldt, ES; Ghadouani, A. Effects of rainfall patterns on toxic cyanobacterial blooms in a changing climate: Between simplistic scenarios and complex dynamics. *Water Research* 2012, 46, 1372–1393.
- Rodriguez, MJ; Serodes, JB. Spatial and temporal evolution of trihalomethanes in three water distribution systems. *Water Res* 2001, 35, 1572–1586.
- Rodriguez, E; Sordo, A; Metcalf, JS; Acero, JL. Kinetics of the oxidation of cylindospermopsin and anatoxin-a with chlorine, monochloramine and permanganate. *Water Res* 2007, 41, 2048–2056.
- Rositano, J; Nicholson, BC; Pieronne, P. Destruction of cyanobacterial toxins by ozone. *Ozone-Sci Eng* 1998, 20, 223–38.

- Rowell, VF; Tangney, P; Hunt, C; Voulvoulis, N. *Estimating levels of micropollutants in municipal wastewater, water, air and soil pollution* 2010, 206, 357-368.
- Shawwa, AR; Smith, DW. Kinetics of microcystin-LR oxidation by ozone. *Ozone-Sci Eng* 2001, 23, 161–170.
- Shin, JY; Spinette, RF; O'melia, CR. Stoichiometry of coagulation revisited. *Environ Sci Technol* 2008, 42, 2582–2589.
- Sinisi, L; Aertgeerts, R. *Guidance on Water Supply and Sanitation in Extreme Weather Events*, World Health Organization, Geneva, Switzerland, 2010.
- Suarez, J; Puertas, J. *Determination of COD, BOD and suspended solids loads during combined sewer overflow (CSO) event in some combined catchments in Spain, ecological engineering* 2005, 24, 199-217.
- Tamtam, F; Mercier, F; Le Bot, B; Eurin, J; Tuc Dinh, Q; Clément, M; Chevreuil, M. Occurrence and fate of antibiotics in the Seine River in various hydrological conditions. *Sci Tot Env* 2008, 393, 84 – 95.
- Ternes, TA. Occurrence of drugs in German sewage treatment plant and rivers, *Water Research* 1998, 32, 3245-3260.
- Ternes, T; Hirsch, R; Mueller, J; Haberer, K. Methods for the determination of neutral drugs as well as betablockers and alpha2-sympathomimetics in aqueous matrices using GC/MS and LC/MS/MS. *Fresen J Anal Chem* 1998, 362, 329–40.
- Ternes, T; Meisenheimer, M; McDowell, D; Sacher, F; Brauch, H-J; Haist-Glude, B; Preuss, G; Wilme, U; Zulei-Seibert, N. Removal of pharmaceuticals during drinking water treatment. *Environ Sci Technol.* 2002, 36, 3855-3863.
- Thomas, KM; Charron, DF; Waltner-Toews, D; Schuster, C; Maarouf, AR; Holt, JD. A role of high impact weather events in waterborne disease outbreaks in Canada, 1975 – 2001. *Int. J. Environ. Health Res.* 2006, 16, 167-180.
- Thomas, PM; Foster, GD. Determination of nonsteroidal anti-inflammatory drugs, caffeine, and triclosan in wastewater by gas chromatography-mass spectrometry. *J Environ Sci Health A* 2004, 39, 1969–1678.
- Tiktak, A; de Nie, DS; Piñeros Garcet, J; Jones, A; Vanclooster, M. Assessment of the pesticide leaching risk at the pan-European level: the EuroPEARL approach. *J. Hydrol.* 2004, 289, 222–238.
- Unc, A; Goss, M. Movement of Faecal Bacteria through the Vadose Zone. *Water Air Soil Pollut.* 2003, 149, 327-337.
- Van Vliet, MTH; Zwolsman, JJG. Impact of Summer Droughts on the Water Quality of the Meuse River. *Journal of Hydrology*, 2008, 353, 1-17.
- Willems, P; Arnbjerg-Nielsen, K; Olsson, J; Nguyen, VTV. Climate change impact assessment on urban rainfall extremes and urban drainage: methods and shortcomings, *Atmospheric Research* 2012, 103, 106-118
- World Health Organization (WHO), Vision 2030 – *The Resilience of Water Supply and Sanitation in the Face of Climate Change*, World Health Organization/Department for International Development, Geneva, Switzerland, 2010 (http://www.who.int/water_sanitation_health/publications/9789241598422_cdrom/en/index.html, accessed 5 January 2011).
- Yu, L ; Fink, G ; Wintgens, T ; Melin, T ; Ternes, TA. Sorption behaviour of potential organic wastewater indicators with soils. *Water Res* 2009, 43, 951–60.
- Zwolsman, JJG; Van Bokhoven, A. Impact of summer droughts on the water quality of the Rhine River - a preview of climate change? *Water Science & Technology*, 2007, 56, 45–55.