

Evolution of Copernicus Land Services based on Sentinel data
3rd EARSeL SIG LU/LC and NASA LCLUC joint Workshop
Wednesday, 11 June 2018
Chania, Crete

ECoLaSS



ECoLaSS:

Evolution of Copernicus Land Services based on Sentinel data

Linda Moser (GAF AG)

Probeck, M., Ramminger, G., Rieke, C., Mack, B., Ickerott, M., Storch, C., Sommer, C., Richter, R., Herrmann, D., Ruiz, I., Kovatsch, M., and Schwab K.














ECoLaSS: "Evolution of Copernicus Land Services based on Sentinel data"


Key Objective = **improve** existing & develop **novel** products/services for future operational pan-European & Global Components of the **CLMS for 2020+**



ECoLaSS

(H2020 Grant Agreement no. 730008)

www.ecolass.eu ©ECoLaSS2020



Test- and Demonstration sites in various biogeographic regions.












1) FOREST: Sentinel-2 Time Features (Mar – Aug 2017)

Demonstration Site: North (Subset: Sweden)

S-2, 2017-05-27, R: B08, G: B04, B: B03

S-2 Time Features (2017-03-15 to 2017-06-15)



- Input: S-2 bands, S-2 indices (NDVI, NDWI, BRIGHTNESS, IRECI)
- Time interval: Mar-Aug 2017
- 60 Features selected












1) FOREST Dominant Leaf Type / Forest Change - Prototype Copernicus HRL

Demonstration Site: North (Subset: Sweden)

S-2, 2017-05-27, R: B08, G: B04, B: B03







DLT Overall Accuracy (OA) = 97%

DLT Producer's Accuracy = 86-98%, DLT User's Accuracy = 90-99%












1) FOREST Dominant Leaf Type / Forest Change - Prototype Copernicus HRL

Demonstration Site: North (Subset: Sweden)

S-2, 2017-05-27, R: B08, G: B04, B: B03





DLT Overall Accuracy (OA) = 97%DLT Producer's Accuracy = 86-98%, DLT User's Accuracy = 90-99%













ernicus Land 2020

2) AGRICULTURE: Sentinel-1 & -2 Time Features

Demonstration Site: Central (Subset: Baden-Württemberg, Germany)



S-2, 2017-03-15 to 2017-11-14 Time Feature: B08_std



Input: S-1 bands, S-1 indices, S-2 bands, S-2 indices (NDVI, NDWI, BRIGHTNESS, IRECI)

Time intervals: Mar-Nov and 2-monthly (Mar-May, May-Jul, Jul-Sep, Sep-Nov)

Forward feature selection: 28 selected time features (from ~500)











Corporations Land 2020X

2) AGRICULTURE: Crop Mask/Type - Potential future Copernicus Agricultural

Demonstration Site: Central (Subset: Baden-Württemberg, Germany)



Crop Type Map (16 crop types)



LPIS data © MLR BW

Produced using Copernicus Sentinel data [2017]

© European Union, Copernicus Land Monitoring Service 2015, European Environment Agency (EEA).

Maize (PA=100%, UA=98%), SugarBeets (PA=98%, UA=100%), WinterCrop (PA=100%, UA=96%), WinterRape (PA=100%, UA=100%), SummerRape (PA=15%, UA=54%), Sunfl/Topinamb (PA=20%, UA=85%)

Overall Accuracy (OA) = 93%Producer's Accuracy = 15-100% User's Accuracy = 54-100%










2) AGRICULTURE: Crop Mask/Type - Potential future Copernicus Agricultural

Demonstration Site: Central (Subset: Baden-Württemberg, Germany)



Maize (PA=100%, UA=98%), SugarBeets (PA=98%, UA=100%), WinterCrop (PA=100%, UA=96%), WinterRape (PA=100%, UA=100%), SummerRape (PA=15%, UA=54%), Sunfl/Topinamb (PA=20%, UA=85%)

Overall Accuracy (OA) = 93%Producer's Accuracy = 15-100%User's Accuracy = 54-100%

LPIS data © MLR BW

Produced using Copernicus Sentinel data [2017]

European Union, Copernicus Land Monitoring Service 2015, European Environment Agency (EEA).











Thank you on behalf of the ECoLaSS team !!

Contact: linda.moser@gaf.de (Project Coordinator)

L. Moser, M. Probeck, G. Ramminger, D. Herrmann, K. Schwab & Team

C. Sannier, S. Villerot, B. Desclée & Development Team

H. Gallaun, M. Schardt, P. Miletich, J. Deutscher, K. Granica

P. Defourny, I. Moreau, J. Wolter, O. Mohamed, X. Blaes

A. Metz-Marconcini, A. Hirner, T. Esch



Follow us:







The research leading to these results has received funding from the European Union's Horizon 2020 Research and Innovation Programme, under Grant Agreement no 730008.



















