Engineering Probiotics for Simultaneous Degradation of Lactose and β-Casomorphin-7 to Alleviate Dairy-Associated Gastrointestinal Symptoms

Sophie Xu

7/11/2026

Abstract

Although dairy intolerance is commonly attributed to lactose intolerance, growing evidence suggests that A1 β-casein may also contribute to gastrointestinal discomfort. During digestion, A1 casein releases β-casemorphin-7 (BCM-7), a bioactive opioid peptide that may produce symptoms such as delayed gastrointestinal transit, slower gut motility, and inflammatory responses through the activation of μ-opioid receptors. This paper reviews the current understanding of A1 β-casein-associated gastrointestinal intolerance, including the difference between A1 and A2 β-casein, the role of BCM-7, the challenges associated with diagnosis, and the limitations of existing management strategies. Building upon this background, a conceptual engineered probiotic approach is proposed that exploits the lactose-inducible lac operon to coordinate expression of β-galactosidase and a BCM-7-degrading peptidase, DPP4, during dairy consumption. Although this proposal remains theoretical, it presents a biologically plausible framework for developing a targeted enzymatic approach to reduce active BCM-7 during digestion. If successfully developed, this strategy could provide an alternative to dietary restriction and contribute to future therapeutic options for individuals experiencing dairy intolerance.

Introduction

Although dairy-associated discomfort is most commonly attributed to lactose intolerance, lactose intolerance represents only one of several conditions that may contribute to adverse reactions following dairy consumption. Milk protein is made up of 80% casein and 20% whey, and it generally contains two types of casein: A1 and A2. There has been an increase in the awareness of other factors that contribute to dairy intolerance, such as A1 β-casein intolerance or allergy. A1 β-casein digestion is associated with increased gastrointestinal (GI) transit time and some inflammatory markers compared with A2 β-casein (Barnett et al., 2014). Common symptoms of A1 β-casein intolerance include eczema, diarrhea, and bloating, which while similar to lactose intolerance, differ in reaction mechanisms and response times. Due to an increase in the population that experiences dairy intolerance and the limited awareness towards casein intolerance, it is necessary to develop a strategy to counter the harmful effects that casein may cause. A1 β-casein releases β-casomorphin-7 (BCM-7), which activates μ-opioid receptors in the GI tract and may contribute to the symptoms (Pal et al., 2015). Dipeptidyl peptidase 4 (DPP-4) is the primary human enzyme known to efficiently break down BCM-7 into X-Pro dipeptide at the N-terminus position, which can potentially relieve the symptoms caused by the production of BCM-7 (Giribaldi et al., 2022). Current management of dairy-associated gastrointestinal symptoms primarily relies on dietary restriction or single-enzyme supplements, leaving limited options for individuals with various kinds of dairy intolerance. This paper proposes the development of an engineered probiotic containing β-galactosidase and a microbial BCM-7-degrading peptidase - DPP4 - as a targeted therapeutic strategy to simultaneously address multiple mechanisms underlying dairy-associated gastrointestinal discomfort.

Background



Figure 1. The evolution history of A1 and A2 milk. Adapted from (Prasad MG et al., 2024). Created with the assistance of ChatGPT.

Bovine milk is valued by many because of its nutritional value, including its contribution to 28 of the 29 nutrients that humans need (de Vasconcelos et al., 2023). Milk is composed mainly of milk proteins, which consist of casein and whey protein. Specifically, casein is further categorized into four subtypes: αs1-casein, αs2-casein, β-casein, and κ-casein (Dmour & Taha, 2018). Each of the four subtypes has dozens of different genetic variants. This paper focuses on A1-β-casein, which is one of the genetic variants of β-casein (Zinßius et al., 2024). Another genetic variant of β-casein, A2-β-casein, is considered to be the ancestral form of A1-β-casein, because it existed before the point mutation - from proline to histidine - that caused the creation of A1-β-casein in some European herds 5,000-10,000 years ago (Pal et al., 2015). *Figure 1* illustrates the evolutionary divergence of A1 and A2 β-casein variants, suggesting that A2 casein emerged before A1 casein.

The presence of histidine allows the enzyme to perform a cleavage of β-CN, which in turn causes the release of β-casomorphin-7 (BCM-7) (Giribaldi et al., 2022). Casomorphin can be defined as any opioid released from casein during digestion (Woodford, 2021). The proline-rich sequence makes it resistant to proteases, resulting in some intact BCM-7 left behind after digestion, which can then bind to the μ-opioid receptors located in the central nervous system and gastrointestinal tracts of humans, delaying GI transit time and decreasing gut motility by reducing intestinal contractions (Asledottir et al., 2019). Moreover, BCM-7 is also known to increase mucus secretion in the gastrointestinal system, contributing to excessive mucus that disrupts normal intestinal function.

Ecem Bolat et al. (2024) reviewed experimental evidence indicating that BCM-7 may activate μ-opioid receptors, increase intestinal permeability, and stimulate the production of

pro-inflammatory cytokines, including IL-6 and TNF-α. However, the authors also emphasized that current evidence remains insufficient to establish a direct causal relationship between BCM-7 and human diseases.

Although considerable research has investigated the relationship between A1 β-casein, BCM-7, and gastrointestinal discomfort, the results are inconclusive. Therefore, current management strategies primarily rely on dietary restriction rather than targeted interventions. This highlights the need for alternative approaches that address BCM-7 directly.

Differentiating Lactose Intolerance, Milk Allergy, and A1 β-casein Intolerance

Although the gastrointestinal symptoms associated with A1 β-casein intolerance may resemble those of lactase intolerance, these conditions differ in their underlying mechanisms. Lactose intolerance results from lactose deficiency and carbohydrate malabsorption, whereas the mechanism of A1 β-casein intolerance remains under investigation and has been hypothesized to involve bioactive peptides such as BCM-7. Distinguishing these conditions is important because they require different diagnostic approaches and management strategies.

Condition	Cause	Trigger	Mechanism	Diagnostic Test
Lactose intolerance	Lactase deficiency	Lactose	Carbohydrate malabsorption	Breath Test
Milk Allergy	Immune response	Milk proteins	IgE/non-IgE immune reaction	Allergy testing
A1 β-casein intolerance	Under investigation	Potentially BCM-7	Opioid/inflammatory response	No validated clinical test

Table 1. This table compares and contrasts the cause, trigger, mechanism, and current diagnostic test of lactose intolerance, milk allergy, and A1 β-casein intolerance.

Lactose intolerance is an enzymatic reaction that can cause symptoms such as abdominal pain, diarrhea, and bloating. People with lactose intolerance don't have enough lactase to digest lactose, leaving some lactose undigested, which is then fermented by bacteria in the large intestine. Lactose intolerance can now be detected by a breath test at most hospitals.

Conversely, a milk allergy is an adverse immune reaction to milk proteins, and most common in infants and children. Three types of reaction mechanisms can be presented in a milk allergy: IgE-mediated allergies, delayed-onset non-IgE cell-mediated allergies, and a mix of both (Hao et al., 2025). Milk proteins can act as antigens, which bind to antibodies and stimulate a series of immune responses. Symptoms include red, blotchy skin that can itch, difficulty breathing, tearing or redness of the eyes, and so on (Spergel, 2025).

Finally, A1 β-casein intolerance is a kind of dairy intolerance that is still being investigated. While the appearance of symptoms of lactose intolerance is often rapid, it can take up to 72 hours for the symptoms of casein intolerance to show. Symptoms often include bloating, stomach cramps, eczema, and rashes (Casein, n.d.).

Structural and Biochemical Differences Between A1 and A2 β-Casein

Compared to conventional milk, studies have shown that bovine milk containing only A2 casein doesn't usually generate gastrointestinal discomfort. *Figure 2* demonstrates the different intermediates that A1/A2 casein release upon enzymatic cleavage, which can be attributed to the

amino acid substitution in Position 67. Since A1 casein has His instead of Pro, it's easier to be broken down by enzymes, which contributes to the release of the shorter intermediate BCM-7.




Figure 2. The structural differences between A1 and A2 casein and the generation of BCM-7 upon enzymatic cleavage. Created via canva.

While BCM-7 is released during the breakdown of A1 casein, a longer peptide, BCM-9, is released when A2 casein is digested. BCM-7 and BCM-9 are both opioid, but they perform different functions: BCM-7 is linked to inflammatory and digestive disorders such as bloating and irritable bowel patterns (University, 2024), while BCM-9 is considered beneficial, potentially both antihypertensive and antioxidant (Woodford, 2021).

Potential Mechanisms Underlying A1 Casein-Associated Intolerance

Currently, there is no confirmed cause of A1 casein intolerance. It is important to note that much research in this field is inconclusive, and no microbiological cause of A1 casein intolerance has been proven as the single explanation. However, there are some hypotheses that could be made between the enzymatic activities or gut permeability and biome and the symptoms of A1 casein intolerance. For example, the lack of DPP4 activity can contribute to impaired degradation of

BCM-7. Variations in DPP4 activities may influence BCM-7 persistence in the GI tract and therefore represent a potential factor contributing to individual differences in tolerance of A1 casein. Asledottir et al., (2019) used GI fluids and porcine jejunal brush border membrane (BBM) peptidases to study the digestion of synthetic BCM-7. Results demonstrated that 5% of BCM-7 remained intact after 24 hours BBM digestion. However, the percentage of BCM-7 that persists after digestion can vary among individuals, and this variability can contribute to different responses from A1 casein, because the intact BCM-7 can interact with the opioid receptors in the gut, contributing to gastrointestinal discomfort.

Furthermore, increased gut permeability also serves as a potential contributor to inflammatory responses towards casein. A possible explanation is that since BCM-7 can trigger inflammation, it can lead to possible barrier dysfunction and thus increased gut permeability. Brooke-Taylor et al., (2017) use rodent models to demonstrate that the ingestion of A1 casein can be linked to the initiation of inflammatory responses. Inflammation and gut permeability reinforce each other, forming a cyclic pathway that attenuates the immune system. As a result, a “leaky gut” can form and contribute to increased exposure of BCM-7 to the gut opioid receptors, which might explain the gastrointestinal discomfort associated with the ingestion of A1 casein.

Current Management Strategies and Their Limitations

Current methods for casein intolerance are limited, with only some medical testing for casein allergy but not intolerance. Moreover, unlike lactose intolerance - which can be diagnosed using hydrogen breath tests; or milk allergy - which can be evaluated using skin prick tests, serum IgE tests, or oral food challenges, there is currently no validated laboratory or clinical test for A1 β-casein-associated gastrointestinal intolerance. Individuals may undergo food sensitivity testing

to determine whether he or she is intolerant to dairy products, but still cannot specify the kind of intolerance. At present, suspected A1 β-casein intolerance is usually evaluated indirectly. If people still exhibit gastrointestinal discomfort after consuming dairy products and being negative in milk allergy tests and lactose intolerance tests, one can try eliminating dairy products in their diets to see whether there are other components in dairy products that may have caused the discomfort.

There are laboratory tests for milk protein allergy but none for casein intolerance, as it is more of a digestive problem rather than immune system reactions. Testing of milk allergy usually involves IgE antibodies, and it is helpful for individuals who suspect they have a milk allergy (Test, 2019). However, as previously noted, A1 β-casein intolerance cannot be revealed with this laboratory test; therefore, there is still room for improvements regarding methods of detecting milk intolerance.

The biggest challenge is that casein intolerance symptoms are largely nonspecific. Most of the symptoms in casein intolerance are also common in lactose intolerance, milk allergy, and other gastrointestinal disorders. Because of this overlap, patients are frequently misdiagnosed or assume they have lactose intolerance when the underlying cause may be different.

One approach to reduce the discomfort associated with A1 casein is to replace conventional milk with A2 milk. In contrast to traditional bovine milk, A2 milk contains only A2 casein, which cuts off the production of BCM-7. By eliminating the dietary source of BCM-7, individuals with intolerance may experience reduced symptoms related to A1 casein. For example, Jianqin et al.

(2016) conducted a randomized crossover study in adults with self-reported milk intolerance and found that consumption of milk containing only A2 β-casein resulted in significantly lower gastrointestinal discomfort and reduced inflammatory markers compared with conventional milk containing both A1 and A2 casein. However, this approach does not address situations in which individuals consume conventional dairy products, such as cheese, yogurt, and cream. Consequently, while A2 milk represents a potential solution, it does not provide a therapeutic approach for individuals who are unable or unwilling to avoid conventional milk and dairy products.

Proposal

Design Rationale

Dipeptidyl peptidase 4 (DPP4) is a proline-specific serine protease that can break down proteins containing proline or alanine in the N terminus next to last (P1) position, resulting in altered function of the original protein (Aljumaah et al., 2025). Human dipeptidyl peptidase 4 (hDPP-4) plays a critical role in homeostasis - it regulates metabolic and systemic processes and modulates the functionality of over 40 bioactive substrates. hDPP4 also has numerous peptide substrates, including the incretin hormones glucagon-like-peptide-1 (GLP-1) and glucose-dependent insulinotropic polypeptide (GIP). (Mulvihill & Drucker, 2014) BCM-7, a peptide released during the digestion of A1 β-casein, is rich in proline and has therefore been investigated as a potential substrate for DPP4. However, the relationship between BCM-7 degradation and gastrointestinal symptoms remains controversial, meaning that BCM-7 degradation should be considered a potential therapeutic strategy rather than an established treatment.

A major challenge in developing a targeted therapy for A1 casein intolerance is identifying an appropriate biological signal that can be used to signal the production of BCM-7 degrading enzymes at the appropriate location and time. BCM-7 itself may be difficult to use as a regulatory signal because current research suggests that there is no established transcriptional system that directly detects BCM-7 and activates production of a degrading enzyme. Although it would be difficult to detect BCM-7, detecting lactose is much easier because lactose binds to the repressor that inactivates the lac operon. Given that casein's presence is often accompanied by lactose, it is reasonable to assume that if structural genes that code for hDPP4 are inserted downstream of the structural genes encoding for β-galactosidase, the presence of lactose can contribute to the production of more DPP4 enzymes. It is important to note that although DPP4 is capable of degrading BCM-7, it does not hydrolyze lactose. Therefore, β-galactosidase remains necessary to digest lactose and alleviate symptoms for those who are lactose intolerant.

A probiotic chassis provides an additional advantage because it could serve as an orally administered, local enzyme-delivery system. Engineered bacteria have been investigated as "living therapeutics" that can deliver biological functions directly within the gastrointestinal tract, potentially allowing therapeutic molecules to be produced locally rather than administered systematically. *E.coli* Nissle 1917 (EcN) is particularly relevant because it has a long history of use as a probiotic and has been investigated as a chassis for engineered microbial therapeutics due to its genetic tractability and ability to survive within the gastrointestinal tract. (Praveschotinunt et al., 2019)

In sum, the proposed system asims to use a food-associated regulatory signal to coordinate two functions: β-galactosidase production to support lactose digestion and a modified DPP4-derived peptidase to investigate BCM-7 degradation. Importantly, the goal is not simply to maximize DPP4 expression. Because DPP4 has numerous peptide substrates, the design must prioritize BCM-7 degradation while minimizing unintended activity toward physiologically important peptides. (Mulvihill & Drucker, 2014)

Proposed Genetic Design

The proposed engineered probiotic would use *E.coli* Nissle 1917 (EcN) as the initial chassis. It is a probiotic strain with a history of human use and has been investigated as a genetically tractable platform for engineered live biotherapeutics, and its use provides an appropriate starting point for an orally administered bacterial therapeutic. Compared to convential laboratory *E.coli*, EcN provides a more appropriate starting platform for an oral live biotherapeutic.

The genetic system will be designed to respond to lactose in the intestinal environment. A lactose-responsive regulatory system based on the lac operon will control the expression of two therapeutic functions: β-galactosidase and a DPP4-derived peptidase. When lactose is present, increased β-galactosidase production would promote lactose breakdown, while the second enzyme would be intended to facilitate the degradation of BCM-7 released from β-casein digestion. This creates a coordinated response in which the presence of a dietary component associated with the proposed symptoms activates the therapeutic system.

Therapeutic Applications

The proposed engineered EcN would be administered orally as a live biotherapeutic intended to act within the gastrointestinal tract. Previous research has demonstrated the feasibility of engineering EcN to produce therapeutic proteins within the gut, supporting its potential as a localized delivery platform. (Praveschotinunt et al., 2019) After reaching the gut, the engineered bacteria would respond to dietary lactose and produce β-galactosidase to support lactose digestion. At the same time, the engineered DPP4-derived peptidase would be intended to degrade BCM-7 generated during the digestion of A1 β-casein. By targeting both lactose and BCM-7, the system is designed to address two proposed contributors to gastrointestinal discomfort associated with A1 milk.

Using an engineered probiotic as the delivery platform could allow the enzymes to be produced locally in the gastrointestinal tract rather than requiring them to remain active throughout the entire digestive process before ingestion. This concept is supported by previous work demonstrating in situ production of therapeutic proteins by engineered EcN. The proposed lactose-responsive design could further restrict enzyme production to conditions in which the dietary target is present.

Advantages

One of the major advantages of this proposed therapeutic is that it simultaneously addresses two potential factors that contribute to symptoms related to intolerance. Current lactase supplements only hydrolyze lactose, so some individuals might still experience discomfort even after taking lactase. By engineering bacteria capable of producing both β-galactosidase and

BCM-7-degrading peptidases during dairy digestion, this probiotic approach may provide broader symptom relief than single-enzyme therapies.

Challenges and Limitations

A major challenge is that human DPP4 is naturally a type II transmembrane glycoprotein and functions as a homodimer. Therefore, expressing the full-length human protein in a bacterial chassis may not reproduce the structure or activity of native DPP4. Instead, the design will investigate a soluble DPP4-derived form that retains the extracellular catalytic domain while avoiding dependence on the native transmembrane region. Protein dimerization, folding, and the effects of glycosylation will need to be evaluated because these structural features may influence enzymatic activity.

The DPP4-derived enzyme will also require evaluation for substrate specificity. Although DPP4 can cleave a range of peptides, including physiologically important substrates such as GLP-1, GIP, and PYY, indiscriminate DPP4 activity could interfere with normal physiological signaling. Therefore, the proposed design should prioritize characterization of the enzyme's activity toward BCM-7 and assessment of potential off-target activity before therapeutic application is considered.

Conclusion

Since the domestication of cattle, humans have consumed milk as a major source of nutrition throughout life. Thus, detecting casein intolerance is difficult yet crucial regarding health and fitness. Many people value milk, seeing it as a beneficial choice to maintain good health and support body development; however, undesired reactions to milk can cause detrimental effects,

such as chronic gastrointestinal distress and systemic inflammation, which can lead to chronic diseases.

While milk is scientifically proven to provide various benefits, the underlying costs are still being explored. For example, Chia et al., 2017 presents a potential correlation between the increased rate of Type 1 diabetes and heightened milk consumption: In Shanghai, the incidence among children aged ⩽15 years increased at a rate of 14.2% per year between 1997 and 2011, which is mirrored by an increase in per capita dairy product consumption among urban residents of 12 kg from nearly 6 kg in 1992 to 18 kg by 2006. Although both environmental factors and genetic factors play important roles in developing Type 1 diseases, this research team argues that the consumption of A1 milk is the key environmental trigger.




Figure 6. Proposed model for progression to Type 1 diabetes in genetically susceptible people. Adapted from (Chia et al., 2017). Created with the assistance of ChatGPT.

This proposal thus offers a novel therapeutic strategy for addressing dairy-associated gastrointestinal discomfort by simultaneously targeting lactose and BCM-7. Future experimental studies should focus on validating the catalytic efficiency of recombinant DPP4 toward BCM-7, evaluating enzyme stability under gastrointestinal conditions, and determining whether dual-enzyme supplementation provides better clinical outcomes compared with conventional lactase supplements. If successfully developed, this dual-enzyme supplement could offer individuals with greater dietary flexibility while reducing reliance on strict dairy avoidance, ultimately contributing to improved quality of life and expanding future therapeutic options for dairy-associated gastrointestinal intolerance.

References

A1 Beta-casein Protein Intolerance. (n.d.-a). Milk Digestion. https://milkdigestion.com/adverse-milk-reactions/a1-beta-casein-protein-intolerance/

A1 Protein Intolerance vs Lactose Intolerance - Betta Nutrition Professional. (2017). Betta Nutrition Professional. https://betaa2professional.com.au/a1-protein-intolerance-vs-lactose-intolerance/

Aljumaah, M. R., Roach, J., Hu, Y., Gunstad, J., & M. Andrea Azcarate-Peril. (2025). Microbial dipeptidyl peptidases of the S9B family as host-microbe isozymes. *Science Advances*, *11*(14), eads5721–eads5721. https://doi.org/10.1126/sciadv.ads5721

Anguita-Ruiz, A., Aguilera, C. M., & Gil, Á. (2020). Genetics of Lactose Intolerance: An Updated Review and Online Interactive World Maps of Phenotype and Genotype Frequencies. *Nutrients*, *12*(9). https://doi.org/10.3390/nu12092689

Asledottir, T., Picariello, G., Mamone, G., Ferranti, P., Røseth, A., Devold, T. G., & Vegarud, G. E. (2019). Degradation of β-casomorphin-7 through in vitro gastrointestinal and jejunal brush border membrane digestion. *Journal of Dairy Science*, *102*(10), 8622–8629. https://doi.org/10.3168/jds.2019-16771

Bayanati, M., Ismail, M., Kabiri, S. S., Mir, B., Elham Rezaee, & Tabatabai, S. A. (2024). Dipeptidyl Peptidase-4 Inhibitors: A Systematic Review of Structure-Activity Relationship Studies. *Deleted Journal*, *23*(1). https://doi.org/10.5812/ijpr-151581

Briden, L. (2021, November 26). *The Inflammation From A1 Milk Is Mind-Boggling*. Lara Briden - The Period Revolutionary. https://www.larabriden.com/the-inflammation-from-a1-milk-is-mind-boggling/

Brooke-Taylor, S., Dwyer, K., Woodford, K., & Kost, N. (2017). Systematic Review of the Gastrointestinal Effects of A1 Compared with A2 β-Casein. *Advances in Nutrition: An International Review Journal*, *8*(5), 739–748. https://doi.org/10.3945/an.116.013953

Casein. (n.d.-b). YorkTest. https://www.yorktest.com/intolerance/casein-intolerance/

Casein Sensitivity | Signs & Symptoms - YorkTest US. (2025, August 12). York Test US. https://www.yorktest.com/us/intolerance/casein/

Chia, J. S. J., McRae, J. L., Kukuljan, S., Woodford, K., Elliott, R. B., Swinburn, B., & Dwyer, K. M. (2017). A1 beta-casein milk protein and other environmental pre-disposing factors for type 1 diabetes. *Nutrition & Diabetes*, *7*(5), e274–e274. https://doi.org/10.1038/nutd.2017.16

de Vasconcelos , M. L., F S Oliveira, L. M., Paul Hill, J., & Centola Vidal, A. M. (2023, August 22). *Difficulties in Establishing the Adverse Effects of β-Casomorphin-7 Released from β-Casein Variants—A Review*. Nih.Gov. https://pmc.ncbi.nlm.nih.gov/articles/PMC10486734/#sec1-foods-12-03151

Dipeptidyl peptidase 4. (2026). NCBI. https://www.ncbi.nlm.nih.gov/datasets/gene/1803/#summary

Dmour, I., & Taha, M. O. (2018). Natural and semisynthetic polymers in pharmaceutical nanotechnology. *Organic Materials as Smart Nanocarriers for Drug Delivery*, 35–100. https://doi.org/10.1016/b978-0-12-813663-8.00002-6

E. coli as a Model Organism: Advantages & Limitations | Boster Bio. (2024). Bosterbio.Com. https://www.bosterbio.com/blog/post/e-coli-as-a-model-organism?srsltid=AfmBOor6sfYLRc3Ja67IAyJyzofzLOZNv6Mcm8d4z-UxtCA8GJb-1zp0

Ecem Bolat, Eker, F., Selin Yılmaz, Sercan Karav, Oz, E., Brennan, C., Charalampos Proestos, Zeng, M., & Oz, F. (2024). BCM-7: Opioid-like Peptide with Potential Role in Disease Mechanisms. *Molecules*, *29*(9), 2161–2161. https://doi.org/10.3390/molecules29092161

Federica Di Vincenzo, Angelo Del Gaudio, Petito, V., Loris Riccardo Lopetuso, & Franco Scaldaferri. (2023). Gut microbiota, intestinal permeability, and systemic inflammation: A narrative review. *Internal and Emergency Medicine*, *19*(2), 275–293. https://doi.org/10.1007/s11739-023-03374-w

Giribaldi, M., Lamberti, C., Cirrincione, S., Giuffrida, M. G., & Cavallarin, L. (2022). A2 Milk and BCM-7 Peptide as Emerging Parameters of Milk Quality. *Frontiers in Nutrition*, *9*. https://doi.org/10.3389/fnut.2022.842375

Hao, L., Wang, S., & Ji, W. (2025). Cow's milk protein allergy: A comprehensive review of epidemiology, pathogenesis, clinical manifestations, diagnostics, and management strategies. *PubMed*, *34*(3), 298–307. https://doi.org/10.6133/apjcn.202506_34(3).0004

Jianqin, S., Leiming, X., Lu, X., Yelland, G. W., Ni, J., & Clarke, A. J. (2016). Effects of milk containing only A2 beta casein versus milk containing both A1 and A2 beta casein proteins on gastrointestinal physiology, symptoms of discomfort, and cognitive behavior of people with self-reported intolerance to traditional cows' milk. *Nutrition Journal*, *15*(1). https://doi.org/10.1186/s12937-016-0147-z

Khan Academy. (2021). *The Lac Operon*. Khan Academy. https://www.khanacademy.org/science/ap-biology/gene-expression-and-regulation/regulation-of-gene-expression-and-cell-specialization/a/the-lac-operon

Lactose Operon - an overview | ScienceDirect Topics. (2016). Sciencedirect.Com. https://www.sciencedirect.com/topics/neuroscience/lactose-operon

Marketingteam Histaminikus. (2024, August). *Lactose intolerance and casein intolerance*. Histaminikus. https://histaminikus.de/en/blogs/blog/lactose-intolerance-and-casein-intolerance

Milk Protein - an overview | ScienceDirect Topics. (n.d.-c). Www.Sciencedirect.Com. https://www.sciencedirect.com/topics/food-science/milk-protein

Mulvihill, E. E., & Drucker, D. J. (2014). Pharmacology, physiology, and mechanisms of action of dipeptidyl Peptidase-4 inhibitors. *Endocrine Reviews*, *35*(6), 992–1019. https://doi.org/10.1210/er.2014-1035

Newmark, L. (2018). *Milk Casein Proteins: Ancient, Diverse, and Essential*. International Milk Genomics Consortium. https://www.milkgenomics.org/?splash=milk-casein-proteins-ancient-diverse-essential

OpenAI. (2023). ChatGPT (Mar 14 version) [Large language model]. https://chat.openai.com/chat

Pal, S., Woodford, K., Kukuljan, S., & Ho, S. (2015). Milk Intolerance, Beta-Casein and Lactose. *Nutrients*, *7*(9), 7285–7297. https://doi.org/10.3390/nu7095339

Park, Y. (2020). *A2 Milk and CMPA | Encyclopedia MDPI*. Encyclopedia.Pub. https://encyclopedia.pub/entry/history/show/21404

Prasad MG, Diksha Purushottam Gourkhede, Vidyarani HB, Shinde, B., Bidyut Prabha Mishra, Pratik Ramesh Wankhade, Balaji Belore, Lalthanmawii, J., & Koneti, P. B. (2024). Delving into the A1/A2 milk hypothesis: A comprehensive analysis of milk proteins and their impact on human health. *International Journal of Veterinary Sciences and Animal Husbandry*, *9*(1S), 594–605. https://doi.org/10.22271/veterinary.2024.v9.i1si.1082

Praveschotinunt, P., Duraj-Thatte, A. M., Gelfat, I., Bahl, F., Chou, D. B., & Joshi, N. S. (2019). Engineered E. coli nissle 1917 for the delivery of matrix-tethered therapeutic domains to the gut. *Nature Communications*, *10*(1). https://doi.org/10.1038/s41467-019-13336-6

Spergel, J. (2025). *Cow's milk protein allergy (CMPA)*. Chop.Edu. https://www.chop.edu/conditions-diseases/cows-milk-protein-allergy

Tellam, R. (2022, February 28). *74 - International Milk Genomics Consortium*. International Milk Genomics Consortium. https://www.milkgenomics.org/issue/74/

Test, C. A. (2019). *Casein Allergy Test*. Healthlabs.Com. https://www.healthlabs.com/casein-allergy-testing?srsltid=AfmBOooMx3cxin4XaG8eGQh7Vm0OczV5okgM8Vvb5Q8jLltn9AC7QDo4

University, S. (2024, September 25). *Understanding A1 and A2 Milk – Unpacking the Controversial Debate*. Sageuniversity.Edu.In; Sage University - Bhopal. https://sageuniversity.edu.in/blogs/understanding-a1-and-a2-milk-unpacking-the-controversial-debate

Vella, A. (2012). Mechanism of Action of DPP-4 Inhibitors—New Insights. *The Journal of Clinical Endocrinology & Metabolism*, *97*(8), 2626–2628. https://doi.org/10.1210/jc.2012-2396

Woodford, K. B. (2021). Casomorphins and Gliadorphins Have Diverse Systemic Effects Spanning Gut, Brain and Internal Organs. *International Journal of Environmental Research and Public Health*, *18*(15), 7911. https://doi.org/10.3390/ijerph18157911

Zinßius, L., Keuter, L., Krischek, C., Jessberger, N., Cramer, B., & Plötz, M. (2024). *Influence of the β-Casein Genotype of Cow's Milk (A1, A2) on the Quality and β-Casomorphin-7*

(BCM-7) Content of a Semi-Hard Cheese During Production. Nih.Gov.

https://pmc.ncbi.nlm.nih.gov/articles/PMC11817730/