

Computational Statistics & Machine Learning

Lecture 3

Lebesgue Integral and Measure

Mark Girolami

`mag92@cam.ac.uk`

Department of Engineering

University of Cambridge

October 19, 2021

Lecture Outline

Riemann Integral

Convergence of
Functions

Lebesgue Integral

- ▶ Riemann Integral.
- ▶ Convergence of Functions.
- ▶ Lebesgue Integral.

Riemann Integral

4M24

M.Girolami

Lecture Outline

Riemann Integral

Convergence of
Functions

Lebesgue Integral

- ▶ Formal definition and construction of Integral
- ▶ Integration is the fundamental operator that appears repeatedly in Machine Learning - it is important



- ▶ Bernhard Riemann 1862

Riemann Integral

4M24

M.Girolami

Lecture Outline

Riemann Integral


Convergence of
Functions

Lebesgue Integral

- ▶ We are familiar with integrals of the form

$$I = \int_a^b f(x) dx$$

- ▶ The domain (\mathbb{R} or \mathbb{R}^d) is split into infinitesimal units
- ▶ In the interval $[a, b] \subset \mathbb{R}$ then $\Delta x_k = [x_k, x_{k+1}]$



- ▶ So $a = x_1 < x_2 < x_3 < \dots < x_{n+1} = b$ defines the partition, P , of the domain of integration
- ▶ Define two summations

$$S_P(x) = \sum_{k=1}^n M_k(x_{k+1} - x_k), \quad s_P(x) = \sum_{k=1}^n m_k(x_{k+1} - x_k)$$

- ▶ With $M_k = \sup_{x \in \Delta x_k} f(x)$ and $m_k = \inf_{x \in \Delta x_k} f(x)$ being the *supremum* and *infimum* of $f(x)$ in Δx_k .
- ▶ What Riemann does is to take limit as $n \rightarrow \infty$ over all possible partitions P to define

$$S(f) = \liminf_{n \rightarrow \infty} S_P \quad s(f) = \limsup_{n \rightarrow \infty} s_P$$

- ▶ Then if $S(f) = s(f) = A$, the value A is the Riemann Integral of $f(x)$ denoted as $A = \int_a^b f(x) dx$
- ▶ The bounded function $f(x)$ must be continuous **almost everywhere** on the domain of integration for the integral to exist

Upper and Lower Riemann Sums

4M24

M.Girolami


Lecture Outline

Riemann Integral



Convergence of
Functions

Lebesgue Integral

Lower Riemann Sum



Upper Riemann Sum



- ▶ For applications in Machine Learning the Riemann Integral has a number of shortcomings
- ▶ The construction of the integral requires a partition of the domain, for \mathbb{R}^d this is natural.
- ▶ What about the domain comprised of continuous functions starting at zero i.e. $C[0, \infty]$, how is this divided into subintervals of equal size?
- ▶ Domains other than \mathbb{R}^d are common in Machine Learning and need to be handled appropriately
- ▶ The Lebesgue Integral provides an ingenious resolution to this issue

- ▶ The other shortcoming is that the exchange of limit operations only applies under stringent conditions



$$\lim_{n \rightarrow \infty} \int_{-\infty}^{+\infty} f_n(x) dx = \int_{-\infty}^{+\infty} \lim_{n \rightarrow \infty} f_n(x) dx$$


- ▶ This is a common operation in Machine Learning applications such as function approximation

Riemann Integral

4M24

M.Girolami

- Think of the function $f_n(x) = 2n^2x \exp(-n^2x^2)$ in the unit interval $[0, 1]$ for all integer n



- The function is continuous on the unit interval and so is Riemann integrable
- $\lim_{n \rightarrow \infty} \int_0^1 f_n(x) dx = \lim_{n \rightarrow \infty} (1 - \exp(-n^2)) = 1$
- $\int_0^1 \lim_{n \rightarrow \infty} f_n(x) dx = \int_0^1 0 dx = 0$
- hmmm what is going on?

Lecture Outline

Riemann Integral

Convergence of
Functions

Lebesgue Integral

Convergence of Functions

4M24

M.Girolami

Lecture Outline

Riemann Integral

Convergence of
Functions

Lebesgue Integral

- ▶ We need to look at the convergence of functions to a limit
- ▶ This is important in optimisation for example when training a Deep Net
- ▶ There are a number of senses in which a sequence of functions will converge
- ▶ The two that we will consider at this point are *Pointwise* and *Uniform* convergence
- ▶ We shall observe the difference in these and why the Riemann Integral requires the stricter form of convergence

- ▶ Consider a sequence of real valued functions f_n with $n = 1, 2, \dots$ i.e. $n \in \mathbb{N}$
- ▶ Each f_n is a real valued function $f_n(x)$ of $x \in D \subseteq \mathbb{R}$
- ▶ If the sequence $f_1(x), f_2(x), \dots, f_n(x), \dots$ converges to $f(x)$ and does so for every $x \in D$
- ▶ This is said to converge *pointwise* - at each point x - on D - the pointwise limit is $f(x) = \lim_{n \rightarrow \infty} f_n(x)$
- ▶ Formally the sequence of functions $f_1(x), f_2(x), \dots, f_n(x), \dots$ converges pointwise to f on D if given $\epsilon > 0$ there is a natural number $N = N(\epsilon, x)$ such that for $n > N$ it holds that $|f_n(x) - f(x)| < \epsilon$
- ▶ Convergence will vary depending on x in a pointwise manner

Pointwise Convergence

- ▶ A simple example illustrates an important facet of pointwise convergence
- ▶ The sequence of continuous functions on $[0, 1]$ is defined as the polynomials $f_n(x) = x^n$
- ▶ The sequence of continuous functions converges to

$$f(x) = \lim_{n \rightarrow \infty} f_n(x) = \begin{cases} 0 & \text{for } 0 \leq x < 1 \\ 1 & \text{at } x = 1 \end{cases}$$

- ▶ This is not a continuous function, the sequence of continuous functions converges to one that is discontinuous at $x = 1$
- ▶ Pointwise convergence **DOES NOT** preserve continuity of functions
- ▶ So

$$\lim_{n \rightarrow \infty} \int_{-\infty}^{+\infty} f_n(x) dx \neq \int_{-\infty}^{+\infty} \lim_{n \rightarrow \infty} f_n(x) dx$$

- ▶ The Riemann Integral requires a stronger form of convergence of functions that does preserve continuity

- ▶ For each $x \in D$ one can select an $N(\epsilon, x)$ in a pointwise manner so convergence is governed by each point selected
- ▶ Alternatively in certain cases one can select $N(\epsilon)$ independent of x so that $|f_n - f| < \epsilon$ for $n > N$ for all $x \in D$. The convergence is uniform over all x
- ▶ Formally the sequence of functions $f_1(x), f_2(x), \dots, f_n(x), \dots$ converges uniformly to f on D if given $\epsilon > 0$ there is a natural number $N = N(\epsilon)$ such that for $n > N$ it holds that
$$|f_n(x) - f(x)| < \epsilon \quad \forall x \in D$$
- ▶ Convergence will be uniform and independent of choice of x . Each f_n is contained in an ϵ tube on D around f
- ▶ Uniform convergence does preserve continuity as required for the limiting process of Riemann Integration

Pointwise vs. Uniform Convergence

4M24

M. Girolami



Lecture Outline

Riemann Integral

Convergence of
Functions

Lebesgue Integral

- ▶ The function $f_n(x) = x^n$, $x \in [0, 1]$ converges pointwise to $f(x) = \begin{cases} 0 & \text{for } x \in [0, 1) \\ 1 & \text{for } x = 1 \end{cases}$
- ▶ The function $g_n(x) = x/n$, $x \in [0, 1]$ converges uniformly to $g(x) = 0$



Lebesgue Integral

4M24

M.Girolami

Lecture Outline

Riemann Integral

Convergence of
Functions

Lebesgue Integral

- ▶ Formal definition and construction of Integral
- ▶ Integration is the fundamental operator that appears repeatedly in Machine Learning - it is important



- ▶ Henri Lebesgue 1875

Lebesgue Integral

4M24

M.Girolami

Lecture Outline

Riemann Integral

Convergence of
Functions

Lebesgue Integral

- ▶ The construction of the Riemann Integral proceeds by making a partition of the domain
- ▶ For the general types of domains we meet in Machine Learning this partitioning is problematic
- ▶ What Lebesgue ingeniously did was to partition the range of the function which would take values in \mathbb{R} and so can be done straightforwardly
- ▶ Once the range is partitioned the size or measure of parts of the domain that correspond to the range partition are identified
- ▶ This exploits the notion of the size or measure of a general set, so for the set of continuous functions a measure needs to be defined - this turns out to be more straightforward than partitioning such domains
- ▶ In addition, as the range of the function is partitioned, the strict continuity requirement of the Riemann integral is relaxed

Lebesgue Integral

4M24

M.Girolami


Lecture Outline

Riemann Integral

Convergence of
Functions

Lebesgue Integral

- ▶ Consider the bounded function $f(x)$ defined on an abstract set X such that $0 \leq f_{\min} \leq f(x) \leq f_{\max}$
- ▶ Now partition the $f(x)$ axis using the sequence $f_{\min} = f_1, f_2, \dots, f_n = f_{\max}$
- ▶ There will be sets of values of x such that $f_k \leq f(x) < f_{k+1}$ for $x \in X_k$
- ▶ The size (we will now start referring to this as the *measure*) of each of these sets X_k will have value $\mu(X_k)$





Lebesgue Integral

4M24

M.Girolami

- Compare the Riemann integral to the Lebesgue integral



Lecture Outline

Riemann Integral

Convergence of
Functions

Lebesgue Integral

- ▶ The Lebesgue sum of products of the function value and the measure of the corresponding set is

$$\sum_{k=1}^n f_k \mu(X_k)$$

- ▶ If the Lebesgue sum is convergent as $n \rightarrow \infty$ such that $|f_k - f_{k+1}| \rightarrow 0$
- ▶ The limiting value is the Lebesgue Integral of the function $f(x)$ over the set X

$$\int_X f d\mu = \lim_{\max |f_k - f_{k-1}| \rightarrow 0} \sum_{k=1}^n f_k \mu(X_k)$$

- ▶ The Riemann Integral is inextricably linked to the natural ordering of \mathbb{R} - i.e. $1 < 2 < 3.4 < 3.5$ etc... integrating on other structures is challenging
- ▶ The Lebesgue integral only requires a measure of sets on the domain of integration so is more general
- ▶ Monotone and Dominated convergence Theorems of Lebesgue Integral allow change of order of limit and integration requiring only pointwise convergence
- ▶ If a function is Riemann integrable it is Lebesgue integrable - the converse is not strictly the case e.g. discontinuous functions