

# Computational Statistics & Machine Learning

## Lecture 10

### Gaussian Measures in Hilbert Space

Mark Girolami

`mag92@cam.ac.uk`

Department of Engineering

University of Cambridge

November 3, 2021

## Lecture Outline

Lebesgue Measure  
in infinite  
dimensions

Measures in  
infinite dimensions

Gaussian Measures  
in infinite  
dimensions

Full covariance  
operators in  
infinite dimensions

Gaussian Measures  
in Hilbert Space

Equivalence of  
Measures

Equivalence of  
Measures

Bayes Rule in  
Hilbert Space

- ▶ Lebesgue Measure in infinite dimensions
- ▶ Gaussian Measure in Hilbert space
- ▶ Equivalence and Singularity of Gaussian Measures in infinite dimensions
- ▶ Bayes formula in Hilbert space

- ▶ We know that Lebesgue measure is a generalisation of notions such as length, area and volume
  - ▶ For every interval  $I = [a, b]$  on  $\mathbb{R}$  then
$$\mu(I) = l(I) = b - a$$
  - ▶ If  $A \subset B \subset \mathbb{R}$  then  $0 \leq \mu(A) \leq \mu(B) \leq \infty$
  - ▶ For each subset  $A$  of  $\mathbb{R}$  and each point  $x_0 \in \mathbb{R}$  then
$$A + x_0 = \{x + x_0 : x \in A\} \text{ and } \mu(A + x_0) = \mu(A) + \mu(x_0) = \mu(A)$$
- ▶ Finiteness
- ▶ Monotonicity
- ▶ Translation invariance

# Lebesgue Measure in infinite dimensions

4M24

M.Girolami

Lecture Outline

Lebesgue Measure  
in infinite  
dimensions

Measures in  
infinite dimensions

Gaussian Measures  
in infinite  
dimensions

Full covariance  
operators in  
infinite dimensions

Gaussian Measures  
in Hilbert Space

Equivalence of  
Measures

Equivalence of  
Measures

Bayes Rule in  
Hilbert Space


- ▶ Lebesgue measure in  $D$  dimensions  $\prod_{d=1}^D (b_d - a_d)$
- ▶ What happens when  $D = \infty$ ?
- ▶ Lets take a Hilbert space,  $H$ , and see what happens
- ▶ We know there is a countable orthonormal set acting as a basis in  $H$ , i.e.  $\{e_i | i = 1, \dots, \infty\}$  with  $\langle e_i, e_j \rangle = \delta_{ij}$
- ▶ How far apart are they?  
 $\|e_i - e_j\|^2 = \|e_i\|^2 + \|e_j\|^2 = 1 + 1 = 2$ , if  $i \neq j$
- ▶ Lets take a ball centered at 0 with radius 2 denoted  $B(0, 2)$
- ▶ Lets place balls of radius  $\frac{1}{2}$  at each  $e_i$  i.e.  
 $\{B(e_i, \frac{1}{2}), i = 1, \dots, \infty\}$
- ▶ It should be clear that  $B(e_i, \frac{1}{2}) \cap B(e_j, \frac{1}{2}) = \emptyset$
- ▶ ... and  $\bigcup_{i \in \mathbb{N}} B(e_i, \frac{1}{2}) \subset B(0, 2)$

# Lebesgue Measure in infinite dimensions

4M24

M.Girolami

- ▶ Lets take a ball centered at 0 with radius 2 denoted  $B(0, 2)$
- ▶ Lets place balls of radius  $\frac{1}{2}$  at each  $e_i$  i.e.  $\{B(e_i, \frac{1}{2}), i = 1, \dots, \infty\}$
- ▶ Then  $B(e_i, \frac{1}{2}) \cap B(e_j, \frac{1}{2}) = \emptyset$  and  $\bigcup_{i \in \mathbb{N}} B(e_i, \frac{1}{2}) \subset B(0, 2)$



Lecture Outline

Lebesgue Measure  
in infinite  
dimensions

Measures in  
infinite dimensions

Gaussian Measures  
in infinite  
dimensions

Full covariance  
operators in  
infinite dimensions

Gaussian Measures  
in Hilbert Space

Equivalence of  
Measures

Equivalence of  
Measures

Bayes Rule in  
Hilbert Space

# Lebesgue Measure in infinite dimensions

- ▶ Now if Lebesgue measure in infinite dimensions is translation invariant then  $\mu(B(e_i, \frac{1}{2})) = \mu(B(e_j, \frac{1}{2}))$ , for all  $i, j \in \mathbb{N}$
- ▶ The monotonicity and countable additivity of Lebesgue measure tells that

$$\sum_{i \in \mathbb{N}} \mu\left(B\left(e_i, \frac{1}{2}\right)\right) \leq \mu(B(0, 2))$$

- ▶ But wait..... Lebesgue measure is (sigma) finite so  $\mu(B(0, 2)) < \infty$
- ▶ But  $\sum_{i \in \mathbb{N}} \mu(B(e_i, \frac{1}{2})) = \text{constant} \times \sum_{i \in \mathbb{N}} 1 = \infty$
- ▶ Meaning that  $\mu(B(0, 2)) \geq \infty$
- ▶ There is no Lebesgue measure in infinite dimensional Hilbert (or Banach) spaces
- ▶ This is a problem discuss why this is a problem (think about how probability densities are defined)

Lecture Outline

Lebesgue Measure  
in infinite  
dimensions

Measures in  
infinite dimensions

Gaussian Measures  
in infinite  
dimensions

Full covariance  
operators in  
infinite dimensions

Gaussian Measures  
in Hilbert Space

Equivalence of  
Measures

Equivalence of  
Measures

Bayes Rule in  
Hilbert Space

- ▶ Most of Machine Learning takes place in infinite dimensional spaces
  - ▶ Function approximation - Kernel methods
  - ▶ Stochastic processes in Signal Processing
  - ▶ Dirichlet Processes in probabilistic modelling
  - ▶ Gaussian Processes in Bayesian Machine Learning
  - ▶ Inverse Problems in all aspects of engineering science
- ▶ Require a reference measure for the function spaces we are reliant upon in Machine Learning

# Gaussian Measures in infinite dimensions

- ▶ Lets take a function we are familiar with the standard Gaussian in  $\mathbb{R}$

$$g(B) = \frac{1}{\sqrt{2\pi}} \int_B \exp\left(-\frac{1}{2}x^2\right) dx$$

- ▶ Define the infinite dimensional product measure on  $\mathbb{R}^\infty$  as  $\mu = \prod_{n=1}^\infty \mu_n$  where each  $\mu_n = g$
- ▶ Consider the function

$$f_\lambda(x) = \exp\left(-\frac{\lambda}{2} \sum_{k=1}^\infty a_k x_k^2\right)$$

- ▶ Well defined only if  $\sum_{k=1}^\infty a_k x_k^2 < \infty$
- ▶  $x \in l_{2,a} = \{x \in \mathbb{R}^\infty; \sum_{k=1}^\infty a_k x_k^2 < \infty\}$
- ▶  $l_{2,a}$  a Hilbert space, inner product  $\langle x, y \rangle = \sum_{k=1}^\infty a_k x_k y_k$

Lecture Outline

Lebesgue Measure  
in infinite  
dimensionsMeasures in  
infinite dimensionsGaussian Measures  
in infinite  
dimensionsFull covariance  
operators in  
infinite dimensionsGaussian Measures  
in Hilbert SpaceEquivalence of  
MeasuresEquivalence of  
MeasuresBayes Rule in  
Hilbert Space



# Gaussian Measures in infinite dimensions

- By using the infinite dimensional standard Gaussian product measure we obtain the result

$$\int_{\mathbb{R}^\infty} f_\lambda d\mu = \frac{1}{\sqrt{\prod_{k=1}^\infty (1 + \lambda a_k)}}$$

- There are a number of things we can see here
  - We have replaced the Lebesgue measure with the product Gaussian measure in  $d\mu$
  - If the infinite product converges then  $\int_{\mathbb{R}^\infty} f_\lambda d\mu$  is finite and positive
  - If  $\sum_{k=1}^\infty a_k < \infty$  then the product converges
  - Note that  $\lambda \rightarrow 0$  then  $f_\lambda(x) = 1$  if  $x \in l_{2,a}$
  - Also as  $\lambda \rightarrow 0$  then
 
$$\int_{\mathbb{R}^\infty} f_\lambda d\mu = \int_{\mathbb{R}^\infty} I_{x \in l_{2,a}} d\mu = \int_{l_{2,a}} d\mu = \mu(l_{2,a}) = 1$$
  - This is awesome - we have a probability measure which gives full measure to the whole sequence space - and as it is isomorphic to  $L^2$  to the corresponding function space

Lecture Outline

Lebesgue Measure  
in infinite  
dimensionsMeasures in  
infinite dimensionsGaussian Measures  
in infinite  
dimensionsFull covariance  
operators in  
infinite dimensionsGaussian Measures  
in Hilbert SpaceEquivalence of  
MeasuresEquivalence of  
MeasuresBayes Rule in  
Hilbert Space

- ▶ Passing over many technical details it follows that Gaussian measure in  $l^2$  (Hilbert space), with mean zero and diagonal covariance operator  $Sx = (a_n x_n)_{n=1}^\infty$ ,  $x \in l^2$  if  $\sum_{k=1}^\infty a_k < \infty$
- ▶ It means that the infinite dimensional sequence  $x \in l^2$  has finite Gaussian measure we can then define a probability space over functions this is getting to be practically very important in Machine learning
- ▶ We have a measure which is finite and well defined in an infinite dimensional space and can be used in taking Radon-Nikodym derivatives in changing measures in e.g. spaces of functions

# Full covariance operators in infinite dimensions

- ▶ We have considered only diagonal covariance operators so far
- ▶ More general full covariance operators can be defined and used in the definition of the Gaussian measure
- ▶ Given the convergence criterion to be met for the summation of the diagonal terms a similar requirement must be met for a full covariance operator
- ▶ If we have a linear operator  $C$  on a Hilbert space,  $H$ , it has to be *trace class* (or nuclear) meaning that  $\text{trace}(C) < \infty$
- ▶ The trace operator in Hilbert space with an orthonormal basis is defined by

$$\text{trace}(C) = \sum_{n=1}^{\infty} \langle e_n, C e_n \rangle$$

what does  $\langle e_n, C e_n \rangle$  look like?

Lecture Outline

Lebesgue Measure  
in infinite  
dimensions

Measures in  
infinite dimensions

Gaussian Measures  
in infinite  
dimensions

Full covariance  
operators in  
infinite dimensions

Gaussian Measures  
in Hilbert Space

Equivalence of  
Measures

Equivalence of  
Measures

Bayes Rule in  
Hilbert Space

# Full covariance operators in infinite dimensions

- ▶ Let  $f$  and  $g$  be two zero-mean functions defined on a domain  $D$  such that  $f, g \in H$
- ▶ Define  $\langle f, Cg \rangle$  as

$$\begin{aligned}
 \langle f, Cg \rangle &= \mathbb{E}(f, u)(u, g) \text{ with } u \in H \\
 &= \mathbb{E} \int_D \int_D f(x)(u(x)u(y))g(y)dydx \\
 &= \mathbb{E} \int_D f(x) \left( \int_D u(x)u(y)g(y)dy \right) dx \\
 &= \int_D f(x) \left( \int_D \mathbb{E}\{u(x)u(y)\}g(y)dy \right) dx \\
 &= \int_D f(x) \left( \int_D c(x, y)g(y)dy \right) dx
 \end{aligned}$$

- ▶ So  $Ce_n(x) = \int_D c(x, y)e_n(y)dy$
- ▶ For those of you studying Probabilistic Machine Learning recognise the *covariance function* defining a Gaussian Process

Lecture Outline

Lebesgue Measure  
in infinite  
dimensionsMeasures in  
infinite dimensionsGaussian Measures  
in infinite  
dimensionsFull covariance  
operators in  
infinite dimensionsGaussian Measures  
in Hilbert SpaceEquivalence of  
MeasuresEquivalence of  
MeasuresBayes Rule in  
Hilbert Space

# Gaussian Measures in Hilbert Space

- ▶ We have a finite measure in Hilbert spaces
- ▶ The Gaussian is a reference measure in Hilbert space
- ▶ It is clearly not translation invariant it is quasi invariant
- ▶ Consider the Radon-Nikodym derivative between two Gaussian measures in  $\mathbb{R}$  i.e.  $\mu = \mathcal{N}(0, \sigma)$  and  $\mu_m = \mathcal{N}(m, \sigma)$

- ▶ Then

$$\frac{d\mu_m}{d\mu}(x) = \exp\left(-\frac{m^2}{2\sigma^2} + \frac{xm}{\sigma^2}\right)$$

- ▶ Both  $\mu$  and  $\mu_m$  have positive densities with respect to Lebesgue measure
- ▶ It follows that for  $\sigma \neq 0$  then for any  $A \in \mathbb{R}$ ,  $\mu(A) = 0$  if and only if  $\mu_m(A) = 0$  if and only if  $\text{Lebesgue}(A) = 0$
- ▶ The measures are *absolutely continuous* (or equivalent) with respect to each other


Lecture Outline

Lebesgue Measure  
in infinite  
dimensionsMeasures in  
infinite dimensionsGaussian Measures  
in infinite  
dimensionsFull covariance  
operators in  
infinite dimensionsGaussian Measures  
in Hilbert SpaceEquivalence of  
MeasuresEquivalence of  
MeasuresBayes Rule in  
Hilbert Space



# Singular Measures

4M24

M.Girolami



$$\nu(A) = 0$$



$$\mu(B) = 0$$

Lecture Outline

Lebesgue Measure  
in infinite  
dimensions

Measures in  
infinite dimensions

Gaussian Measures  
in infinite  
dimensions

Full covariance  
operators in  
infinite dimensions

Gaussian Measures  
in Hilbert Space

**Equivalence of  
Measures**

Equivalence of  
Measures

Bayes Rule in  
Hilbert Space

- ▶ Things get a little more extreme in infinite dimensions
- ▶ Two measures are absolutely continuous if they both assign zero measure to the same sets
- ▶ Two measures are singular with respect to each other if one measure assigns zero to a set and the other measure assigns zero measure to the set complement
- ▶ In Hilbert space two Gaussian measures will either be equivalent or singular see Jupyter Notebook for example
- ▶ This has serious implications in the design of various Computational Statistical methods and Machine Learning algorithms as the following lecture will demonstrate

[Lecture Outline](#)[Lebesgue Measure in infinite dimensions](#)[Measures in infinite dimensions](#)[Gaussian Measures in infinite dimensions](#)[Full covariance operators in infinite dimensions](#)[Gaussian Measures in Hilbert Space](#)[Equivalence of Measures](#)[Equivalence of Measures](#)[Bayes Rule in Hilbert Space](#)

- ▶ Given a Gaussian measure  $\mathcal{N}(0, C)$  on  $l^2$  the shifted Gaussian measure  $\mathcal{N}(v, C)$  is either equivalent to  $\mathcal{N}(0, C)$  or they are singular
- ▶ There is a subspace containing all  $v$  for which the measures are equivalent
- ▶ The subspace for all translations  $v$  for which the measures are absolutely continuous, is the image of the space  $l^2$  under the operator  $C^{\frac{1}{2}}$
- ▶ This space is known as the Cameron-Martin space (also the Reproducing kernel Hilbert Space - RKHS)

Lecture Outline

Lebesgue Measure  
in infinite  
dimensions

Measures in  
infinite dimensions

Gaussian Measures  
in infinite  
dimensions

Full covariance  
operators in  
infinite dimensions

Gaussian Measures  
in Hilbert Space

Equivalence of  
Measures

Equivalence of  
Measures

Bayes Rule in  
Hilbert Space



# Bayes Rule in Hilbert Space

- ▶ Bayes formula defines a posterior probability measure in terms of a likelihood and prior probability measure

$$\mathbb{P}(u|y) = \frac{\mathbb{P}(y|u)\mathbb{P}(u)}{\mathbb{P}(y)}$$

- ▶ defining this on an infinite dimensional Hilbert space the corresponding densities cannot be obtained
- ▶ However the Radon-Nikodym derivative between a posterior measure  $\mu^y$  and a prior measure  $\mu^0$  can define the corresponding *likelihood function*

$$\frac{d\mu^y}{d\mu^0} \propto \mathbb{P}(y|u)$$

- ▶ The reference measure for a Hilbert space is the Gaussian Measure and so the prior measure over functions  $\mu^0 = \mathcal{N}(m, c)$
- ▶ This is formal definition of a Gaussian Process prior - one of the major tools in Machine Learning

Lecture Outline

Lebesgue Measure  
in infinite  
dimensionsMeasures in  
infinite dimensionsGaussian Measures  
in infinite  
dimensionsFull covariance  
operators in  
infinite dimensionsGaussian Measures  
in Hilbert SpaceEquivalence of  
MeasuresEquivalence of  
MeasuresBayes Rule in  
Hilbert Space