

The Hedin's equation

Ref: <http://www.teor.mi.infn.it/~molinari/NOTES/hedin2017.pdf>

In 1965, Lars Hedin derived the formally closed set of equations for propagator G , self-energy Σ_0 , effective potential W , proper polarization Π_0 and dressed vertex Γ .

$$G(1, 2) = G_0(1, 2) + \int d1'2' G_0(1, 1') \Sigma_0(1', 2') G(2', 2) \quad (1)$$

$$W(1, 2) = V(1, 2) + \int d1'2' V(1, 1') \Pi_0(1', 2') W(2', 2) \quad (2)$$

$$\Sigma_0(1, 2) = i \int d34 W(1, 3) \Gamma(3; 4, 2) G(1, 4) \quad (3)$$

$$\Pi_0(1, 2) = -i \int d34 \Gamma(1; 3, 4) G(2, 4) G(4, 2) \quad (4)$$

$$\Gamma(1; 2, 3) = \delta(1, 2) \delta(1, 3) + \int d4567 \Gamma(1; 4, 5) G(6, 4) G(5, 7) \frac{\partial \Sigma_0(2, 3)}{\partial G(6, 7)} \quad (5)$$

Here, $1 = (x_1, t_1)$, $V(1, 2) = V_0(x_1, x_2) \delta(t_1 - t_2)$ and G_0 is Green function with Hartree approximation ("tadpole" diagrams with **exact** particle density).

In Landau's book, the self energy is decoupled to two parts. One is "branched" diagrams Σ_a , which is absorbed into G_0 , and another is Σ_b , which is Σ_0 in our notation.

就是说我们从决定 Σ (粒子间是成对相互作用) 的全部图形集合中分离出各种“分枝”图, 这些“分枝”图是用一条虚线连接到各外线上的: 它们之和以 Σ_a 标记. 所有这些图形都包含在如下形状的一个骨架图形之中①:

Diagram (14.8) shows a self-energy diagram (a circle with a dashed line branch) equal to a diagram with a dashed line branch (a circle with a dashed line branch). The equation is labeled (14.8).

Σ 的其余部分用 Σ_b 标记. 这样, 在一级和二级图形中, 属于第一种的图形如下:

Diagram (14.9) shows three diagrams (a), (b), and (c) representing the first part of the self-energy. Diagram (a) is a circle with a dashed line branch. Diagram (b) is a circle with a dashed line branch and a dashed line loop. Diagram (c) is a circle with a dashed line branch and a dashed line loop. The equation is labeled (14.9).

EQU1 and EQU2 are just Dyson's equation. EQU3 and EQU4 gives the skeleton diagram, and EQU5 contains a functional derivative, which is the main difficulty of many-body systems.

We now derive it from a field perspective. We assume that EQU1 is already proved.

Dyson-Schwinger equation

Consider adding a source term ϕ to the Hamiltonian:

$$H(t) = H_0 + U + \int d\vec{x} \phi(\vec{x}, t) n(\vec{x})$$

Where n is density operator. We will take $\phi = 0$ in the results. The Dyson-Schwinger equation show that

$$\frac{\delta}{\delta \phi(x)} \frac{\langle \Omega | \mathcal{T} S F | \Omega \rangle}{\langle \Omega | S | \Omega \rangle} = -i \frac{\langle \Omega | \mathcal{T} S F \delta n(x) | \Omega \rangle}{\langle \Omega | S | \Omega \rangle}$$

- F is product of field operators
- $\delta n = n - \langle n \rangle$
- Ω is ground state of $H_I = H_0 + U$
- $S = \mathcal{T} \exp \frac{1}{i\hbar} \int d^4x \phi(x) n(x)$ is the evolution operator
- $n(x) = n(\vec{x}, t) = e^{iH_I t} n(\vec{x}) e^{-iH_I t}$.
- $\langle n(x) \rangle = \frac{\langle \Omega | \mathcal{T} S n(x) | \Omega \rangle}{\langle \Omega | S | \Omega \rangle}$

This equation is easily verified by

$$\frac{\delta \mathcal{T}[SF]}{\delta \phi(x)} = \sum_{k=1}^{\infty} \frac{(-i)^k k}{k!} \int d1' \dots k' n(1') \dots n(k') \delta(1 - 1') \phi(2') \dots \phi(k') = -i \mathcal{T}[SF n(x)]$$

and the definition of $\langle n \rangle$.

Derivation of Hedin's equation

The first equation is easiest to verify: because $G = g + g(\Sigma_H + \Sigma_0)G$ and $G_0 = g + g\Sigma_0 G_0$, then $G = G_0 + G_0 \Sigma_0 G$. Σ_H is contribution from Hartree terms and g is Green function with no interaction.

We define the time-ordered propagator and full polarization as

$$G(1, 2) = -i \frac{\langle \Omega | \mathcal{T} S \psi(1) \psi^\dagger(2) | \Omega \rangle}{\langle \Omega | S | \Omega \rangle}$$

$$\Pi(1, 2) = -i \frac{\langle \Omega | \mathcal{T} S \delta n(1) \delta n(2) | \Omega \rangle}{\langle \Omega | S | \Omega \rangle}$$

Choosing $F = n$ in DS equation, we have

$$\frac{\delta}{\delta \phi(2)} \frac{\langle \Omega | \mathcal{T} S n(1) | \Omega \rangle}{\langle \Omega | S | \Omega \rangle} = -i \frac{\langle \Omega | \mathcal{T} S n(1) \delta n(2) | \Omega \rangle}{\langle \Omega | S | \Omega \rangle} = -i \frac{\langle \Omega | \mathcal{T} S \delta n(1) \delta n(2) | \Omega \rangle}{\langle \Omega | S | \Omega \rangle}$$

We express it in Green's function: because $\langle n(1) \rangle = -iG(1, 1^+)$, then

$$\Pi(1, 2) = -i \frac{\delta G(1, 1^+)}{\delta \phi(2)}$$

We define a "modified" local potential by adding in Hartree term:

$$v(1) = \phi(1) - i \int d2 V(1, 2) G(2, 2^+)$$

Then

$$\frac{\delta}{\delta\phi(1)} = \int d2 \frac{\delta v(2)}{\delta\phi(1)} \frac{\delta}{\delta v(2)} = \frac{\delta}{\delta v(1)} + \int d23 V(2,3) \Pi(3,1) \frac{\delta}{\delta v(2)}$$

Impose this equation on $-iG$ and defining the proper polarization as $\Pi_0(1,2) = -i \frac{\delta G(1,1^+)}{\delta v(2)}$, we have the Dyson's equation for Π :

$$\Pi(1,2) = \Pi_0(1,2) + \int d34 \Pi_0(1,3) V(3,4) \Pi(4,2)$$

Define the two-body effective potential as

$$\begin{aligned} W(1,3) &= \int d2 \frac{\delta v(1)}{\delta\phi(2)} V(2,3) \\ &= V(1,3) + \int d24 V(1,4) \Pi(4,2) V(2,3) \\ &= V(1,3) + \int d24 V(1,4) \Pi_0(4,2) W(2,3) \end{aligned}$$

In the last equation, we used $\int d1 V(4,1) \frac{\delta}{\delta\phi(1)} = \int d1 W(4,1) \frac{\delta}{\delta v(1)}$. **This is the second Hedin's equation.**

We now define the inverse Green function

$$\int d3 G^{-1}(1,3) G(3,2) = \delta(1,2)$$

The derivative of $v(4)$ gives

$$\begin{aligned} 0 &= \int d3 \frac{\delta G^{-1}(1,3)}{\delta v(4)} G(3,2) + \int d3 G^{-1}(1,3) \frac{\delta G(3,2)}{\delta v(4)} \\ &= \int d13 G(5,1) \frac{\delta G^{-1}(1,3)}{\delta v(4)} G(3,2) + \frac{\delta G(5,2)}{\delta v(4)} \end{aligned}$$

In the second equation, we inserted $\int d1 G(5,1)$. Define vertex as

$$\Gamma(1;2,3) = - \frac{\delta G^{-1}(2,3)}{\delta\phi(1)}$$

Then let $2 = 5^+$:

$$\Pi_0(1,2) = -i \int d34 \Gamma(1;3,4) G(3,2) G(4,2)$$

This is the fourth Hedin's equation.

Now because $G^{-1}(1,2) = G_0^{-1}(1,2) - \Sigma_0(1,2)$, and the equation of motion for G_0 is

$$(i\partial_{t_1} - h(1) - \phi(1) - \int d2 V(1,2) n(2)) G_0(1,2) = \delta(1,2)$$

Then $G_0^{-1}(1,2) = i\partial_{t_1} - h(1) - v(1)$, and

$$\Gamma(1;2,3) = \delta(1,2)\delta(1,3) + \frac{\partial \Sigma_0(2,3)}{\partial v(1)}$$

Because of Hohenberg-Kohn theorem, G is uniquely determined by V , then

$$\frac{\delta}{\delta v(5)} = \int d12 \frac{\delta G(1,2)}{\delta v(5)} \frac{\delta}{\delta G(1,2)} = \int d1234 G(1,4) \Gamma(5; 4, 3) G(3, 2) \frac{\delta}{\delta G(1, 2)}$$

Then

$$\Gamma(1; 2, 3) = \delta(1, 2)\delta(1, 3) + \int d4567 \Gamma(1; 4, 5) G(6, 4) G(5, 7) \frac{\partial \Sigma_0(2, 3)}{\partial G(6, 7)}$$

This is the fifth Hedin's equation.

Lastly, we consider the equation of motion for G . By expanding the time-derivative of the expression of G , we obtain

$$\begin{aligned} (i\partial_{t_1} - h(1) - v(1))G(1, 2) &= \delta(1, 2) + \int d3 V(1, 3) \frac{\langle \Omega | \mathcal{T} S \psi(1) \psi^\dagger(2) \delta n(3) | \Omega \rangle}{\langle \Omega | S | \Omega \rangle} \\ &= \delta(1, 2) + i \int d3 V(1, 3) \frac{\delta G(1, 2)}{\delta \phi(3)} \end{aligned}$$

Let $G = G_0 + G_0 \Sigma_0 G$, we have

$$\int d3 \Sigma_0(1, 3) G(3, 2) = i \int d3 V(1, 3) \frac{\delta G(1, 2)}{\delta \phi(3)} = i \int d3 W(1, 3) \frac{\delta G(1, 2)}{\delta v(3)}$$


Inserting $\int d2 G^{-1}(2, 4)$:

$$\begin{aligned} \Sigma_0(1, 2) &= i \int d34 W(1, 3) \frac{\delta G(1, 4)}{\delta v(3)} G^{-1}(4, 2) \\ &= -i \int d34 W(1, 3) G(1, 4) \frac{\delta G^{-1}(4, 2)}{\delta v(3)} \\ &= i \int d34 W(1, 3) \Gamma(3; 4, 2) G(1, 4) \end{aligned}$$


This is the third Hedin's equation.

Diagrammatic expression of Hedin's equation

EQU1 and EQU2:



EQU3 and EQU4 (the left is Σ_0 and the right is Π_0):




GW approximation

The second term in (5) can be neglected in GW, then

$$\begin{aligned}\Gamma(1; 2, 3) &= \delta(1, 2)\delta(1, 3) \\ \Sigma_0(1, 2) &= iG(1, 2)W(1, 2) \\ \Pi_0(1, 2) &= -iG(1, 2)G(2, 1)\end{aligned}$$

And W is determined by Π_0 . This is like the RPA approximation:



(The first term in the first equation above is absorbed into G_0).

Figures from <https://www.cond-mat.de/events/correl11/manuscripts/held.pdf>