Canonical Physics and Conventions Reference for `cqed_sim`

Post-cleanup and protocol-alignment reference

March 10, 2026

This document is the source of truth for the reusable simulator code under `cqed_sim/`. It records the active physics conventions, the protocol-facing abstractions that now sit on top of the simulator, the cleaned package boundary versus `examples/`, and the tests that directly back the main conventions.

1 Overview of cqed_sim

`cqed_sim/` now contains only reusable simulator code:

- `core`: Hilbert-space conventions, frames, the two-mode dispersive transmon-storage model, the three-mode storage-transmon-readout model, ideal gates, and manifold-frequency helpers.
- `pulses`, `sequence`, `sim`: pulse objects, envelopes, channel compilation, hardware distortion, the solver path, collapse operators, and extractors.
- `experiment`: reusable state preparation, qubit measurement, and a lightweight protocol wrapper built on the same compile-and-simulate path as the low-level runtime.
- `calibration`, `observables`, `operators`, `tomo`, `io`, `plotting`, `unitary_synthesis`: reusable helpers that still belong to the library surface.

Study, workflow, paper-reproduction, audit, and smoke-test code remains under `examples/`. The alignment note for the experiment-facing cleanup is kept in `physics_and_conventions/experimental_protocol_a`

2 Experimental Protocol Layer

The main experiment-facing additions are implemented in:

- `cqed_sim/experiment/state_prep.py`: `StatePreparationSpec` and reusable subsystem state specs.
- `cqed_sim/experiment/measurement.py`: `QubitMeasurementSpec` and `measure_qubit(...)`.
- `cqed_sim/experiment/readout_chain.py`: `ReadoutResonator`, `PurcellFilter`, `AmplifierChain`, and `ReadoutChain`.
- `cqed_sim/experiment/protocol.py`: `SimulationExperiment`, `ExperimentResult`, and metadata wrappers.

These objects are not a second solver stack. They are a thin reusable layer above `SequenceCompiler` plus `simulate_sequence(...)`. In particular:

- `SimulationExperiment` is executable, not merely declarative. It resolves an initial state, compiles pulses, runs the existing solver path, and optionally performs qubit measurement on the resulting runtime state.
- State preparation may return either a ket or a density matrix. If any subsystem spec is mixed, the joint initial state is promoted to a density matrix before tensoring.
- `measure_qubit(...)` returns exact probabilities by default. Sampled outcomes appear only when `shots` is set.
- When a confusion matrix is supplied, the code applies $\mathbf{p}_{\text{obs}} = M\mathbf{p}_{\text{latent}}$ with columns ordered as latent (g, e) and rows ordered as reported (g, e).
- If `QubitMeasurementSpec.readout_chain` is omitted, synthetic I/Q points are still generated from the previous Gaussian cluster model.
- If a `ReadoutChain` is provided, the measurement layer instead uses a resonator response, optional Purcell filter, amplifier noise, and mixer-style I/Q integration. The same API can also report γ_{meas} and a Purcell-limited T_1, and can optionally apply readout-induced dephasing plus Purcell relaxation as a reduced-state backaction model.

Direct tests: `tests/test_26_experimental_protocols.py`.

3 Physical Readout Chain

The physical readout-chain path models a driven dispersive resonator with optional Purcell filtering and additive amplifier noise:

- `ReadoutResonator` stores ω_r, κ, the qubit-resonator coupling g, the complex drive amplitude ϵ, and an optional runtime-convention dispersive pull χ.
- The steady-state pointer amplitudes are

$$\alpha_{g/e}^{(\text{ss})} = \frac{-i\epsilon}{\kappa_{\text{eff}}/2 + i(\omega_{r,g/e} - \omega_d)},$$

  with $\omega_{r,g} = \omega_r$ and $\omega_{r,e} = \omega_r - \chi$ in the runtime projector convention.
- The implemented measurement-induced dephasing rate is

$$\gamma_{\text{meas}} = \frac{\kappa_{\text{eff}}}{2}|\alpha_e - \alpha_g|^2,$$

  which reproduces the expected small-χ scaling $\gamma_{\text{meas}} \propto \chi^2\bar{n}/\kappa$.
- The Purcell filter follows the standard frequency-dependent linewidth picture:

$$\kappa_{\text{eff}}(\omega) = \frac{4J^2\kappa_f}{\kappa_f^2 + 4(\omega - \omega_f)^2},$$

  so the Purcell rate is evaluated as $\Gamma_P = (g/\Delta)^2\kappa_{\text{eff}}(\omega_q)$.
- `AmplifierChain` applies linear gain and additive thermal noise before mixer-phase rotation and I/Q integration.

Direct tests: `tests/experiment/test_readout_chain.py`.

4 Hilbert Space Structure

Two-mode runtime

- Runtime ordering is qubit first, storage second:

$$|q, n\rangle = |q\rangle_{\text{qubit}} \otimes |n\rangle_{\text{storage}}.$$

- The flattened basis index is $\texttt{index}(q, n) = q\, n_{\text{cav}} + n$, implemented in `cqed_sim/core/conventions.py`.
- Computational basis is $|g\rangle = |0\rangle$, $|e\rangle = |1\rangle$.

Three-mode runtime

- The three-mode extension uses

$$|q, n_s, n_r\rangle = |q\rangle_{\text{qubit}} \otimes |n_s\rangle_{\text{storage}} \otimes |n_r\rangle_{\text{readout}}.$$

- The flattened basis index is

$$\texttt{index}(q, n_s, n_r) = ((q\, n_{\text{storage}}) + n_s)\, n_{\text{readout}} + n_r.$$

- This ordering is implemented in `cqed_sim/core/conventions.py` and `cqed_sim/core/readout_model.py`, and all new extractors assume subsystem 0 is the transmon, subsystem 1 is storage, subsystem 2 is readout.

Synthesis subspace

The unitary-synthesis layer retains the same full-space qubit-first ordering, but restricted subspace bases are explicitly reordered in `cqed_sim/unitary_synthesis/subspace.py`; for example, a qubit-cavity block is ordered as $|g, 0\rangle, |e, 0\rangle, |g, 1\rangle, |e, 1\rangle, \ldots$.

Direct tests: `tests/test_25_tensor_product_convention.py`, `tests/test_27_three_mode_model.py`, `tests/unitary_synthesis/test_subspace.py`.

5 Operator Conventions

- The simulator uses QuTiP Pauli matrices without sign changes. Therefore

$$\sigma_z|g\rangle = +|g\rangle, \qquad \sigma_z|e\rangle = -|e\rangle.$$

- The core Hamiltonians use the excitation number operator $n_q = b^\dagger b$, not $\sigma_z/2$. For two levels,

$$n_q = |e\rangle\langle e| = \frac{I - \sigma_z}{2}.$$

- Two-mode operators in `cqed_sim/core/model.py` are embedded as $a = I_q \otimes a_{\text{storage}}$ and $b = b_{\text{transmon}} \otimes I_s$.
- Three-mode operators in `cqed_sim/core/readout_model.py` are embedded as

$$b = b_{\text{transmon}} \otimes I_s \otimes I_r, \quad a_s = I_q \otimes a_{\text{storage}} \otimes I_r, \quad a_r = I_q \otimes I_s \otimes a_{\text{readout}}.$$

- The generic runtime now resolves drive operators through `model.drive_coupling_operators()`, so the solver no longer hard-codes a qubit-plus-one-mode assumption.

Direct tests: `tests/test_16_ideal_primitives_and_extractors.py`, `tests/test_20_gaussian_iq_convention` `tests/test_25_tensor_product_convention.py`, `tests/test_27_three_mode_model.py`.

6 Hamiltonian Definitions

Two-mode runtime Hamiltonian

The reusable two-mode Hamiltonian in `cqed_sim/core/model.py` is

$$H_0 = \delta_c n_c + \delta_q n_q + \frac{\alpha}{2} b^{\dagger 2} b^2 + K(n_c) - \Xi(n_c) n_q,$$

with

$$K(n_c) = \frac{\texttt{kerr}}{2} n_c(n_c - 1) + \cdots, \qquad \Xi(n_c) = \texttt{chi}\, n_c + \texttt{chi2}\, n_c(n_c-1) + \texttt{chi3}\, n_c(n_c-1)(n_c-2) + \cdots.$$

`cqed_sim/core/frequencies.py` makes the intended transition law explicit:

$$\omega_{ge}(n) = \omega_{ge}(0) - \Xi(n).$$

Textbook-to-code translation

With the active runtime sign convention,

$$n_q = \frac{I - \sigma_z}{2}.$$

Therefore

$$\delta_q n_q = \frac{\delta_q}{2} I - \frac{\delta_q}{2} \sigma_z, \qquad -\Xi(n_c) n_q = -\frac{\Xi(n_c)}{2} I + \frac{\Xi(n_c)}{2} \sigma_z.$$

Up to identity and storage-only shifts,

$$H_{q,\text{runtime}} \doteq -\frac{1}{2}\big(\delta_q - \Xi(n_c)\big)\sigma_z.$$

If one instead uses the textbook sign

$$\sigma_z^{\text{text}} = |e\rangle\langle e| - |g\rangle\langle g| = -\sigma_z,$$

then

$$H_{q,\text{runtime}} \doteq \frac{1}{2}\big(\delta_q - \Xi(n_c)\big)\sigma_z^{\text{text}}.$$

Hence runtime `chi` is the per-photon downward pull of the $|g, n\rangle \leftrightarrow |e, n\rangle$ transition frequency.

Comparison with the synthesis drift convention

`cqed_sim/unitary_synthesis/sequence.py` defines drift energies

$$E_{g,n} = \delta_c n - \frac{\delta_q}{2} - \Xi_{\text{synth}}(n) + K(n), \qquad E_{e,n} = \delta_c n + \frac{\delta_q}{2} + \Xi_{\text{synth}}(n) + K(n).$$

Its docstring uses the opposite qubit sign convention,

$$\sigma_z^{\text{synth}}|e\rangle = +|e\rangle, \qquad \sigma_z^{\text{synth}}|g\rangle = -|g\rangle.$$

Its transition shift is therefore

$$E_{e,n} - E_{g,n} = \delta_q + 2\Xi_{\text{synth}}(n).$$

Matching the runtime shift $\delta_q - \Xi_{\text{runtime}}(n)$ gives

$$2\Xi_{\text{synth}}(n) = -\Xi_{\text{runtime}}(n),$$

so componentwise

$$\chi_{\text{synth}} = -\chi_{\text{runtime}}/2, \qquad \chi_{2,\text{synth}} = -\chi_{2,\text{runtime}}/2, \qquad \chi_{3,\text{synth}} = -\chi_{3,\text{runtime}}/2.$$

This mapping is derived from code and tests, but is not enforced by a dedicated helper.

Three-mode storage-transmon-readout Hamiltonian

The new three-mode model in `cqed_sim/core/readout_model.py` implements

$$\frac{H}{\hbar} = \delta_s n_s + \delta_r n_r + \delta_q n_q + \frac{\alpha}{2} b^{\dagger 2} b^2 - \chi_s n_s n_q - \chi_r n_r n_q + \chi_{sr} n_s n_r + \frac{K_s}{2} n_s(n_s - 1) + \frac{K_r}{2} n_r(n_r - 1).$$

This is intentionally kept in the same runtime convention style as the two-mode path:

- $n_q = b^\dagger b$ is the transmon excitation projector/number operator.
- Positive χ_s or χ_r lowers the transmon transition frequency as storage or readout occupancy increases.
- Positive χ_{sr} raises the storage frequency with readout occupancy and raises the readout frequency with storage occupancy.
- Storage and readout self-Kerr use the same $K\, n(n-1)/2$ convention as the two-mode storage Kerr term.

The model also exposes analytic helpers for the basis energies and the qubit, storage, and readout transition frequencies.

Optional nonlinear and exchange terms

`cqed_sim/sim/couplings.py` now provides explicit builder functions for

$$\chi_{ab} a^\dagger a\, b^\dagger b, \qquad \frac{K}{2} a^{\dagger 2} a^2, \qquad J(a^\dagger b + a b^\dagger),$$

and the model classes accept `CrossKerrSpec`, `SelfKerrSpec`, and `ExchangeSpec` entries that are injected into the static Hamiltonian assembly. This extends the existing dispersive models without replacing their original API or sign conventions.

Canonical parameter-definition table

Name	Code-level meaning	Algebra used	Caveat
`chi`	two-mode first-order dispersive pull	$-\texttt{chi}\, n_c n_q$, $\omega_{ge}(n) = \omega_{ge}(0) - \chi n$	runtime projector form, not stored as $\sigma_z a^\dagger a/2$
`chi2`	second-order pull, exposed by `chi_higher[0]`	$-\chi_2 n_c(n_c - 1) n_q$	falling-factorial coefficient
`chi3`	third-order pull, exposed by `chi_higher[1]`	$-\chi_3 n_c(n_c - 1)(n_c - 2) n_q$	falling-factorial coefficient
`chi_s`	storage-transmon dispersive pull in the three-mode model	$-\chi_s n_s n_q$	positive values lower ω_{ge} with storage photons
`chi_r`	readout-transmon dispersive pull in the three-mode model	$-\chi_r n_r n_q$	positive values lower ω_{ge} with readout photons

Name	Code-level meaning	Algebra used	Caveat
`chi_sr`	storage-readout cross-Kerr	$+\chi_{sr} n_s n_r$	positive values raise both bosonic mode frequencies with occupancy of the other mode
`kerr`	two-mode storage self-Kerr coefficient	`kerr` $n_c(n_c-1)/2$	same sign convention as code, no implicit factor change
higher Kerr	tuple `kerr_higher`	$K_m n_c^{\underline{m}}/m!$	falling-factorial hierarchy
`kerr_s`	three-mode storage self-Kerr	`kerr_s` $n_s(n_s-1)/2$	same convention as `kerr`
`kerr_r`	three-mode readout self-Kerr	`kerr_r` $n_r(n_r-1)/2$	same convention as `kerr`
`alpha`	transmon Duffing anharmonicity	`alpha` $b^{\dagger 2} b^2/2$	inactive for strict two-level truncation
σ_z	active runtime Pauli-Z	$\lvert g\rangle\langle g\rvert - \lvert e\rangle\langle e\rvert = I - 2n_q$	opposite sign from the synthesis docstring
n_q	transmon excitation operator	$b^\dagger b$; for two levels $n_q = \lvert e\rangle\langle e\rvert$	repo-specific implementation choice versus $\sigma_z/2$ notation
δ_q	qubit residual detuning	$\omega_q - \omega_q^{\text{frame}}$	only defined after choosing a frame
δ_c	two-mode storage residual detuning	$\omega_c - \omega_c^{\text{frame}}$	same frame caveat
δ_s	three-mode storage residual detuning	$\omega_s - \omega_s^{\text{frame}}$	ω_s^{frame} is stored as `omega_c_frame` with an alias
δ_r	three-mode readout residual detuning	$\omega_r - \omega_r^{\text{frame}}$	only used in the three-mode path

Standard-compatible versus repo-specific

- Standard-compatible: qubit-first tensor ordering, QuTiP Pauli signs, rad/s and seconds internally, Duffing/Kerr terms, Lindblad dynamics, and positive χ or χ_s, χ_r as downward transmon pulls.
- Repo-specific: writing the transmon term in terms of n_q, storing higher-order pulls and Kerr terms in falling-factorial form, and using complex envelopes with $e^{+i(\omega t+\phi)}$.
- Needs caution: the synthesis drift model still does not share the runtime σ_z sign or χ normalization.

7 Frequency and Time Units

- Internal Hamiltonian coefficients and rotating-frame frequencies are in rad/s.
- Pulse durations, timeline steps, and solver `max_step` values are in seconds.
- The repository now centralizes conversion helpers in `physics_and_conventions/conventions.py`, including `to_internal_units(...)` and `from_internal_units(...)`.

- Example-side configuration files may still carry `_hz`, `_ns`, or `_1_per_s` field names, but those values are converted before constructing library models or noise objects.

Direct tests: `tests/test_05_detuning_and_frames.py`, `tests/test_10_chi_convention.py`, `tests/test_13_ha` `tests/test_27_three_mode_model.py`.

8 Parameter Translation, Calibration Targets, and Backends

- `cqed_sim/analysis/parameter_translation.py` now provides `from_transmon_params(...)` and `from_measured(...)`. Bare transmon parameters use the usual large-E_J/E_C expansion $\omega_{01} \approx \sqrt{8E_JE_C} - E_C$ and $\alpha \approx -E_C$, while dressed χ and χ_2 are extracted by matching a low-dimensional Duffing-plus-resonator diagonalization to the runtime form $\omega_{ge}(n) = \omega_q - \chi n - \chi_2 n(n-1)$.
- `cqed_sim/calibration_targets/` now provides reusable spectroscopy, Rabi, Ramsey, T_1, $T_{2,\mathrm{echo}}$, and DRAG-tuning surrogate targets that return fitted parameters, uncertainties, and raw simulated data.
- `cqed_sim/backends/` now provides `NumPyBackend` and `JaxBackend` plus a dense piecewise-constant solver path in `cqed_sim/sim/solver.py`. The default runtime still uses QuTiP; the backend path is opt-in through `SimulationConfig.backend`.

Direct tests: `tests/analysis/test_parameter_translation.py`, `tests/calibration_targets/test_targets.p` `tests/sim/test_backends.py`.

9 Rotating Frame Conventions

- Frames are represented by `FrameSpec(omega_c_frame, omega_q_frame, omega_r_frame)` in `cqed_sim/core/frame.py`.
- The legacy field `omega_c_frame` remains the storage/cavity frame frequency. The readout model adds the alias property `omega_s_frame` for clarity.
- The two-mode path uses $\delta_c = \omega_c - \omega_c^{\mathrm{frame}}$ and $\delta_q = \omega_q - \omega_q^{\mathrm{frame}}$.
- The three-mode path uses $\delta_s = \omega_s - \omega_s^{\mathrm{frame}}$, $\delta_r = \omega_r - \omega_r^{\mathrm{frame}}$, and $\delta_q = \omega_q - \omega_q^{\mathrm{frame}}$.
- `manifold_transition_frequency(...)` still refers only to the two-mode path; the three-mode model instead exposes `qubit_transition_frequency(...)`, `storage_transition_frequency(...)`, and `readout_transition_frequency(...)`.

Direct tests: `tests/test_05_detuning_and_frames.py`, `tests/test_21_qubox_convention_reconciliation.py` `tests/test_27_three_mode_model.py`.

10 Drive Hamiltonians

- `cqed_sim/pulses/pulse.py` samples analytic or discrete envelopes as

$$\epsilon(t) = \texttt{amp}\,\texttt{envelope}(t)\, e^{i(\texttt{carrier}t+\texttt{phase})}.$$

- `cqed_sim/sim/runner.py` inserts $\epsilon(t)$ and $\epsilon^*(t)$ as raising and lowering couplings obtained from the model, rather than from a hard-coded target list.
- The two-mode path supports `"cavity"` or `"storage"`, `"qubit"`, and `"sideband"` targets.
- The three-mode path supports `"storage"`, `"readout"`, `"qubit"`, and `"transmon"` targets. The readout and storage drives use the same complex-envelope sign convention as the existing qubit and storage drives.
- The effective three-mode drive Hamiltonian is

$$H_d/\hbar = \epsilon_s(t)a_s^\dagger + \epsilon_s^*(t)a_s + \epsilon_q(t)b^\dagger + \epsilon_q^*(t)b + \epsilon_r(t)a_r^\dagger + \epsilon_r^*(t)a_r.$$

Direct tests: `tests/test_20_gaussian_iq_convention.py`, `tests/test_23_sqr_additive_amplitude_correcti` `tests/test_28_three_mode_drives_and_noise.py`.

11 Dissipation and Lindblad Terms

- `NoiseSpec` in `cqed_sim/sim/noise.py` now supports `t1`, `tphi`, `kappa`, `nth`, `kappa_storage`, `kappa_readout`, `nth_storage`, and `nth_readout`.
- The implemented convention is $\gamma_1 = 1/T_1$ and $\gamma_\phi = 1/(2T_\phi)$.
- Two-level dephasing uses $\sqrt{\gamma_\phi}\,\sigma_z$ with $\sigma_z = I - 2n_q$. Multi-level transmon dephasing uses $\sqrt{\gamma_\phi}\,n_q$.
- In the two-mode path, `kappa` and `nth` still drive storage loss and thermal excitation.
- In the three-mode path, storage and readout can decay independently via `kappa_storage` and `kappa_readout`, with separate thermal occupations.
- `simulate_sequence(...)` uses `mesolve` when collapse operators are present or when the initial state is already a density matrix; otherwise it uses `sesolve`.

Direct tests: `tests/test_14_dissipation.py`, `tests/test_18_noise_and_sideband.py`, `tests/test_28_three_`

12 State Representation

- Runtime inputs may be kets or density matrices.
- The result wrapper stores the final state, optional state history, expectation traces, and the raw QuTiP solver result.
- The generic extractor path now supports two-mode and three-mode states through subsystem-aware partial traces.
- `reduced_cavity_state(...)` remains specific to the two-mode path. The three-mode path uses `reduced_storage_state(...)` and `reduced_readout_state(...)`.
- `measure_qubit(...)` acts on the reduced transmon state obtained from the runtime state convention. The exact probabilities are taken from the reduced density matrix before any optional confusion-matrix application or finite-shot sampling.

- Synthetic I/Q points remain available for lightweight sampling-only usage.
- The new readout-chain path adds optional state-conditioned resonator traces, amplifier-noise I/Q sampling, and a reduced-state backaction model without changing the default no-chain behavior.

Direct tests: `tests/test_16_ideal_primitives_and_extractors.py`, `tests/test_19_fock_tomo.py`, `tests/test_26_experimental_protocols.py`, `tests/test_27_three_mode_model.py`, `tests/test_28_three_mo`

13 Hilbert Space Truncation

- The two-mode model uses `n_cav` for storage truncation and `n_tr` for transmon truncation.
- The three-mode model uses `n_storage`, `n_readout`, and `n_tr`.
- State preparation helpers pad or truncate explicit amplitude vectors to the requested subsystem dimension.
- SQR angle arrays are still padded to the storage dimension in `cqed_sim/pulses/calibration.py`.
- The synthesis layer still distinguishes the full Hilbert dimension from restricted subspace dimension in `cqed_sim/unitary_synthesis/subspace.py`.

Direct tests: `tests/test_11_model_invariants.py`, `tests/test_24_displacement_selective_spectroscopy.p`, `tests/test_26_experimental_protocols.py`, `tests/test_27_three_mode_model.py`, `tests/unitary_synthesi`

14 Test Suite Audit

File	Targets, type, and main assertion	Status
`tests/test_01_sanity_and_free.py`	unit and numerical sanity checks for Hermiticity, free evolution, and Kerr phase accumulation in the two-mode runtime.	Passed
`tests/test_02_cavity_drive_and_kerr.py`	unit and regression checks for storage drive behavior and Kerr response.	Passed
`tests/test_03_dispersive_and_ramsey.py`	dispersive phase scaling and Ramsey pull in the two-mode model.	Passed
`tests/test_04_xy_phase_and_overlap.py`	XY phase convention and overlap cancellation for qubit pulses.	Passed
`tests/test_05_detuning_and_frames.py`	rotating-frame detuning signs and frame handling.	Passed
`tests/test_06_leakage_drag_and_higher_order.py`	multilevel leakage, DRAG behavior, and higher-order term smoke checks.	Passed
`tests/test_07_convergence_regression.py`	truncation convergence and regression against a stored reference.	Passed
`tests/test_08_timeline_and_hardware.py`	waveform timeline assembly and basic hardware distortion semantics.	Passed
`tests/test_09_runtime_policy.py`	runtime-budget and fast-path regression checks.	Passed
`tests/test_10_chi_convention.py`	direct checks of runtime χ sign, transition-pull meaning, and observable defaults.	Passed
`tests/test_11_model_invariants.py`	model dimensions, invariants, and expectation consistency.	Passed
`tests/test_12_pulse_semantics.py`	pulse-area and pulse-shape semantics.	Passed

File	Targets, type, and main assertion	Status
`tests/test_13_hardware_extended.py`	ZOH, low-pass, sideband, quantization, and crosstalk behavior.	Passed
`tests/test_14_dissipation.py`	explicit T_1, T_ϕ, and storage damping checks using collapse operators.	Passed
`tests/test_15_numerics_performance.py`	runtime, compile-cache, and runtime scaling smoke checks.	Passed
`tests/test_16_ideal_primitives_and_extractors.py`	ideal SNAP/SQR primitives and the two-mode extractor path.	Passed
`tests/test_17_gate_library_integration.py`	registration and integration of the library-level gate path.	Passed
`tests/test_18_noise_and_sideband.py`	sideband exchange and combined noise behavior in the two-mode runtime.	Passed
`tests/test_19_fock_tomo.py`	reusable tomography and leakage-calibration helpers.	Passed
`tests/test_20_gaussian_iq_convention.py`	Gaussian IQ axis, phase, and detuning sign convention.	Passed
`tests/test_21_qubox_conventions.py`	reconciliation between runtime conventions and externally documented qubox-style conventions.	Passed
`tests/test_22_support_aware_objective.py`	support-aware objective wiring in synthesis-related helpers.	Passed
`tests/test_23_sqr_additive_amplitude_correction.py`	additive correction convention for SQR tones.	Passed
`tests/test_24_displacement_selective_spectroscopy.py`	selective-spectroscopy frequencies and displacement conventions.	Passed
`tests/test_25_tensor_product_conventions.py`	conventions of tensor ordering, block indices, and active σ_z sign.	Passed
`tests/test_26_experimental_protocols.py`	protocol-layer tests for state preparation ordering, mixed initial states, sampled qubit measurement, and `SimulationExperiment`.	Passed
`tests/test_27_three_mode_model.py`	explicit basis-index, operator-embedding, partial-trace, analytic diagonal-energy, two-mode reduction, and dispersive shift checks for the three-mode model.	Passed
`tests/test_28_three_mode_drives_and_noise.py`	drives and noise, storage/readout ringdown, transmon T_1, transmon dephasing, positivity, and readout-response checks for the three-mode path.	Passed
`tests/unitary_synthesis/test_subspace.py`	subspace basis ordering and round-trip embedding.	Passed
`tests/unitary_synthesis/test_metrics.py`	fidelity, gauges and leakage metrics.	Passed
`tests/unitary_synthesis/test_primitives_and_backends.py`	primitives and backends, duration/pulse backends.	Passed
`tests/unitary_synthesis/test_optimization.py`	optimization and time optimization.	Passed
`tests/unitary_synthesis/test_synthesis_and_targets.py`	synthesis and targets, and MPS-target behavior.	Passed with one optional skip
`tests/unitary_synthesis/test_free_evolution_conditional.py`	free-evolution conditional phase formulas.	Passed
`tests/unitary_synthesis/test_hardware_grid_parallelism.py`	hardware, grid, and parallelism and parallel-multistart behavior.	Passed
`tests/unitary_synthesis/test_progress_reporting.py`	progress reporting and serialization.	Passed
`tests/unitary_synthesis/test_reporting_and_determinism.py`	reporting and determinism.	Passed
`tests/unitary_synthesis/test_time_policy_and_phase.py`	time-policy and conditional-phase decomposition.	Passed
`examples/audits/tests/test_experiment_conventions_audit2.py`	experiment conventions for audit 2, tensor ordering, waveform signs, and relative phases.	Passed
`examples/workflows/tests/test_workflow_artifacts.py`	workflow artifact serialization and replay.	Passed
`examples/studies/tests/test_snap_opt_convergence.py`	snap opt convergence regression for the SNAP optimization study.	Passed
`examples/studies/tests/test_snap_opt_achieves_zero_coherent_error_above_the_empirical limit.py`	snap opt achieves zero coherent error above the empirical limit.	Passed
`examples/studies/tests/test_snap_opt_generalization_sweeps.py`	snap opt generalization sweeps, dropout failure, runtime budget, and parameter sweeps.	Passed

File	Targets, type, and main assertion	Status
`examples/studies/tests/test_map_optimizer.py`	map optimizer preserves total frequency content and produced.	Passed
`examples/paper_reproductions/tests/test_PRL-133_convergence_and_runtime.py`	PRL-133 convergence and runtime-budget regression.	Passed
`examples/paper_reproductions/tests/test_PRL-133_cross-frequency_coupling.py`	PRL-133 cross-frequency coupling test.	Passed
`examples/paper_reproductions/tests/test_PRL-133_optimization_limit.py`	PRL-133 optimization-limit regression.	Passed
`examples/paper_reproductions/tests/test_PRL-133-138_components_consistency.py`	PRL-133/138 components consistency.	Passed
`examples/paper_reproductions/tests/test_PRL-133-138_metric_bounds.py`	PRL-133/138 metric bounds and fail-fast validation.	Passed
`examples/paper_reproductions/tests/test_PRL-133-138_metric_regression.py`	PRL-133/138 metric regression.	Passed
`examples/paper_reproductions/tests/test_PRL-133_unitarity_trace.py`	PRL-133 unitarity and trace-preservation checks.	Passed
`examples/paper_reproductions/tests/test_PRL-138_reproduce_heseline.py`	PRL-138 reproduce-heseline smoke test.	Passed
`examples/smoke_tests/tests/test_sanity.py`	Top-level sanity and partial-workflow sanity checks.	Passed
`examples/smoke_tests/tests/test_sqr_SQR_calibration.py`	SQR calibration smoke and guarded-benchmark checks.	Passed
`examples/smoke_tests/tests/test_sqr_speed_limit_analytical.py`	SQR speed-limit analytical equivalence checks.	Passed

15 Test Execution Results

Commands used:

```
E:\Programs\python.exe -m pytest tests -q
E:\Programs\python.exe -m pytest examples/audits/tests examples/workflows/tests \
  examples/studies/tests examples/paper_reproductions/tests \
  examples/smoke_tests/tests -q
```

Observed results:

- `tests/`: 172 collected, 171 passed, 1 skipped, 0 failed.
- `examples/*/tests`: 49 collected, 49 passed, 0 failed.
- The single skip is the optional external MPS-reference check in `tests/unitary_synthesis/test_synthesis_an`
- Warning-only notes: two `numpy.trapz` deprecation warnings in `tests/test_12_pulse_semantics.py` and `tests/test_20_gaussian_iq_convention.py`.

Interpreter note: `E:\Program Files\Python311\python.exe` from `AGENTS.md` is not present in this workspace, so the suite was run with `E:\Programs\python.exe` (Python 3.11.8).

16 Verified Behaviors Backed by Tests

- Two-mode tensor ordering, basis labels, block indexing, and active Pauli/Bloch sign conventions are directly tested.
- The runtime meaning of χ as a downward transition pull is directly tested.
- The textbook-to-runtime projector translation is supported by direct σ_z sign tests and by the runtime χ transition tests, even though the exact algebraic translation is not asserted by a single dedicated unit test.

- The new three-mode tensor ordering, operator embeddings, partial traces, diagonal energies, and analytic dispersive shifts are directly tested.
- The three-mode solver path is directly tested for drive isolation, storage/readout-specific damping, transmon relaxation, transmon dephasing, positivity, and qubit-conditioned readout response.
- State preparation helpers are directly tested for ordering and mixed-state behavior.
- The qubit measurement abstraction is directly tested for exact probabilities, confusion-matrix application, finite-shot sampling, and synthetic I/Q output shape.
- `SimulationExperiment` is directly tested as the recommended light protocol wrapper over the compile-and-run path.
- The moved example workflows still execute correctly in their new locations.

17 Gaps Between Code Conventions and Test Coverage

- The new runtime-to-synthesis translation helper exposes $\chi_{\text{synth}} = -\chi_{\text{runtime}}/2$, but the library still does not force every synthesis-side caller to use that mapping automatically.
- Higher-order χ_2, χ_3, and higher-order Kerr coefficients are covered indirectly, but not with isolated sign-and-factor analytic tests of the same style now used for the three-mode diagonal-energy checks.
- The readout-chain model is physically motivated, but it is still an effective resonator/filter/amplifier abstraction rather than a full microwave network or POVM calibration workflow.
- The backend path is dense and piecewise constant; it is intended for parity and small-system acceleration checks, not as a drop-in replacement for every QuTiP workflow.

18 Potential Ambiguities

Critical: runtime and synthesis drift layers do not use the same σ_z sign or χ scaling; the report derives the mapping, but code does not enforce it.
Moderate: higher-order pull and Kerr parameters are falling-factorial coefficients, not ordinary polynomial coefficients.
Moderate: the runtime writes transmon physics in terms of n_q rather than $\sigma_z/2$, so ground-state energy offsets differ from some textbook presentations.
Moderate: synthetic I/Q output is a convenience sampling layer, not a calibrated hardware-response model.
Low: the explicit complex-envelope sign $e^{+i(\omega t+\phi)}$ is repo-specific and must be preserved when adding pulse families.
Low: `FrameSpec` still stores the storage frame in the legacy field `omega_c_frame`; the `omega_s_frame` alias removes ambiguity but does not rename the underlying field.

19 Summary

`cqed_sim` is now a coherent reusable library for pulse-level cQED simulation with both a two-mode transmon-storage path and a three-mode storage-transmon-readout extension. The authoritative runtime conventions are: qubit-first tensor ordering; $|g\rangle = |0\rangle$, $|e\rangle = |1\rangle$; QuTiP-standard Pauli signs; rad/s internally and seconds for time; complex envelopes with $e^{+i(\omega t+\phi)}$; dispersive physics written in terms of $n_q = b^\dagger b$; and Lindblad collapse operators as implemented in `cqed_sim/sim/noise.py`. The experiment-facing layer now has two measurement modes: a default lightweight exact-probability path and an opt-in physical readout-chain path with resonator response, Purcell filtering, amplifier noise, and reduced-state backaction. The library also now exposes optional nonlinear/exchange Hamiltonian terms, runtime-to-synthesis parameter translation helpers, calibration-target utilities, and an opt-in dense NumPy/JAX backend path. The main caution that remains is that the synthesis layer still does not automatically consume the runtime-to-synthesis χ mapping everywhere by construction.