Errata File for Second Edition

Daniel Sorensen

October 24, 2026

This errata file contains a list of errors, short clarifications and modifications to the manuscript's second edition.

1. Page 7, first sentence below equation (1.5): "This means that there is a 95% probability that this interval contains the true parameter θ".

   This statement is technically incorrect for the following reason. A $100\,(1-\alpha)\,\%$ confidence interval for θ is an interval defined by the lower and upper bounds $(l\,(Y)\,, u\,(Y))$, where Y represents the data, a random variable, such that

   $$\Pr\left[l\,(Y) < \theta < u\,(Y)\,|\theta\right] = 1-\alpha,$$

   where θ is the fixed, unknown parameter and α is the significance level of the test. The probability is computed with respect to the distribution of Y. Once the data are observed and $Y = y$, the expression

   $$\Pr\left[l\,(y) < \theta < u\,(y)\,|\theta\right] = 1-\alpha,$$

   does not make sense since there are no random variables involved within the square brackets. It is therefore technically incorrect to assert that the observed interval $(l\,(y)\,, u\,(y))$ contains θ with probability $(1-\alpha)$. The interval $(l\,(y)\,, u\,(y))$ either contains or does not contain θ. However, if $\alpha = 0.05$, one can be quite confident that the unobserved θ is within the observed $(l\,(y)\,, u\,(y))$ and act accordingly.

   The remaining statements in the paragraph are correct.

2. Page 170, bottom half of the page, bracket missing:
   CHANGE:

   $$\mathrm{Var}\,(\hat{a}-a) = \mathrm{E}_y(\mathrm{Var}(a|y)$$

   TO.

   $$\mathrm{Var}\,(\hat{a}-a) = \mathrm{E}_y\,(\mathrm{Var}(a|y))$$

3. Page 171, the first partial derivative below Eq. (4.62).
   CHANGE:

$$\frac{\partial \ln p\left(a,b|y,b_0,a_0,B,G,R\right)}{\partial b} = -B^{-1}(b-b_0) + XR^{-1}\left(y - Xb - Za\right)$$

   TO:

$$\frac{\partial \ln p\left(a,b|y,b_0,a_0,B,G,R\right)}{\partial b} = -B^{-1}(b-b_0) + X'R^{-1}\left(y - Xb - Za\right)$$

4. Page 172, top.
   CHANGE: "When B^{-1} implying $B = \infty$ ..."
   TO: "When and if $B^{-1} = 0$ implying $B = \infty$ ..."

5. Page 172,
   CHANGE:
   Finally, add and subtract $\hat{\theta}'\left(W'R^{-1}W + \Sigma\right)\hat{\theta}$ and rearrange to yield

$$y'R^{-1}y + \left(\theta - \widehat{\theta}\right)'\left(W'R^{-1}W + \Sigma\right)\left(\theta - \widehat{\theta}\right)' - \widehat{\theta}'\left(W'R^{-1}W + \Sigma\right)\widehat{\theta} + \theta_0'\Sigma\theta_0$$

   TO:
   Finally, add and subtract $\hat{\theta}'\left(W'R^{-1}W + \Sigma\right)\hat{\theta}$ and rearrange to yield

$$y'R^{-1}y + \left(\theta - \widehat{\theta}\right)'\left(W'R^{-1}W + \Sigma\right)\left(\theta - \widehat{\theta}\right) - \widehat{\theta}'\left(W'R^{-1}W + \Sigma\right)\widehat{\theta} + \theta_0'\Sigma\theta_0$$

6. Page 172, equation at the bottom,
   CHANGE:

$$\ln p\left(\theta|y,\theta_0,B,G,R\right) \propto -\frac{1}{2}\left(\theta - \widehat{\theta}\right)'\left(W'R^{-1}W + \Sigma\right)\left(\theta - \widehat{\theta}\right)'$$

   TO:

$$\ln p\left(\theta|y,\theta_0,B,G,R\right) \propto -\frac{1}{2}\left(\theta - \widehat{\theta}\right)'\left(W'R^{-1}W + \Sigma\right)\left(\theta - \widehat{\theta}\right)$$

7. Page 263,
   CHANGE,

$$v_\beta^* = v + 1,$$

   TO:

$$v_\beta^* = v_\beta + 1,$$

8. Page 265,
   Delete Subsection "An Alternative Parametrisation" (up to and including eqn (5.82)).

9. Page 353, Eq. (7.41), remove "comma" before "$i = 1, \ldots, m$"
   CHANGE:

$$y|\mu, \alpha, \delta, \sigma^2 \sim N\left(1\mu + \sum\nolimits_{i=1}^{m} X_i\left(\alpha_i \delta_i\right), I\sigma^2\right), \quad , i = 1, \ldots, m \tag{7.41}$$

   TO:

$$y|\mu, \alpha, \delta, \sigma^2 \sim N\left(1\mu + \sum\nolimits_{i=1}^{m} X_i\left(\alpha_i \delta_i\right), I\sigma^2\right), \quad i = 1, \ldots, m \tag{7.41}$$

10. Page 361,
    CHANGE:
    "In terms of $p-$values, a threshold of 0.05 guarantees that the expected number of false positives over the whole family of m tests is less than or equal to $0.05\,m$, far too large for modern genomewide studies."
    TO:
    "In terms of $p-$values, a threshold of 0.05 guarantees that the expected number of false positives over the whole family of m tests is $0.05\,m$, far too large for modern genomewide studies."

11. Page 361,
    CHANGE:
    "Then for the whole family of m tests, when all the m hypotheses are true nulls, the expected number of false positives is less than or equal to $m(\alpha/m) = \alpha$."
    TO:
    "Then for the whole family of m tests, when all the m hypotheses are true nulls, the expected number of false positives is equal to $m(\alpha/m) = \alpha$."

12. Page 363,
    CHANGE:
    Then the expected number of false discoveries over the m tests is

$$\mathrm{E}\left(V\right) = \mathrm{E}\left(\sum\nolimits_{i \in M_0} V_i\right) = \sum\nolimits_{i \in M_0} \mathrm{E}\left(V_i\right) \leq m_0 \alpha_B.$$

    TO:
    Then, for continuously distributed tests statistics, the expected number of false discoveries over the m tests is

$$\mathrm{E}\left(V\right) = \mathrm{E}\left(\sum\nolimits_{i \in M_0} V_i\right) = \sum\nolimits_{i \in M_0} \mathrm{E}\left(V_i\right) = m_0 \alpha_B.$$

13. Page 369,
    CHANGE:

"This is in contrast with the meaning attached to the $p-$value for $i = 19$ equal to 0.02845: in conceptual replications of the experiment where a total of m hypotheses are tested, the expected number of false discoveries among the hypotheses rejected at the cutoff 0.02845 or smaller is $0.02845 \times m$ or smaller."
TO:
"This is in contrast with the meaning attached to the $p-$value for $i = 19$ equal to 0.02845: in conceptual replications of the experiment where a total of m hypotheses are tested of which m_0 are true nulls, for continuously distributed test statistics, the expected number of false discoveries is $0.02845\, m_0$."

14. Page 373, paragraph above Eq. (8.23), defines the number of false discoveries $V(A)$, given that k hypotheses are rejected, as binomially distributed $Bi(k, \Pr(H = 0|z \in A)$. Another definition can be found on page 376, paragraph above Eq. (8.33), concerning the expected number of false discoveries. Both definitions may benefit from a little explanation that is provided below.

Background

Let the binary random variable $V_i = 1$ if hypothesis i is rejected and if hypothesis i is a true null; $V_i = 0$, otherwise. Using the notation on page 363, we can write $V_i = I(H_i = 0, R_i = 1)$, where $R_i = 1$ if $z_i \in A$ and z_i is the observed value of the scalar random variable Z_i. Treating the hypotheses as random, it follows that V_i is a Bernoulli random variable with probability $\Pr(H_i = 0, Z_i \in A)$; that is $V_i \sim Br(\Pr(H_i = 0, Z_i \in A)$.

$$V_i \sim Br(\Pr(H_i = 0, Z_i \in A).$$

- **Page 373**. Consider the binary random variable $V_i|R_i = 1$ that takes the value 1 if $H_i = 0$ and 0 otherwise. Then $[V_i|R_i = 1] \sim Br(\Pr(H_i = 0|Z_i \in A)$, a Bernoulli distributed random variable. If in a series of independent trials the total number of rejections is $R = k$, among these k rejections, the total number of false discoveries is binomially distributed:

  $$[V|R = k] \sim Bi(k, \Pr(H = 0|Z \in A).$$

  The expected number of false discoveries among the k rejections is

  $$\mathrm{E}(V|R = k) = k\ \Pr(H = 0|Z \in A).$$

  The subscript i has been omitted because $\Pr(H_i = 0|Z_i \in A)$ is the same for each i.

- **Page 376**. Consider an experiment where m independent hypotheses are tested. Given the definition of V_i and the independence assumption, the total number of false discoveries, $V = \sum_{i=1}^{m} V_i$, is then binomially distributed

  $$V \sim Bi(m, \Pr(H = 0, Z \in A)),$$

with expected value

$$\mathrm{E}(V) = m\ \Pr(H = 0, Z \in A)) = m\ \Pr(Z \in A | H = 0)\pi_0,$$

where $\pi_0 = \Pr(H = 0)$, the prior probability of a true null.

Note that: (i) hypotheses H_i are treated as random (with a common prior π_0), (ii) random vectors (H_i, Z_i), $i = 1, 2, \ldots, m$, are independently distributed, (iii) the rejection region A is data-independent and fixed in advance.

15. Page 382, CHANGE

$$\frac{\widehat{\mathrm{E}}\left(V | r, \widehat{\Pr}\left(\delta_i = 0 | y\right)\right)}{R} = \frac{\sum_{i=1}^{m} \widehat{\Pr}\left(\delta_i = 0 | y\right) r_i}{R}, \tag{8.42}$$

TO:

$$\frac{\widehat{\mathrm{E}}\left(V | r\right)}{R} = \frac{\sum_{i=1}^{m} \widehat{\Pr}\left(\delta_i = 0 | y\right) r_i}{R}, \tag{8.42}$$

16. Page 382, REPLACE THE CONTENT OF THE SUBSECTION:
"Posterior Distribution of False Discovery Rate"
WITH:
The following approach provides a Monte Carlo estimator of the complete posterior distribution of the fully Bayesian FDP. This distribution reflects posterior uncertainty of the parameters of the Bayesian model that propagates on to the FDP. The protocol is as follows. First execute the McMC sampler and store the draws $\delta_i^{[j]}$ at each cycle, where i refers to the marker genotype and j to the McMC cycle. On completion choose a discovery set; an informal approach could be based on selecting the marker genotypes whose Monte Carlo estimates of the posterior probability of non-zero effects is larger than a chosen value $\varphi^\star$. The resulting discovery set includes R marker genotypes. Then, from the stored draws $\delta_i^{[j]}$ *belonging to the discovery set*, compute at each McMC cycle the false discovery proportion for the discovery set. For example, at round j compute

$$\begin{aligned} Q^{[j]} &= \frac{\sum_{i=1}^{m} \left(1 - \delta_i^{[j]}\right) r_i}{R}, \\ Q^{[j]} &\in \left\{0, \frac{1}{R}, \frac{2}{R}, \ldots, 1\right\}, \quad j = 1, 2, \ldots, T, \end{aligned} \tag{8.43}$$

where m is the number of markers and T is the number of samples. The numerator in the first line counts the number of indices i for which $\delta_i^{[j]} = 0$ and $r_i = 1$ in the jth iteration of the McMC chain. The quantity $Q^{[j]}$ is the jth McMC draw from the discrete posterior distribution of the Bayesian FDP for the discovery set chosen.

Importantly, the discovery set is chosen only once and used repeatedly across all the draws of the McMC chain. With enough McMC samples the discreteness of the distribution $[Q \mid y, r]$ should become visible.

The MC (ordinary) Bayesian estimator of FDR is obtained by averaging over $Q^{[j]}$:

$$\hat{\mathrm{E}}(Q|y,r) = \frac{1}{T}\sum_{j=1}^{T} Q^{[j]},$$

that is an alternative way of expressing (8.42).

The two-component normal mixture model implemented in the example below uses rules of the type $r_i = I\left(y_i \geq y^{\star}\right)$ or $r_i = I\left(\hat{\varphi}_i \geq \varphi^{\star}\right)$.

A recent McMC implementation of the Bayesian FDR was described by de los Campos et al (2022).