

CHAPTER 3

0301.93.21	0301.93.31	0301.99.11	0301.99.22	0301.99.24
0301.99.31	0301.99.42			
BREEDING CARP, OTHER THAN FRY; BREEDING MILKFISH AND GROUPER, FRY; OTHER BREEDING FISH FRY; OTHER CARP, FOR BREEDING; BREEDING MILKFISH, OTHER THAN FRY				
Breeding fish are accompanied by certification from the competent authorities as provided for under the national law.				
General requirements on appearance:				
Well-proportioned body, no deformity, full and normal fins, no scratches, no grease loss, fish of even size, with no sign of disease and certified by competent authorities as fit for breeding				
Activeness:				
Fish should be active, swift, swimming under the water in groups.				
Weight and size:				
Depends on each species and hatchery time.				
(Source: Viet Nam)				

0301.99.22	0301.99.23	0301.99.42
OTHER CARP		
Other carp included under 0301.99.22, 0301.99.23 and 0301.99.42 are those carps not included under subheading 0301.93, for example:		
- Genus <i>Abramis</i>: Carp bream (<i>Abramis brama</i>) - Genus <i>Aristichthys</i>: Bighead carp (<i>Aristichthys nobilis</i>) - Genus <i>Barbodes</i>: Carnatic carp (<i>Barbodes carnaticus</i>) - Genus <i>Culter</i>: Predatory carp (<i>Culter erythropterus</i>) - Genus <i>Epalzeorhynchus</i>: Red-tailed black shark (<i>Epalzeorhynchus bicolor</i>) and Red-finned black shark or Rainbow shark (<i>Epalzeorhynchus frenatus</i>) - Genus <i>Henicorhynchus</i>: Siamese mud carp (<i>Henicorhynchus siamensis</i>)		
Source: http://www.newworldencyclopedia.org/entry/Carp		
Refer to the link below to view a more detailed list of subfamilies and genera of carp. https://en.wikipedia.org/wiki/Cyprinidae#Subfamilies_and_genera		
(Source: Philippines)		

0301.99.31	0301.99.32	0303.89.23
MILKFISH (<i>CHANOS CHANOS</i>)		
Kingdom: <i>Animalia</i> Class: <i>Actinopterygii</i> Order: <i>Gonorynchiformes</i> Family: <i>Chanidae</i> Genus: <i>Chanos</i> Species: <i>Chanos chanos</i> Weight is not less than 800 grams. The milkfish is found in fresh, marine as well as brackish water. Size : up to 180 cm		
		
(Source: Indonesia)		

Picture 1. Milkfish

0301.99.33
LEOPARD CORAL GROUPER (<i>PLECTROPOMUS LEOPARDUS</i>)
Kingdom: <i>Animalia</i> Phylum: <i>Chordata</i> Class: <i>Actinopterygii</i> Order: <i>Perciformes</i> Family: <i>Serranidae</i> Genus: <i>Plectropomus</i> Species: <i>Plectropomus leopardus</i> Weight is not less than 600 grams. Size: 35-120 cm



Picture 1. Leopard Coral Grouper

(Source: Indonesia)

0301.99.34

BROWN-MARbled GROUper (EPINEPHELUS FUSCOGUTTATUS)

Kingdom: *Animalia*
 Phylum: *Chordata*
 Class: *Actinopterygii*
 Order: *Perciformes*
 Family: *Serranidae*
 Genus: *Epinephelus*
 Species: *Epinephelus fuscoguttatus*
 Weight is not less than 600 grams.
 Size : 50-120 cm



Picture 1. Brown-Marbled Grouper

(Source: Indonesia)

0301.99.35

HUMPBAC GROUPE (CROMILEPTES ALTIVELIS)

Kingdom: *Animalia*
Phylum: *Chordata*
Class: *Actinopterygii*
Order: *Perciformes*
Family: *Serranidae*
Genus: *Cromileptes*
Species: *Cromileptes altivelis*
Weight is not less than 600 grams.
Size : up to 70 cm



Picture 1. Humpback Grouper

(Source: Indonesia)

0301.99.41

TILAPIAS (OREOCHROMIS SPP.)

Categorized as freshwater fish for consumption

- **Shape:** elongated and compressed body, big eye and bluish green at the edge.
- **Color:** silver with olive/grey/black bars, and often reddish during the breeding season.
- **Size:** up to 60 cm



Picture 1. Tilapia

(Source: Indonesia)

0303.49.10

LONGTAIL TUNA (*THUNNUS TONGGOL*)

Kingdom: Animalia

Phylum: Chordata

Class: Actinopterygii

Order: Perciformes

Family: Scrombidae

Genus: Thunnus

Species: *Thunnus tonggol*

Weight is generally not less than 1000 grams.

Size around 70-145 cm.

Longtail tuna is one of several fish species of northern bluefin tuna.

Reference: FAO species catalogue. Vol. 2. Scrombrids of the world. p. 92-93.




Picture 1. Longtail Tuna


(Source: Indonesia)

0303.54.10

MACKEREL (*SCOMBER SCOMBRUS*, *SCOMBER AUSTRALASICUS*)



Picture 1. Atlantic Mackerel (*Scomber scombrus*)



Picture 2. Blue Mackerel (*Scomber australasicus*)


(Source: Indonesia)

0303.54.20

PACIFIC MACKEREL (*SCOMBER JAPONICUS*)

- Categorized to marine fish, in Atlantic, Indian and Pacific Oceans
- The shape: head is conical, the mouth is slightly tilted, and the eye is completed with eyelid.

- The color: silver on the sides and blackish pattern on the head to the caudal fin.



Picture 1. Pacific Mackerel (*Scomber japonicus*)

(Source: Indonesia)

0304.89.10

MAHI-MAHI (*CORYPHAENA HIPPURUS*)

Kingdom: *Animalia*
 Class: *Actinopterygii*
 Order: *Perciformes*
 Family: *Coryphaenidae*
 Genus: *Coryphaena*
 Species: *Coryphaena hippurus*

The appearance of mahi-mahi fillet is identified by blood-line on the middle of fillet slice elongated to wholly fish body structure (tail-head).



Picture 1. Mahi-mahi



Picture 2. Mahi-mahi, processed to be fillet

(Source: Indonesia)

0304.99.10

SURIMI (MINCED FISH MEAT)

Frozen surimi is a fish protein product made for further processing. It is made from headed, gutted, cleaned and mechanically deboned fresh fish. Deboned fish meat is then washed, refined, minced and dewatered, and then mixed with cryoprotective food ingredients and frozen.

Cryoprotectants are added to prevent protein denaturation in the frozen state, so that the minced fish meat can be frozen and yet retain the capacity to form gel when heat-treated after thawing. Sugars and/or polyhydric alcohols are commonly used as cryoprotectants.

Reference :

1. Code of Practice for Fish and Fishery Products First Edition World Health Organization Food and Agriculture Organization of the United Nations (FAO).
2. Benjakul et al., 2004; Guenneugues and Morrissey, 2005; Perez-Mateos and Lanier, 2006; Rawdkuen et al., 2008; Campo-Deano and Tovar, 2009.




Picture 1. Surimi

(Source: Indonesia)

0305.59.21	
ANCHOVIES (STOLEPHORUS SPP., COILIA SPP., SETIPINNA SPP., LYCOTRISSA SPP., THRYSSA SPP. AND ENCRASICHOLINA SPP.)	
Family: Engraulidae	
FAO Names: <i>E. Anchovies</i>	
1. Anchovies (<i>Engraulis spp</i> (8 species) are anchovies species that are found in European countries, California, Argentina, Australia, South Africa and Japan. 2. Anchovies (<i>Stolephous spp.</i> (18 species), <i>coilia spp.</i> (12 species), <i>Setipinne spp.</i> (7 species), <i>Lycotrissa crocodilus</i> and <i>Thryssa spp.</i> (24 species), <i>Encrasicholina spp.</i> (5 species)) are mostly found in ASEAN waters	
There are other species such as <i>Anchoviella spp.</i>(15 species), <i>Anchoa spp.</i>(34 species), <i>Anchovia spp.</i> (3 species) <i>Centgraulia spp.</i>(2 species), <i>Jurengrualiajurunesis.</i>, <i>Peterngualisatherinoides</i>, <i>Lycengraulis spp.</i>(3 species), <i>Amazonsprattus scintilla</i>, <i>Encrasicholina spp.</i>(5 species) and <i>Papuengraulismicropinna</i>. No habitat in Asian waters.	
(Based on FAO Species Catalogue Vol. 7, Clupeoid Fishes Of The World).	
(Source: Malaysia)	

0305.72.11	0305.72.19
FISH MAWS	
The fish maw or swim bladder, gas bladder or air bladder is an internal gas-filled organ that contributes to the ability of a fish to maintain its buoyancy, and thus to stay at the current water depth without having to waste energy in swimming. The swim bladder is also used as a stabilizing agent because in the upright position the center of mass is below the center of volume due to the dorsal position of the swim bladder.	
In the trade of fishery industries, whole fresh fish including part of fish (e.g. fillets, meat, liver, maw, roes, tails etc.) are separated by type of fish. The sorting or separating according to type of fish is very important in terms or pricing, identification, for processing to make into other product such as minced fish, dried fish product, smoked product, salted fish product or other prepared product based on fish.	
Quality fish maw should have an amber and clear colour, smooth and delightful texture. The darker the colour (golden yellow), less translucency (dark brown), less white tissues and more wrinkles, meaning older the fish maw is, thus, the more medical benefits. The medical benefit in fish maw works best after storage in a dry place for 1 to 3 years when the fishy smell and stickiness of gluten are reduced.	



Picture 1. General Fish Anatomy





Picture 1. Fresh Eel Maws



Picture 2. Fresh Cod Maws



Picture 3. Dried Cod fish maws



(Source: Malaysia)

0306.35.10	0306.36.11	0306.36.12	0306.36.13	0306.36.19
SHRIMPS AND PRAWNS, BREEDING				
Breeding shrimps and prawns are accompanied by certification from the competent authorities as provided for under the national law.				
General requirements on appearance: well-proportioned body, no deformity, no damage to the crust and no sign of disease.				

ใช้ในราชการมตฺลการเท่านั้น
 ไม่มีผลผูกพันทางกฎหมาย