



MAT1320-Linear Algebra Lecture Notes

Basis, Dimension and Coordinates

Mehmet E. KÖROĞLU

Fall 2020

YILDIZ TECHNICAL UNIVERSITY, DEPARTMENT OF MATHEMATICS

mkoroglu@yildiz.edu.tr

Table of contents

1. Basis and Dimension

2. Coordinates

Basis and Dimension

Basis and Dimension

Definition

A set $\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ is a **base** of V if it has the following two properties:

Basis and Dimension

Definition

A set $\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ is a **base** of V if it has the following two properties:

1. \mathcal{B} is linearly independent,

Basis and Dimension

Definition

A set $\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ is a **base** of V if it has the following two properties:

1. \mathcal{B} is linearly independent,

Basis and Dimension

Definition

A set $\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ is a **base** of V if it has the following two properties:

1. \mathcal{B} is linearly independent,
2. $\langle \mathcal{B} \rangle = V$ i.e., \mathcal{B} spans V .

Basis and Dimension

Definition

A set $\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ is a **base** of V if it has the following two properties:

1. \mathcal{B} is linearly independent,
2. $\langle \mathcal{B} \rangle = V$ i.e., \mathcal{B} spans V .

Basis and Dimension

Definition

A set $\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ is a **base** of V if it has the following two properties:

1. \mathcal{B} is linearly independent,
2. $\langle \mathcal{B} \rangle = V$ i.e., \mathcal{B} spans V .

Note: All bases of a vector space V have the same number of elements.

Basis and Dimension

Definition

A set $\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ is a **base** of V if it has the following two properties:

1. \mathcal{B} is linearly independent,
2. $\langle \mathcal{B} \rangle = V$ i.e., \mathcal{B} spans V .

Note: All bases of a vector space V have the same number of elements.

Definition

A vector space V is said to be of **finite dimension n** or **n -dimensional** or **n -space**, written $\dim(V) = n$, if V has a basis with n elements.

Basis and Dimension

Definition

A set $\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ is a **base** of V if it has the following two properties:

1. \mathcal{B} is linearly independent,
2. $\langle \mathcal{B} \rangle = V$ i.e., \mathcal{B} spans V .

Note: All bases of a vector space V have the same number of elements.

Definition

A vector space V is said to be of **finite dimension n** or **n -dimensional** or **n -space**, written $\dim(V) = n$, if V has a basis with n elements. The vector space $\{0\}$ is defined to have dimension 0.

Example

1. $\mathcal{B} = \{1\} \subset V = \mathbb{R} \Rightarrow \mathbb{R} = \langle 1 \rangle$ and $\dim(V) = 1$.

Example

1. $\mathcal{B} = \{1\} \subset V = \mathbb{R} \Rightarrow \mathbb{R} = \langle 1 \rangle$ and $\dim(V) = 1$.
2. $\mathcal{B} = \{\vec{e}_1 = (1, 0), \vec{e}_2 = (0, 1)\} \subset V = \mathbb{R}^2 \Rightarrow \mathbb{R}^2 = \langle \mathcal{B} \rangle$
and $\dim(V) = 2$.

Example

1. $\mathcal{B} = \{1\} \subset V = \mathbb{R} \Rightarrow \mathbb{R} = \langle 1 \rangle$ and $\dim(V) = 1$.
2. $\mathcal{B} = \{\vec{e}_1 = (1, 0), \vec{e}_2 = (0, 1)\} \subset V = \mathbb{R}^2 \Rightarrow \mathbb{R}^2 = \langle \mathcal{B} \rangle$
and $\dim(V) = 2$.
3. $\mathcal{B} = \{\vec{e}_1 = (1, 0, 0), \vec{e}_2 = (0, 1, 0), \vec{e}_3 = (0, 0, 1)\} \subset V = \mathbb{R}^3 \Rightarrow \mathbb{R}^3 = \langle \mathcal{B} \rangle$ and $\dim(V) = 3$.

Example

1. $\mathcal{B} = \{1\} \subset V = \mathbb{R} \Rightarrow \mathbb{R} = \langle 1 \rangle$ and $\dim(V) = 1$.
2. $\mathcal{B} = \{\vec{e}_1 = (1, 0), \vec{e}_2 = (0, 1)\} \subset V = \mathbb{R}^2 \Rightarrow \mathbb{R}^2 = \langle \mathcal{B} \rangle$
and $\dim(V) = 2$.
3. $\mathcal{B} = \{\vec{e}_1 = (1, 0, 0), \vec{e}_2 = (0, 1, 0), \vec{e}_3 = (0, 0, 1)\} \subset$
 $V = \mathbb{R}^3 \Rightarrow \mathbb{R}^3 = \langle \mathcal{B} \rangle$ and $\dim(V) = 3$.
4. $\mathcal{B} = \{\vec{e}_1 = (1, 0, \dots, 0), \dots, \vec{e}_n = (0, \dots, 0, 1)\} \subset V$
 $= \mathbb{R}^n \Rightarrow \mathbb{R}^n = \langle \mathcal{B} \rangle$ and $\dim(V) = n$.

Example

1. $\mathcal{B} = \{1\} \subset V = \mathbb{R} \Rightarrow \mathbb{R} = \langle 1 \rangle$ and $\dim(V) = 1$.
2. $\mathcal{B} = \{\vec{e}_1 = (1, 0), \vec{e}_2 = (0, 1)\} \subset V = \mathbb{R}^2 \Rightarrow \mathbb{R}^2 = \langle \mathcal{B} \rangle$ and $\dim(V) = 2$.
3. $\mathcal{B} = \{\vec{e}_1 = (1, 0, 0), \vec{e}_2 = (0, 1, 0), \vec{e}_3 = (0, 0, 1)\} \subset V = \mathbb{R}^3 \Rightarrow \mathbb{R}^3 = \langle \mathcal{B} \rangle$ and $\dim(V) = 3$.
4. $\mathcal{B} = \{\vec{e}_1 = (1, 0, \dots, 0), \dots, \vec{e}_n = (0, \dots, 0, 1)\} \subset V = \mathbb{R}^n \Rightarrow \mathbb{R}^n = \langle \mathcal{B} \rangle$ and $\dim(V) = n$.

Example

1. $\mathcal{B} = \{1\} \subset V = \mathbb{R} \Rightarrow \mathbb{R} = \langle 1 \rangle$ and $\dim(V) = 1$.
2. $\mathcal{B} = \{\vec{e}_1 = (1, 0), \vec{e}_2 = (0, 1)\} \subset V = \mathbb{R}^2 \Rightarrow \mathbb{R}^2 = \langle \mathcal{B} \rangle$ and $\dim(V) = 2$.
3. $\mathcal{B} = \{\vec{e}_1 = (1, 0, 0), \vec{e}_2 = (0, 1, 0), \vec{e}_3 = (0, 0, 1)\} \subset V = \mathbb{R}^3 \Rightarrow \mathbb{R}^3 = \langle \mathcal{B} \rangle$ and $\dim(V) = 3$.
4. $\mathcal{B} = \{\vec{e}_1 = (1, 0, \dots, 0), \dots, \vec{e}_n = (0, \dots, 0, 1)\} \subset V = \mathbb{R}^n \Rightarrow \mathbb{R}^n = \langle \mathcal{B} \rangle$ and $\dim(V) = n$.

Note: All vector spaces given above are of finite dimension.

Example

$$1. \mathcal{B} = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right\} \subset V = \mathcal{M}_{2 \times 2}(\mathbb{R}) \Rightarrow \mathcal{M}_{2 \times 2}(\mathbb{R}) = \langle \mathcal{B} \rangle \text{ and } \dim(\mathcal{M}_{2 \times 2}(\mathbb{R})) = 4.$$

Example

1. $\mathcal{B} = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right\} \subset V = \mathcal{M}_{2 \times 2}(\mathbb{R}) \Rightarrow \mathcal{M}_{2 \times 2}(\mathbb{R}) = \langle \mathcal{B} \rangle$ and $\dim(\mathcal{M}_{2 \times 2}(\mathbb{R})) = 4$.
2. $\mathcal{B} = \{1, x, x^2, \dots, x^{n-1}\} \subset V = \mathcal{P}_{n-1}(x) = \{a_0 + a_1x + \dots + a_{n-1}x^{n-1} \mid a_i \in \mathbb{R}\} \Rightarrow \mathcal{P}_{n-1}(x) = \langle \mathcal{B} \rangle$ and $\dim(\mathcal{P}_{n-1}(x)) = n$.

Example

1. $\mathcal{B} = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right\} \subset V = \mathcal{M}_{2 \times 2}(\mathbb{R}) \Rightarrow \mathcal{M}_{2 \times 2}(\mathbb{R}) = \langle \mathcal{B} \rangle$ and $\dim(\mathcal{M}_{2 \times 2}(\mathbb{R})) = 4$.
2. $\mathcal{B} = \{1, x, x^2, \dots, x^{n-1}\} \subset V = \mathcal{P}_{n-1}(x) = \{a_0 + a_1x + \dots + a_{n-1}x^{n-1} \mid a_i \in \mathbb{R}\} \Rightarrow \mathcal{P}_{n-1}(x) = \langle \mathcal{B} \rangle$ and $\dim(\mathcal{P}_{n-1}(x)) = n$.
3. $\mathcal{B} = \{1, x, x^2, \dots\} \subset V = \mathcal{P}(x) = \{\text{the set of all polynomials}\} \Rightarrow \mathcal{P}(x) = \langle \mathcal{B} \rangle$ and $\dim(\mathcal{P}(x)) = \infty$.

Example

1. $\mathcal{B} = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right\} \subset V = \mathcal{M}_{2 \times 2}(\mathbb{R}) \Rightarrow \mathcal{M}_{2 \times 2}(\mathbb{R}) = \langle \mathcal{B} \rangle$ and $\dim(\mathcal{M}_{2 \times 2}(\mathbb{R})) = 4$.
2. $\mathcal{B} = \{1, x, x^2, \dots, x^{n-1}\} \subset V = \mathcal{P}_{n-1}(x) = \{a_0 + a_1x + \dots + a_{n-1}x^{n-1} \mid a_i \in \mathbb{R}\} \Rightarrow \mathcal{P}_{n-1}(x) = \langle \mathcal{B} \rangle$ and $\dim(\mathcal{P}_{n-1}(x)) = n$.
3. $\mathcal{B} = \{1, x, x^2, \dots\} \subset V = \mathcal{P}(x) = \{\text{the set of all polynomials}\} \Rightarrow \mathcal{P}(x) = \langle \mathcal{B} \rangle$ and $\dim(\mathcal{P}(x)) = \infty$.
4. Let $V = \left\{ \vec{\mathbf{0}} \right\}$ be the zero space, then $V = \left\langle \vec{\mathbf{0}} \right\rangle$ and $\dim(V) = 0$.

Example

Determine whether the set

$\mathcal{B} = \{\vec{v}_1 = (1, 2, 0), \vec{v}_2 = (2, 0, 1), \vec{v}_3 = (1, 2, 1)\}$ is a basis for \mathbb{R}^3 .

Example

Determine whether the set

$\mathcal{B} = \{\vec{v}_1 = (1, 2, 0), \vec{v}_2 = (2, 0, 1), \vec{v}_3 = (1, 2, 1)\}$ is a basis for \mathbb{R}^3 .

1. Linear Independence: For $x_1, x_2, x_3 \in \mathbb{R}$

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3 = (0, 0, 0)$$

Example

Determine whether the set

$\mathcal{B} = \{\vec{v}_1 = (1, 2, 0), \vec{v}_2 = (2, 0, 1), \vec{v}_3 = (1, 2, 1)\}$ is a basis for \mathbb{R}^3 .

1. Linear Independence: For $x_1, x_2, x_3 \in \mathbb{R}$

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3 = (0, 0, 0)$$

$$\Rightarrow x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1) = (0, 0, 0)$$

Basis and Dimension

Example

Determine whether the set

$\mathcal{B} = \{\vec{v}_1 = (1, 2, 0), \vec{v}_2 = (2, 0, 1), \vec{v}_3 = (1, 2, 1)\}$ is a basis for \mathbb{R}^3 .

1. Linear Independence: For $x_1, x_2, x_3 \in \mathbb{R}$

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3 = (0, 0, 0)$$

$$\Rightarrow x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1) = (0, 0, 0)$$

$$\Rightarrow \begin{cases} x_1 + 2x_2 + x_3 = 0 \\ 2x_1 + 2x_3 = 0 \\ x_2 + x_3 = 0 \end{cases}$$

Basis and Dimension

Example

Determine whether the set

$\mathcal{B} = \{\vec{v}_1 = (1, 2, 0), \vec{v}_2 = (2, 0, 1), \vec{v}_3 = (1, 2, 1)\}$ is a basis for \mathbb{R}^3 .

1. Linear Independence: For $x_1, x_2, x_3 \in \mathbb{R}$

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3 = (0, 0, 0)$$

$$\Rightarrow x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1) = (0, 0, 0)$$

$$\Rightarrow \begin{cases} x_1 + 2x_2 + x_3 = 0 \\ 2x_1 + 2x_3 = 0 \\ x_2 + x_3 = 0 \end{cases} \Rightarrow \begin{pmatrix} 1 & 2 & 1 \\ 2 & 0 & 2 \\ 0 & 1 & 1 \end{pmatrix}$$

Basis and Dimension

Example

Determine whether the set

$\mathcal{B} = \{\vec{v}_1 = (1, 2, 0), \vec{v}_2 = (2, 0, 1), \vec{v}_3 = (1, 2, 1)\}$ is a basis for \mathbb{R}^3 .

1. Linear Independence: For $x_1, x_2, x_3 \in \mathbb{R}$

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3 = (0, 0, 0)$$

$$\Rightarrow x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1) = (0, 0, 0)$$

$$\Rightarrow \begin{cases} x_1 + 2x_2 + x_3 = 0 \\ 2x_1 + 2x_3 = 0 \\ x_2 + x_3 = 0 \end{cases} \Rightarrow \begin{pmatrix} 1 & 2 & 1 \\ 2 & 0 & 2 \\ 0 & 1 & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Basis and Dimension

Example

Determine whether the set

$\mathcal{B} = \{\vec{v}_1 = (1, 2, 0), \vec{v}_2 = (2, 0, 1), \vec{v}_3 = (1, 2, 1)\}$ is a basis for \mathbb{R}^3 .

1. Linear Independence: For $x_1, x_2, x_3 \in \mathbb{R}$

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3 = (0, 0, 0)$$

$$\Rightarrow x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1) = (0, 0, 0)$$

$$\Rightarrow \begin{cases} x_1 + 2x_2 + x_3 = 0 \\ 2x_1 + 2x_3 = 0 \\ x_2 + x_3 = 0 \end{cases} \Rightarrow \begin{pmatrix} 1 & 2 & 1 \\ 2 & 0 & 2 \\ 0 & 1 & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$\Rightarrow x_1 = x_2 = x_3 = 0.$$

Basis and Dimension

Example

Determine whether the set

$\mathcal{B} = \{\vec{v}_1 = (1, 2, 0), \vec{v}_2 = (2, 0, 1), \vec{v}_3 = (1, 2, 1)\}$ is a basis for \mathbb{R}^3 .

1. Linear Independence: For $x_1, x_2, x_3 \in \mathbb{R}$

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3 = (0, 0, 0)$$

$$\Rightarrow x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1) = (0, 0, 0)$$

$$\Rightarrow \begin{cases} x_1 + 2x_2 + x_3 = 0 \\ 2x_1 + 2x_3 = 0 \\ x_2 + x_3 = 0 \end{cases} \Rightarrow \begin{pmatrix} 1 & 2 & 1 \\ 2 & 0 & 2 \\ 0 & 1 & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$\Rightarrow x_1 = x_2 = x_3 = 0.$$

Thus $\vec{v}_1, \vec{v}_2, \vec{v}_3$ are linearly independent.

2. Span: For all $(x, y, z) \in \mathbb{R}^3$

$$(x, y, z) = x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3$$

2. Span: For all $(x, y, z) \in \mathbb{R}^3$

$$(x, y, z) = x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3$$

$$(x, y, z) = x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1)$$

2. Span: For all $(x, y, z) \in \mathbb{R}^3$

$$(x, y, z) = x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3$$

$$(x, y, z) = x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1)$$

$$x_1 + 2x_2 + x_3 = x$$

$$2x_1 + 2x_3 = y$$

$$x_2 + x_3 = z$$

2. Span: For all $(x, y, z) \in \mathbb{R}^3$

$$(x, y, z) = x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3$$

$$(x, y, z) = x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1)$$

$$\begin{array}{rcl} x_1 + 2x_2 + x_3 & = & x \\ 2x_1 + 2x_3 & = & y \\ x_2 + x_3 & = & z \end{array} \Rightarrow \left(\begin{array}{ccc|c} 1 & 2 & 1 & x \\ 2 & 0 & 2 & y \\ 0 & 1 & 1 & z \end{array} \right)$$

Basis and Dimension

2. Span: For all $(x, y, z) \in \mathbb{R}^3$

$$(x, y, z) = x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3$$

$$(x, y, z) = x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1)$$

$$\begin{array}{rcl} x_1 + 2x_2 + x_3 = x \\ 2x_1 + 2x_3 = y \\ x_2 + x_3 = z \end{array} \Rightarrow \left(\begin{array}{ccc|c} 1 & 2 & 1 & x \\ 2 & 0 & 2 & y \\ 0 & 1 & 1 & z \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 0 & 0 & \frac{2x+y-4z}{4} \\ 0 & 1 & 0 & \frac{2x-y}{4} \\ 0 & 0 & 1 & \frac{-2x+y+4z}{4} \end{array} \right).$$

2. Span: For all $(x, y, z) \in \mathbb{R}^3$

$$(x, y, z) = x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3$$

$$(x, y, z) = x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1)$$

$$\begin{aligned} x_1 + 2x_2 + x_3 &= x \\ 2x_1 + 2x_3 &= y \\ x_2 + x_3 &= z \end{aligned} \Rightarrow \left(\begin{array}{ccc|c} 1 & 2 & 1 & x \\ 2 & 0 & 2 & y \\ 0 & 1 & 1 & z \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 0 & 0 & \frac{2x+y-4z}{4} \\ 0 & 1 & 0 & \frac{2x-y}{4} \\ 0 & 0 & 1 & \frac{-2x+y+4z}{4} \end{array} \right).$$

Hence, we have $x_1 = \frac{2x+y-4z}{4}$, $x_2 = \frac{2x-y}{4}$, $x_3 = \frac{-2x+y+4z}{4}$

Basis and Dimension

2. Span: For all $(x, y, z) \in \mathbb{R}^3$

$$(x, y, z) = x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3$$

$$(x, y, z) = x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1)$$

$$\begin{aligned} x_1 + 2x_2 + x_3 &= x \\ 2x_1 + 2x_3 &= y \\ x_2 + x_3 &= z \end{aligned} \Rightarrow \left(\begin{array}{ccc|c} 1 & 2 & 1 & x \\ 2 & 0 & 2 & y \\ 0 & 1 & 1 & z \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 0 & 0 & \frac{2x+y-4z}{4} \\ 0 & 1 & 0 & \frac{2x-y}{4} \\ 0 & 0 & 1 & \frac{-2x+y+4z}{4} \end{array} \right).$$

Hence, we have $x_1 = \frac{2x+y-4z}{4}$, $x_2 = \frac{2x-y}{4}$, $x_3 = \frac{-2x+y+4z}{4}$ and

$$(x, y, z) = x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1).$$

Basis and Dimension

2. Span: For all $(x, y, z) \in \mathbb{R}^3$

$$(x, y, z) = x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3$$

$$(x, y, z) = x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1)$$

$$\begin{aligned} x_1 + 2x_2 + x_3 &= x \\ 2x_1 + 2x_3 &= y \\ x_2 + x_3 &= z \end{aligned} \Rightarrow \left(\begin{array}{ccc|c} 1 & 2 & 1 & x \\ 2 & 0 & 2 & y \\ 0 & 1 & 1 & z \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 0 & 0 & \frac{2x+y-4z}{4} \\ 0 & 1 & 0 & \frac{2x-y}{4} \\ 0 & 0 & 1 & \frac{-2x+y+4z}{4} \end{array} \right).$$

Hence, we have $x_1 = \frac{2x+y-4z}{4}$, $x_2 = \frac{2x-y}{4}$, $x_3 = \frac{-2x+y+4z}{4}$ and

$$(x, y, z) = x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1).$$

Consequently, $\mathbb{R}^3 = \langle \mathcal{B} \rangle$.

Theorem

Let V be a vector space of dimension n and

$\mathcal{B} = \{\vec{\mathbf{v}}_1, \vec{\mathbf{v}}_2, \dots, \vec{\mathbf{v}}_n\} \subset V$. Then the following statements hold:

1. *If \mathcal{B} is a linearly independent set, then $\langle \mathcal{B} \rangle = V$.*

Theorem

Let V be a vector space of dimension n and

$\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\} \subset V$. Then the following statements hold:

1. If \mathcal{B} is a linearly independent set, then $\langle \mathcal{B} \rangle = V$.
2. If $\langle \mathcal{B} \rangle = V$, then \mathcal{B} is a linearly independent set.

Theorem

Let V be a vector space of dimension n and

$\mathcal{B} = \{\vec{\mathbf{v}}_1, \vec{\mathbf{v}}_2, \dots, \vec{\mathbf{v}}_n\} \subset V$. Then the following statements hold:

1. If \mathcal{B} is a linearly independent set, then $\langle \mathcal{B} \rangle = V$.
2. If $\langle \mathcal{B} \rangle = V$, then \mathcal{B} is a linearly independent set.

Theorem

Let V be a vector space of dimension n and

$\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\} \subset V$. Then the following statements hold:

1. If \mathcal{B} is a linearly independent set, then $\langle \mathcal{B} \rangle = V$.
2. If $\langle \mathcal{B} \rangle = V$, then \mathcal{B} is a linearly independent set.

Note: By the above Theorem, in order to check that $\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\} \subset V$ is a basis of the finite dimensional vector space V , it is just sufficient to check linearly independence.

Example

Determine whether the set $\mathcal{B} = \{\vec{\mathbf{v}}_1 = (1, 2), \vec{\mathbf{v}}_2 = (2, 0)\}$ is a basis for \mathbb{R}^2 .

Basis and Dimension

Example

Determine whether the set $\mathcal{B} = \{\vec{\mathbf{v}}_1 = (1, 2), \vec{\mathbf{v}}_2 = (2, 0)\}$ is a basis for \mathbb{R}^2 .

1. Solution: For $x_1, x_2 \in \mathbb{R}$

$$\Rightarrow x_1 \vec{\mathbf{v}}_1 + x_2 \vec{\mathbf{v}}_2 = (0, 0)$$

Basis and Dimension

Example

Determine whether the set $\mathcal{B} = \{\vec{\mathbf{v}}_1 = (1, 2), \vec{\mathbf{v}}_2 = (2, 0)\}$ is a basis for \mathbb{R}^2 .

1. Solution: For $x_1, x_2 \in \mathbb{R}$

$$\Rightarrow x_1 \vec{\mathbf{v}}_1 + x_2 \vec{\mathbf{v}}_2 = (0, 0)$$

$$\Rightarrow x_1(1, 2) + x_2(2, 0) = (0, 0)$$

Basis and Dimension

Example

Determine whether the set $\mathcal{B} = \{\vec{v}_1 = (1, 2), \vec{v}_2 = (2, 0)\}$ is a basis for \mathbb{R}^2 .

1. Solution: For $x_1, x_2 \in \mathbb{R}$

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 = (0, 0)$$

$$\Rightarrow x_1(1, 2) + x_2(2, 0) = (0, 0)$$

$$\Rightarrow \begin{cases} x_1 + 2x_2 = 0 \\ 2x_1 = 0 \end{cases}$$

Basis and Dimension

Example

Determine whether the set $\mathcal{B} = \{\vec{v}_1 = (1, 2), \vec{v}_2 = (2, 0)\}$ is a basis for \mathbb{R}^2 .

1. Solution: For $x_1, x_2 \in \mathbb{R}$

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 = (0, 0)$$

$$\Rightarrow x_1(1, 2) + x_2(2, 0) = (0, 0)$$

$$\Rightarrow \begin{cases} x_1 + 2x_2 = 0 \\ 2x_1 = 0 \end{cases} \Rightarrow \begin{pmatrix} 1 & 2 \\ 2 & 0 \end{pmatrix}$$

Basis and Dimension

Example

Determine whether the set $\mathcal{B} = \{\vec{\mathbf{v}}_1 = (1, 2), \vec{\mathbf{v}}_2 = (2, 0)\}$ is a basis for \mathbb{R}^2 .

1. Solution: For $x_1, x_2 \in \mathbb{R}$

$$\Rightarrow x_1 \vec{\mathbf{v}}_1 + x_2 \vec{\mathbf{v}}_2 = (0, 0)$$

$$\Rightarrow x_1(1, 2) + x_2(2, 0) = (0, 0)$$

$$\Rightarrow \begin{cases} x_1 + 2x_2 = 0 \\ 2x_1 = 0 \end{cases} \Rightarrow \begin{pmatrix} 1 & 2 \\ 2 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Basis and Dimension

Example

Determine whether the set $\mathcal{B} = \{\vec{v}_1 = (1, 2), \vec{v}_2 = (2, 0)\}$ is a basis for \mathbb{R}^2 .

1. Solution: For $x_1, x_2 \in \mathbb{R}$

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 = (0, 0)$$

$$\Rightarrow x_1(1, 2) + x_2(2, 0) = (0, 0)$$

$$\Rightarrow \begin{cases} x_1 + 2x_2 = 0 \\ 2x_1 = 0 \end{cases} \Rightarrow \begin{pmatrix} 1 & 2 \\ 2 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\Rightarrow x_1 = x_2 = 0.$$

Basis and Dimension

Example

Determine whether the set $\mathcal{B} = \{\vec{\mathbf{v}}_1 = (1, 2), \vec{\mathbf{v}}_2 = (2, 0)\}$ is a basis for \mathbb{R}^2 .

1. Solution: For $x_1, x_2 \in \mathbb{R}$

$$\Rightarrow x_1 \vec{\mathbf{v}}_1 + x_2 \vec{\mathbf{v}}_2 = (0, 0)$$

$$\Rightarrow x_1(1, 2) + x_2(2, 0) = (0, 0)$$

$$\Rightarrow \begin{cases} x_1 + 2x_2 = 0 \\ 2x_1 = 0 \end{cases} \Rightarrow \begin{pmatrix} 1 & 2 \\ 2 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\Rightarrow x_1 = x_2 = 0.$$

Therefore, the vectors $\vec{\mathbf{v}}_1, \vec{\mathbf{v}}_2$ are linearly independent. Since $\dim(\mathbb{R}^2) = 2$ we have $\langle \mathcal{B} \rangle = \mathbb{R}^2$ and we conclude that \mathcal{B} is a basis for \mathbb{R}^2 .

2. Solution: For all $(x, y) \in \mathbb{R}^2$

$$(x, y) = x_1 \vec{v}_1 + x_2 \vec{v}_2$$

2. Solution: For all $(x, y) \in \mathbb{R}^2$

$$(x, y) = x_1 \vec{v}_1 + x_2 \vec{v}_2$$

$$(x, y) = x_1(1, 2) + x_2(2, 0)$$

2. Solution: For all $(x, y) \in \mathbb{R}^2$

$$(x, y) = x_1 \vec{v}_1 + x_2 \vec{v}_2$$

$$(x, y) = x_1(1, 2) + x_2(2, 0)$$

$$\begin{cases} x_1 + 2x_2 = x \\ 2x_1 = y \end{cases}$$

2. Solution: For all $(x, y) \in \mathbb{R}^2$

$$(x, y) = x_1 \vec{v}_1 + x_2 \vec{v}_2$$

$$(x, y) = x_1(1, 2) + x_2(2, 0)$$

$$\begin{cases} x_1 + 2x_2 = x \\ 2x_1 = y \end{cases} \Rightarrow \left(\begin{array}{cc|c} 1 & 2 & x \\ 2 & 0 & y \end{array} \right)$$

2. Solution: For all $(x, y) \in \mathbb{R}^2$

$$(x, y) = x_1 \vec{v}_1 + x_2 \vec{v}_2$$

$$(x, y) = x_1(1, 2) + x_2(2, 0)$$

$$\begin{cases} x_1 + 2x_2 = x \\ 2x_1 = y \end{cases} \Rightarrow \left(\begin{array}{cc|c} 1 & 2 & x \\ 2 & 0 & y \end{array} \right) \sim \left(\begin{array}{cc|c} 1 & 0 & \frac{y}{2} \\ 0 & 1 & \frac{2x-y}{4} \end{array} \right)$$

2. Solution: For all $(x, y) \in \mathbb{R}^2$

$$(x, y) = x_1 \vec{v}_1 + x_2 \vec{v}_2$$

$$(x, y) = x_1(1, 2) + x_2(2, 0)$$

$$\begin{cases} x_1 + 2x_2 = x \\ 2x_1 = y \end{cases} \Rightarrow \left(\begin{array}{cc|c} 1 & 2 & x \\ 2 & 0 & y \end{array} \right) \sim \left(\begin{array}{cc|c} 1 & 0 & \frac{y}{2} \\ 0 & 1 & \frac{2x-y}{4} \end{array} \right)$$

This means that, we have $x_1 = \frac{y}{2}$ and $x_2 = \frac{2x-y}{4}$.

2. Solution: For all $(x, y) \in \mathbb{R}^2$

$$(x, y) = x_1 \vec{v}_1 + x_2 \vec{v}_2$$

$$(x, y) = x_1(1, 2) + x_2(2, 0)$$

$$\begin{cases} x_1 + 2x_2 = x \\ 2x_1 = y \end{cases} \Rightarrow \left(\begin{array}{cc|c} 1 & 2 & x \\ 2 & 0 & y \end{array} \right) \sim \left(\begin{array}{cc|c} 1 & 0 & \frac{y}{2} \\ 0 & 1 & \frac{2x-y}{4} \end{array} \right)$$

This means that, we have $x_1 = \frac{y}{2}$ and $x_2 = \frac{2x-y}{4}$. Hence

$$(x, y) = \frac{y}{2}(1, 2) + \left(\frac{2x-y}{4} \right) (2, 0).$$

2. Solution: For all $(x, y) \in \mathbb{R}^2$

$$\begin{aligned}(x, y) &= x_1 \vec{v}_1 + x_2 \vec{v}_2 \\(x, y) &= x_1(1, 2) + x_2(2, 0) \\ \begin{cases} x_1 + 2x_2 = x \\ 2x_1 = y \end{cases} &\Rightarrow \left(\begin{array}{cc|c} 1 & 2 & x \\ 2 & 0 & y \end{array} \right) \sim \left(\begin{array}{cc|c} 1 & 0 & \frac{y}{2} \\ 0 & 1 & \frac{2x-y}{4} \end{array} \right)\end{aligned}$$

This means that, we have $x_1 = \frac{y}{2}$ and $x_2 = \frac{2x-y}{4}$. Hence

$$(x, y) = \frac{y}{2}(1, 2) + \left(\frac{2x-y}{4} \right) (2, 0).$$

Since, $\mathbb{R}^2 = \langle \mathcal{B} \rangle$ the vectors \vec{v}_1, \vec{v}_2 are linearly independent. As a result, \mathcal{B} is a basis for \mathbb{R}^2 .

Example

Let $\mathcal{U} = \{(x, y, z) \mid x + 2z = 0\} \subset \mathbb{R}^3$.

Example

Let $\mathcal{U} = \{(x, y, z) \mid x + 2z = 0\} \subset \mathbb{R}^3$.

1. Show that \mathcal{U} is a subspace of \mathbb{R}^3 .

Example

Let $\mathcal{U} = \{(x, y, z) \mid x + 2z = 0\} \subset \mathbb{R}^3$.

1. Show that \mathcal{U} is a subspace of \mathbb{R}^3 .
2. Find a basis for \mathcal{U} .

Example

Let $\mathcal{U} = \{(x, y, z) \mid x + 2z = 0\} \subset \mathbb{R}^3$.

1. Show that \mathcal{U} is a subspace of \mathbb{R}^3 .
2. Find a basis for \mathcal{U} .
3. $\dim(\mathcal{U}) = ?$

1. **Subspace:** For all $(x, y, z), (a, b, c) \in \mathcal{U}$ and $\alpha, \beta \in \mathbb{R}$

1. Subspace: For all $(x, y, z), (a, b, c) \in \mathcal{U}$ and $\alpha, \beta \in \mathbb{R}$

$$\Rightarrow \alpha(x, y, z) + \beta(a, b, c) = (\alpha x + \beta a, \alpha y + \beta b, \alpha z + \beta c)$$

1. Subspace: For all $(x, y, z), (a, b, c) \in \mathcal{U}$ and $\alpha, \beta \in \mathbb{R}$

$$\Rightarrow \alpha(x, y, z) + \beta(a, b, c) = (\alpha x + \beta a, \alpha y + \beta b, \alpha z + \beta c)$$

$$\Rightarrow \begin{cases} x + 2z = 0 \Rightarrow \alpha(x + 2z) = 0 \\ a + 2c = 0, \Rightarrow \beta(a + 2c) = 0 \end{cases}$$

1. Subspace: For all $(x, y, z), (a, b, c) \in \mathcal{U}$ and $\alpha, \beta \in \mathbb{R}$

$$\Rightarrow \alpha(x, y, z) + \beta(a, b, c) = (\alpha x + \beta a, \alpha y + \beta b, \alpha z + \beta c)$$

$$\Rightarrow \begin{cases} x + 2z = 0 \Rightarrow \alpha(x + 2z) = 0 \\ a + 2c = 0, \Rightarrow \beta(a + 2c) = 0 \end{cases}$$

$$\Rightarrow \alpha(x + 2z) + \beta(a + 2c) = 0$$

1. Subspace: For all $(x, y, z), (a, b, c) \in \mathcal{U}$ and $\alpha, \beta \in \mathbb{R}$

$$\Rightarrow \alpha(x, y, z) + \beta(a, b, c) = (\alpha x + \beta a, \alpha y + \beta b, \alpha z + \beta c)$$

$$\Rightarrow \begin{cases} x + 2z = 0 \Rightarrow \alpha(x + 2z) = 0 \\ a + 2c = 0, \Rightarrow \beta(a + 2c) = 0 \end{cases}$$

$$\Rightarrow \alpha(x + 2z) + \beta(a + 2c) = 0$$

$$\Rightarrow \alpha(x, y, z) + \beta(a, b, c) \in \mathcal{U}.$$

1. Subspace: For all $(x, y, z), (a, b, c) \in \mathcal{U}$ and $\alpha, \beta \in \mathbb{R}$

$$\Rightarrow \alpha(x, y, z) + \beta(a, b, c) = (\alpha x + \beta a, \alpha y + \beta b, \alpha z + \beta c)$$

$$\Rightarrow \begin{cases} x + 2z = 0 \Rightarrow \alpha(x + 2z) = 0 \\ a + 2c = 0, \Rightarrow \beta(a + 2c) = 0 \end{cases}$$

$$\Rightarrow \alpha(x + 2z) + \beta(a + 2c) = 0$$

$$\Rightarrow \alpha(x, y, z) + \beta(a, b, c) \in \mathcal{U}.$$

This says that \mathcal{U} is a subspace of \mathbb{R}^3 .

2. Base: For all $(x, y, z) \in \mathcal{U}$ we have $x + 2z = 0 \Rightarrow x = -2z$, a linear equation with three unknowns.

2. Base: For all $(x, y, z) \in \mathcal{U}$ we have $x + 2z = 0 \Rightarrow x = -2z$, a linear equation with three unknowns. This system has infinitely many solutions depending on two parameters.

2. Base: For all $(x, y, z) \in \mathcal{U}$ we have $x + 2z = 0 \Rightarrow x = -2z$, a linear equation with three unknowns. This system has infinitely many solutions depending on two parameters.

$$\Rightarrow \begin{cases} x = -2z \\ y = t \\ z = s \end{cases} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -2s \\ t \\ s \end{bmatrix}$$

Basis and Dimension

2. Base: For all $(x, y, z) \in \mathcal{U}$ we have $x + 2z = 0 \Rightarrow x = -2z$, a linear equation with three unknowns. This system has infinitely many solutions depending on two parameters.

$$\begin{aligned} \Rightarrow \begin{cases} x = -2z \\ y = t \\ z = s \end{cases} &\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -2s \\ t \\ s \end{bmatrix} \\ &= s \begin{bmatrix} -2 \\ 0 \\ 1 \end{bmatrix} + t \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \quad t, s \in \mathbb{R} \end{aligned}$$

Basis and Dimension

2. Base: For all $(x, y, z) \in \mathcal{U}$ we have $x + 2z = 0 \Rightarrow x = -2z$, a linear equation with three unknowns. This system has infinitely many solutions depending on two parameters.

$$\begin{aligned} \Rightarrow \begin{cases} x = -2z \\ y = t \\ z = s \end{cases} &\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -2s \\ t \\ s \end{bmatrix} \\ &= s \begin{bmatrix} -2 \\ 0 \\ 1 \end{bmatrix} + t \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \quad t, s \in \mathbb{R} \end{aligned}$$

It can be easily seen that the vectors $(-2, 0, 1)$ and $(0, 1, 0)$ are linearly independent and spans \mathcal{U} . Thus, $\mathcal{B} = \{(-2, 0, 1), (0, 1, 0)\}$ is a basis for \mathcal{U} .

Basis and Dimension

2. Base: For all $(x, y, z) \in \mathcal{U}$ we have $x + 2z = 0 \Rightarrow x = -2z$, a linear equation with three unknowns. This system has infinitely many solutions depending on two parameters.

$$\begin{aligned} \Rightarrow \begin{cases} x = -2z \\ y = t \\ z = s \end{cases} &\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -2s \\ t \\ s \end{bmatrix} \\ &= s \begin{bmatrix} -2 \\ 0 \\ 1 \end{bmatrix} + t \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \quad t, s \in \mathbb{R} \end{aligned}$$

It can be easily seen that the vectors $(-2, 0, 1)$ and $(0, 1, 0)$ are linearly independent and spans \mathcal{U} . Thus, $\mathcal{B} = \{(-2, 0, 1), (0, 1, 0)\}$ is a basis for \mathcal{U} .

3. Dimension: $\dim(\mathcal{U}) = 2$.

Example

$$\text{Let } \mathcal{U} = \left\{ \begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix} \mid a, b \in \mathbb{R} \right\} \subset \mathcal{M}_{2 \times 3}(\mathbb{R}).$$

Example

$$\text{Let } \mathcal{U} = \left\{ \begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix} \mid a, b \in \mathbb{R} \right\} \subset \mathcal{M}_{2 \times 3}(\mathbb{R}).$$

1. Show that \mathcal{U} is a subspace of $\mathcal{M}_{2 \times 3}(\mathbb{R})$.

Example

$$\text{Let } \mathcal{U} = \left\{ \begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix} \mid a, b \in \mathbb{R} \right\} \subset \mathcal{M}_{2 \times 3}(\mathbb{R}).$$

1. Show that \mathcal{U} is a subspace of $\mathcal{M}_{2 \times 3}(\mathbb{R})$.
2. Find a basis for \mathcal{U} .

Example

Let $\mathcal{U} = \left\{ \begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix} \mid a, b \in \mathbb{R} \right\} \subset \mathcal{M}_{2 \times 3}(\mathbb{R})$.

1. Show that \mathcal{U} is a subspace of $\mathcal{M}_{2 \times 3}(\mathbb{R})$.
2. Find a basis for \mathcal{U} .
3. $\dim(\mathcal{U}) = ?$

1. Subspace: For all $\begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix}, \begin{pmatrix} x & y & 0 \\ 0 & 0 & x+y \end{pmatrix} \in \mathcal{U}$ and $\alpha, \beta \in \mathbb{R}$

1. Subspace: For all $\begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix}, \begin{pmatrix} x & y & 0 \\ 0 & 0 & x+y \end{pmatrix} \in \mathcal{U}$ and $\alpha, \beta \in \mathbb{R}$

$$\Rightarrow \alpha \begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix} + \beta \begin{pmatrix} x & y & 0 \\ 0 & 0 & x+y \end{pmatrix}$$

1. Subspace: For all $\begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix}, \begin{pmatrix} x & y & 0 \\ 0 & 0 & x+y \end{pmatrix} \in \mathcal{U}$ and $\alpha, \beta \in \mathbb{R}$

$$\begin{aligned} &\Rightarrow \alpha \begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix} + \beta \begin{pmatrix} x & y & 0 \\ 0 & 0 & x+y \end{pmatrix} \\ &= \begin{pmatrix} \alpha a + \beta x & \alpha b + \beta y & 0 \\ 0 & 0 & \alpha(a+b) + \beta(x+y) \end{pmatrix} \in \mathcal{U}. \end{aligned}$$

1. Subspace: For all $\begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix}, \begin{pmatrix} x & y & 0 \\ 0 & 0 & x+y \end{pmatrix} \in \mathcal{U}$ and $\alpha, \beta \in \mathbb{R}$

$$\begin{aligned} &\Rightarrow \alpha \begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix} + \beta \begin{pmatrix} x & y & 0 \\ 0 & 0 & x+y \end{pmatrix} \\ &= \begin{pmatrix} \alpha a + \beta x & \alpha b + \beta y & 0 \\ 0 & 0 & \alpha(a+b) + \beta(x+y) \end{pmatrix} \in \mathcal{U}. \end{aligned}$$

This says that \mathcal{U} is a subspace of $\mathcal{M}_{2 \times 3}(\mathbb{R})$.

2. **Base:** For all $\begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix} \in \mathcal{U},$

2. Base: For all $\begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix} \in \mathcal{U},$

$$\begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix} = a \underbrace{\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{E_1} + b \underbrace{\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{E_2}.$$

2. Base: For all $\begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix} \in \mathcal{U},$

$$\begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix} = a \underbrace{\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{E_1} + b \underbrace{\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{E_2}.$$

The matrices E_1, E_2 are linearly independent and spans \mathcal{U} . Hence, $\mathcal{B} = \{E_1, E_2\}$ is a basis for \mathcal{U} .

2. Base: For all $\begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix} \in \mathcal{U},$

$$\begin{pmatrix} a & b & 0 \\ 0 & 0 & a+b \end{pmatrix} = a \underbrace{\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{E_1} + b \underbrace{\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{E_2}.$$

The matrices E_1, E_2 are linearly independent and spans \mathcal{U} . Hence, $\mathcal{B} = \{E_1, E_2\}$ is a basis for \mathcal{U} .

3. Dimension: $\dim(\mathcal{U}) = 2.$

Coordinates

Theorem

Let V be vector space of dimension n and with the ordered basis $\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$. Then any vector $\vec{w} \in V$ can be expressed uniquely as a linear combination of basis vectors in \mathcal{B} , say

$$\vec{w} = x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n.$$

Theorem

Let V be vector space of dimension n and with the ordered basis $\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$. Then any vector $\vec{w} \in V$ can be expressed uniquely as a linear combination of basis vectors in \mathcal{B} , say

$$\vec{w} = x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n.$$

Note: These n scalars x_1, x_2, \dots, x_n are called the **coordinates** of \vec{w} relative to the basis \mathcal{B} , and they form a vector (x_1, x_2, \dots, x_n) in \mathbb{R}^n called the **coordinate vector** of \vec{w} relative to \mathcal{B} . We denote this vector by $[\vec{w}]_{\mathcal{B}}$, or simply $[\vec{w}]$, when \mathcal{B} is understood. Thus,

$$[\vec{w}]_{\mathcal{B}} = (x_1, x_2, \dots, x_n).$$

Proof.

Let $\mathcal{B} = \{\vec{\mathbf{v}}_1, \vec{\mathbf{v}}_2, \dots, \vec{\mathbf{v}}_n\}$ be an ordered basis of n -space V and $x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n \in \mathbb{R}$.

Proof.

Let $\mathcal{B} = \{\vec{\mathbf{v}}_1, \vec{\mathbf{v}}_2, \dots, \vec{\mathbf{v}}_n\}$ be an ordered basis of n -space V and $x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n \in \mathbb{R}$. Assume that $\vec{\mathbf{w}}$ has two expressions as follows:

$$\vec{\mathbf{w}} = x_1 \vec{\mathbf{v}}_1 + x_2 \vec{\mathbf{v}}_2 + \dots + x_n \vec{\mathbf{v}}_n$$

Proof.

Let $\mathcal{B} = \{\vec{\mathbf{v}}_1, \vec{\mathbf{v}}_2, \dots, \vec{\mathbf{v}}_n\}$ be an ordered basis of n -space V and $x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n \in \mathbb{R}$. Assume that $\vec{\mathbf{w}}$ has two expressions as follows:

$$\vec{\mathbf{w}} = x_1 \vec{\mathbf{v}}_1 + x_2 \vec{\mathbf{v}}_2 + \dots + x_n \vec{\mathbf{v}}_n$$

$$\vec{\mathbf{w}} = y_1 \vec{\mathbf{v}}_1 + y_2 \vec{\mathbf{v}}_2 + \dots + y_n \vec{\mathbf{v}}_n$$

Proof.

Let $\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ be an ordered basis of n -space V and $x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n \in \mathbb{R}$. Assume that \vec{w} has two expressions as follows:

$$\vec{w} = x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n$$

$$\vec{w} = y_1 \vec{v}_1 + y_2 \vec{v}_2 + \dots + y_n \vec{v}_n$$

Since basis elements are linearly independent we have

$$\Rightarrow (x_1 - y_1) \vec{v}_1 + (x_2 - y_2) \vec{v}_2 + \dots + (x_n - y_n) \vec{v}_n = \vec{0}$$

Proof.

Let $\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ be an ordered basis of n -space V and $x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n \in \mathbb{R}$. Assume that \vec{w} has two expressions as follows:

$$\vec{w} = x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n$$

$$\vec{w} = y_1 \vec{v}_1 + y_2 \vec{v}_2 + \dots + y_n \vec{v}_n$$

Since basis elements are linearly independent we have

$$\Rightarrow (x_1 - y_1) \vec{v}_1 + (x_2 - y_2) \vec{v}_2 + \dots + (x_n - y_n) \vec{v}_n = \vec{0}$$

$$\Rightarrow x_1 - y_1 = 0, x_2 - y_2 = 0, \dots, x_n - y_n = 0$$

Proof.

Let $\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ be an ordered basis of n -space V and $x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n \in \mathbb{R}$. Assume that \vec{w} has two expressions as follows:

$$\vec{w} = x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n$$

$$\vec{w} = y_1 \vec{v}_1 + y_2 \vec{v}_2 + \dots + y_n \vec{v}_n$$

Since basis elements are linearly independent we have

$$\Rightarrow (x_1 - y_1) \vec{v}_1 + (x_2 - y_2) \vec{v}_2 + \dots + (x_n - y_n) \vec{v}_n = \vec{0}$$

$$\Rightarrow x_1 - y_1 = 0, x_2 - y_2 = 0, \dots, x_n - y_n = 0$$

$$\Rightarrow x_1 = y_1, x_2 = y_2, \dots, x_n = y_n.$$



Example

Find coordinate vector $[\vec{\mathbf{w}}]_{\mathcal{B}}$ of $\vec{\mathbf{w}} = (1, 2, 3) \in \mathbb{R}^3$ relative to ordered basis $\mathcal{B} = \{(1, 2, 0), (2, 0, 1), (1, 2, 1)\}$.

Example

Find coordinate vector $[\vec{w}]_{\mathcal{B}}$ of $\vec{w} = (1, 2, 3) \in \mathbb{R}^3$ relative to ordered basis $\mathcal{B} = \{(1, 2, 0), (2, 0, 1), (1, 2, 1)\}$. For all $(x, y, z) \in \mathbb{R}^3$

$$(x, y, z) = x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1)$$

Example

Find coordinate vector $[\vec{w}]_{\mathcal{B}}$ of $\vec{w} = (1, 2, 3) \in \mathbb{R}^3$ relative to ordered basis $\mathcal{B} = \{(1, 2, 0), (2, 0, 1), (1, 2, 1)\}$. For all $(x, y, z) \in \mathbb{R}^3$

$$(x, y, z) = x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1)$$

$$x_1 + 2x_2 + x_3 = x$$

$$2x_1 + 2x_3 = y$$

$$x_2 + x_3 = z$$

Coordinates

Example

Find coordinate vector $[\vec{w}]_{\mathcal{B}}$ of $\vec{w} = (1, 2, 3) \in \mathbb{R}^3$ relative to ordered basis $\mathcal{B} = \{(1, 2, 0), (2, 0, 1), (1, 2, 1)\}$. For all $(x, y, z) \in \mathbb{R}^3$

$$\begin{aligned}(x, y, z) &= x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1) \\ \begin{aligned} x_1 + 2x_2 + x_3 &= x \\ 2x_1 + 2x_3 &= y \\ x_2 + x_3 &= z \end{aligned} &\Rightarrow \left(\begin{array}{ccc|c} 1 & 2 & 1 & x \\ 2 & 0 & 2 & y \\ 0 & 1 & 1 & z \end{array} \right)\end{aligned}$$

Coordinates

Example

Find coordinate vector $[\vec{w}]_{\mathcal{B}}$ of $\vec{w} = (1, 2, 3) \in \mathbb{R}^3$ relative to ordered basis $\mathcal{B} = \{(1, 2, 0), (2, 0, 1), (1, 2, 1)\}$. For all $(x, y, z) \in \mathbb{R}^3$

$$\begin{aligned}(x, y, z) &= x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1) \\ \begin{aligned} x_1 + 2x_2 + x_3 &= x \\ 2x_1 + 2x_3 &= y \\ x_2 + x_3 &= z \end{aligned} &\Rightarrow \left(\begin{array}{ccc|c} 1 & 2 & 1 & x \\ 2 & 0 & 2 & y \\ 0 & 1 & 1 & z \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 0 & 0 & \frac{2x+y-4z}{4} \\ 0 & 1 & 0 & \frac{2x-y}{4} \\ 0 & 0 & 1 & \frac{-2x+y+4z}{4} \end{array} \right).\end{aligned}$$

Coordinates

Example

Find coordinate vector $[\vec{w}]_{\mathcal{B}}$ of $\vec{w} = (1, 2, 3) \in \mathbb{R}^3$ relative to ordered basis $\mathcal{B} = \{(1, 2, 0), (2, 0, 1), (1, 2, 1)\}$. For all $(x, y, z) \in \mathbb{R}^3$

$$\begin{aligned} (x, y, z) &= x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1) \\ \begin{aligned} x_1 + 2x_2 + x_3 &= x \\ 2x_1 + 2x_3 &= y \\ x_2 + x_3 &= z \end{aligned} &\Rightarrow \left(\begin{array}{ccc|c} 1 & 2 & 1 & x \\ 2 & 0 & 2 & y \\ 0 & 1 & 1 & z \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 0 & 0 & \frac{2x+y-4z}{4} \\ 0 & 1 & 0 & \frac{2x-y}{4} \\ 0 & 0 & 1 & \frac{-2x+y+4z}{4} \end{array} \right). \end{aligned}$$

Then we have $x_1 = \frac{2x+y-4z}{4}$, $x_2 = \frac{2x-y}{4}$, $x_3 = \frac{-2x+y+4z}{4}$

Coordinates

Example

Find coordinate vector $[\vec{w}]_{\mathcal{B}}$ of $\vec{w} = (1, 2, 3) \in \mathbb{R}^3$ relative to ordered basis $\mathcal{B} = \{(1, 2, 0), (2, 0, 1), (1, 2, 1)\}$. For all $(x, y, z) \in \mathbb{R}^3$

$$\begin{aligned}(x, y, z) &= x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(1, 2, 1) \\ \begin{aligned} x_1 + 2x_2 + x_3 &= x \\ 2x_1 + 2x_3 &= y \\ x_2 + x_3 &= z \end{aligned} &\Rightarrow \left(\begin{array}{ccc|c} 1 & 2 & 1 & x \\ 2 & 0 & 2 & y \\ 0 & 1 & 1 & z \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 0 & 0 & \frac{2x+y-4z}{4} \\ 0 & 1 & 0 & \frac{2x-y}{4} \\ 0 & 0 & 1 & \frac{-2x+y+4z}{4} \end{array} \right).\end{aligned}$$

Then we have $x_1 = \frac{2x+y-4z}{4}$, $x_2 = \frac{2x-y}{4}$, $x_3 = \frac{-2x+y+4z}{4}$ and hence,

$$(1, 2, 3) = -2 \cdot (1, 2, 0) + 0 \cdot (2, 0, 1) + 3(1, 2, 1).$$

?