



# MAT1320-Linear Algebra

## Lecture Notes

Determinants and Properties

---

Mehmet E. KÖROĞLU  
Summer 2020

YILDIZ TECHNICAL UNIVERSITY, DEPARTMENT OF MATHEMATICS  
*mkoroglu@yildiz.edu.tr*

# Table of contents

1. Determinants
2. Minors and Cofactors
3. Properties of Determinants

# Determinants

---

# Determinants

Let  $\mathbf{A} = [a_{ij}]_{n \times n}$  be a square matrix of order  $n$ . Then the **determinant** of  $\mathbf{A}$  is denoted by  $\det(\mathbf{A})$  or  $|\mathbf{A}|$  and defined as follows:

# Determinants

Let  $\mathbf{A} = [a_{ij}]_{n \times n}$  be a square matrix of order  $n$ . Then the **determinant** of  $\mathbf{A}$  is denoted by  $\det(\mathbf{A})$  or  $|\mathbf{A}|$  and defined as follows:

- For  $n = 1$ ,  $A = [a]_{1 \times 1}$  and  $\det(A) = a$ .

# Determinants

Let  $\mathbf{A} = [a_{ij}]_{n \times n}$  be a square matrix of order  $n$ . Then the **determinant** of  $\mathbf{A}$  is denoted by  $\det(\mathbf{A})$  or  $|\mathbf{A}|$  and defined as follows:

- For  $n = 1$ ,  $A = [a]_{1 \times 1}$  and  $\det(A) = a$ .

- For  $n = 2$ ,  $A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}_{2 \times 2}$  and

$$\det(A) = a_{11}a_{22} - a_{12}a_{21}.$$

# Determinants

- For  $n = 3$ ,  $A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}_{3 \times 3}$  and

$$\det(A) = a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$

# Determinants

▪ For  $n = 3$ ,  $A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}_{3 \times 3}$  and

$$\det(A) = a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$

$$\begin{aligned} &= a_{11} (a_{22}a_{33} - a_{23}a_{32}) - a_{12} (a_{21}a_{33} - a_{23}a_{31}) \\ &\quad + a_{13} (a_{21}a_{32} - a_{22}a_{31}) \end{aligned}$$



# Determinants

- For  $n = 3$ ,  $A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}_{3 \times 3}$  and

$$\det(A) = a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$

$$\begin{aligned} &= a_{11} (a_{22}a_{33} - a_{23}a_{32}) - a_{12} (a_{21}a_{33} - a_{23}a_{31}) \\ &\quad + a_{13} (a_{21}a_{32} - a_{22}a_{31}) \end{aligned}$$

In order to generalize this concept for  $n > 3$ , we need to give definition of the minors and cofactors.

# Minors and Cofactors

---

# Minors and Cofactors

## Definition

Given an  $n \times n$  matrix  $\mathbf{A}$ , the  $(i, j)$ -th minor, denoted  $M_{ij}$ , is the determinant of the  $(n - 1) \times (n - 1)$  matrix obtained from  $\mathbf{A}$  by deleting the  $i$ -th row and the  $j$ -th column.

# Minors and Cofactors

## Definition

Given an  $n \times n$  matrix  $\mathbf{A}$ , the  $(i, j)$ -th minor, denoted  $M_{ij}$ , is the determinant of the  $(n - 1) \times (n - 1)$  matrix obtained from  $\mathbf{A}$  by deleting the  $i$ -th row and the  $j$ -th column. Similarly, the  $(i, j)$ -th cofactor of  $\mathbf{A}$  is denoted by  $A_{ij}$  is defined in terms of the minor by  $A_{ij} = (-1)^{i+j} M_{ij}$ .

# Minors and Cofactors

## Definition

Given an  $n \times n$  matrix  $\mathbf{A}$ , the  $(i, j)$ -th minor, denoted  $M_{ij}$ , is the determinant of the  $(n - 1) \times (n - 1)$  matrix obtained from  $\mathbf{A}$  by deleting the  $i$ -th row and the  $j$ -th column. Similarly, the  $(i, j)$ -th cofactor of  $\mathbf{A}$  is denoted by  $A_{ij}$  is defined in terms of the minor by  $A_{ij} = (-1)^{i+j} M_{ij}$ .

## Example

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}$$

we have

$$M_{23} = \det \begin{pmatrix} 1 & 2 \\ 7 & 8 \end{pmatrix} = 8 - 14 = -6$$

We also find  $A_{23} = (-1)^{2+3}(-6) = 6$

# Laplace Expansion

**Algorithm (Laplace Expansion):** To compute the determinant of a square  $\mathbf{A}$  matrix, do the following.

# Laplace Expansion

**Algorithm (Laplace Expansion):** To compute the determinant of a square  $\mathbf{A}$  matrix, do the following.

1. Choose any row or column of  $\mathbf{A}$ .

# Laplace Expansion

**Algorithm (Laplace Expansion):** To compute the determinant of a square  $\mathbf{A}$  matrix, do the following.

1. Choose any row or column of  $\mathbf{A}$ .
2. For each element  $a_{ij}$  of this row or column, compute the associated cofactor  $A_{ij}$ .



# Laplace Expansion

**Algorithm (Laplace Expansion):** To compute the determinant of a square  $\mathbf{A}$  matrix, do the following.

1. Choose any row or column of  $\mathbf{A}$ .
2. For each element  $a_{ij}$  of this row or column, compute the associated cofactor  $A_{ij}$ .
3. Multiply each cofactor by the associated matrix entry  $a_{ij}$ .

# Laplace Expansion

**Algorithm (Laplace Expansion):** To compute the determinant of a square  $\mathbf{A}$  matrix, do the following.

1. Choose any row or column of  $\mathbf{A}$ .
2. For each element  $a_{ij}$  of this row or column, compute the associated cofactor  $A_{ij}$ .
3. Multiply each cofactor by the associated matrix entry  $a_{ij}$ .
4. The sum of these products is  $\det(\mathbf{A})$ .

# Laplace Expansion

**Algorithm (Laplace Expansion):** To compute the determinant of a square  $\mathbf{A}$  matrix, do the following.

1. Choose any row or column of  $\mathbf{A}$ .
2. For each element  $a_{ij}$  of this row or column, compute the associated cofactor  $A_{ij}$ .
3. Multiply each cofactor by the associated matrix entry  $a_{ij}$ .
4. The sum of these products is  $\det(\mathbf{A})$ .

Let  $\mathbf{A} = [a_{ij}]_{n \times n}$  be a square matrix of order  $n$ .

# Laplace Expansion

**Algorithm (Laplace Expansion):** To compute the determinant of a square  $\mathbf{A}$  matrix, do the following.

1. Choose any row or column of  $\mathbf{A}$ .
2. For each element  $a_{ij}$  of this row or column, compute the associated cofactor  $A_{ij}$ .
3. Multiply each cofactor by the associated matrix entry  $a_{ij}$ .
4. The sum of these products is  $\det(\mathbf{A})$ .

Let  $\mathbf{A} = [a_{ij}]_{n \times n}$  be a square matrix of order  $n$ . Then

$$\det(\mathbf{A}) = \sum_{k=1}^n a_{ik} A_{ik} \text{ (LE along } i\text{-th row)}$$

# Laplace Expansion

**Algorithm (Laplace Expansion):** To compute the determinant of a square  $\mathbf{A}$  matrix, do the following.

1. Choose any row or column of  $\mathbf{A}$ .
2. For each element  $a_{ij}$  of this row or column, compute the associated cofactor  $A_{ij}$ .
3. Multiply each cofactor by the associated matrix entry  $a_{ij}$ .
4. The sum of these products is  $\det(\mathbf{A})$ .

Let  $\mathbf{A} = [a_{ij}]_{n \times n}$  be a square matrix of order  $n$ . Then

$$\det(\mathbf{A}) = \sum_{k=1}^n a_{ik} A_{ik} \text{ (LE along } i\text{-th row)}$$

$$\det(\mathbf{A}) = \sum_{k=1}^n a_{kj} A_{kj} \text{ (LE along } j\text{-th column)}$$

# Laplace Expansion

**Note:** For  $i \neq j$  we have

$$\sum_{k=1}^n a_{ik} A_{jk} = 0$$

# Laplace Expansion

**Note:** For  $i \neq j$  we have

$$\sum_{k=1}^n a_{ik}A_{jk} = 0 \text{ and } \sum_{k=1}^n a_{ki}A_{kj} = 0$$

# Laplace Expansion

**Note:** For  $i \neq j$  we have

$$\sum_{k=1}^n a_{ik}A_{jk} = 0 \text{ and } \sum_{k=1}^n a_{ki}A_{kj} = 0$$

## Example

We find the determinant of

$$A = \begin{pmatrix} 2 & 1 & 3 \\ -1 & 2 & 1 \\ -2 & 2 & 3 \end{pmatrix}.$$



# Laplace Expansion

## Example (1)

We make the arbitrary choice to expand along the first row. We compute the cofactors as

# Laplace Expansion

## Example (1)

We make the arbitrary choice to expand along the first row. We compute the cofactors as

$$A_{11} = (-1)^{1+1} \det \begin{pmatrix} 2 & 1 \\ 2 & 3 \end{pmatrix} = 4$$

# Laplace Expansion

## Example (1)

We make the arbitrary choice to expand along the first row. We compute the cofactors as

$$A_{11} = (-1)^{1+1} \det \begin{pmatrix} 2 & 1 \\ 2 & 3 \end{pmatrix} = 4$$

$$A_{12} = (-1)^{1+2} \det \begin{pmatrix} -1 & 1 \\ -2 & 3 \end{pmatrix} = 1$$

# Laplace Expansion

## Example (1)

We make the arbitrary choice to expand along the first row. We compute the cofactors as

$$A_{11} = (-1)^{1+1} \det \begin{pmatrix} 2 & 1 \\ 2 & 3 \end{pmatrix} = 4$$

$$A_{12} = (-1)^{1+2} \det \begin{pmatrix} -1 & 1 \\ -2 & 3 \end{pmatrix} = 1$$

$$A_{13} = (-1)^{1+3} \det \begin{pmatrix} -1 & 2 \\ -2 & 2 \end{pmatrix} = 2$$

# Laplace Expansion

## Example (1)

We make the arbitrary choice to expand along the first row. We compute the cofactors as

$$A_{11} = (-1)^{1+1} \det \begin{pmatrix} 2 & 1 \\ 2 & 3 \end{pmatrix} = 4$$

$$A_{12} = (-1)^{1+2} \det \begin{pmatrix} -1 & 1 \\ -2 & 3 \end{pmatrix} = 1$$

$$A_{13} = (-1)^{1+3} \det \begin{pmatrix} -1 & 2 \\ -2 & 2 \end{pmatrix} = 2$$

Thus,

$$\det(A) = a_{11}A_{11} + a_{12}A_{12} + a_{13}A_{13} = 2(4) + (1) + 3(2) = 15$$

## Example (1-cont.)

$$A_{11} = (-1)^{1+1} \det \begin{pmatrix} 2 & 1 \\ 2 & 3 \end{pmatrix} = 4$$

# Laplace Expansion

## Example (1-cont.)

$$A_{11} = (-1)^{1+1} \det \begin{pmatrix} 2 & 1 \\ 2 & 3 \end{pmatrix} = 4$$

$$A_{12} = (-1)^{1+2} \det \begin{pmatrix} -1 & 1 \\ -2 & 3 \end{pmatrix} = 1$$

# Laplace Expansion

## Example (1-cont.)

$$A_{11} = (-1)^{1+1} \det \begin{pmatrix} 2 & 1 \\ 2 & 3 \end{pmatrix} = 4$$

$$A_{12} = (-1)^{1+2} \det \begin{pmatrix} -1 & 1 \\ -2 & 3 \end{pmatrix} = 1$$

$$A_{13} = (-1)^{1+3} \det \begin{pmatrix} -1 & 2 \\ -2 & 2 \end{pmatrix} = 2$$



# Laplace Expansion

## Example (1-cont.)

$$A_{11} = (-1)^{1+1} \det \begin{pmatrix} 2 & 1 \\ 2 & 3 \end{pmatrix} = 4$$

$$A_{12} = (-1)^{1+2} \det \begin{pmatrix} -1 & 1 \\ -2 & 3 \end{pmatrix} = 1$$

$$A_{13} = (-1)^{1+3} \det \begin{pmatrix} -1 & 2 \\ -2 & 2 \end{pmatrix} = 2$$

Notice that

$$a_{21}A_{11} + a_{22}A_{12} + a_{23}A_{13} = (-1)(4) + 2(1) + 1(2) = 0$$

# Laplace Expansion

## Example (1-cont.)

$$A_{11} = (-1)^{1+1} \det \begin{pmatrix} 2 & 1 \\ 2 & 3 \end{pmatrix} = 4$$

$$A_{12} = (-1)^{1+2} \det \begin{pmatrix} -1 & 1 \\ -2 & 3 \end{pmatrix} = 1$$

$$A_{13} = (-1)^{1+3} \det \begin{pmatrix} -1 & 2 \\ -2 & 2 \end{pmatrix} = 2$$

Notice that

$$a_{21}A_{11} + a_{22}A_{12} + a_{23}A_{13} = (-1)(4) + 2(1) + 1(2) = 0$$

$$a_{31}A_{11} + a_{32}A_{12} + a_{33}A_{13} = (-2)(4) + 2(1) + 3(2) = 0$$

# Laplace Expansion

## Example (2)

We make the arbitrary choice to expand along the second column.

We compute the cofactors as:

# Laplace Expansion

## Example (2)

We make the arbitrary choice to expand along the second column.

We compute the cofactors as:

$$A_{12} = (-1)^{1+2} \det \begin{pmatrix} -1 & 1 \\ -2 & 3 \end{pmatrix} = 1$$

# Laplace Expansion

## Example (2)

We make the arbitrary choice to expand along the second column.

We compute the cofactors as:

$$A_{12} = (-1)^{1+2} \det \begin{pmatrix} -1 & 1 \\ -2 & 3 \end{pmatrix} = 1$$

$$A_{22} = (-1)^{2+2} \det \begin{pmatrix} 2 & 3 \\ -2 & 3 \end{pmatrix} = 12$$

# Laplace Expansion

## Example (2)

We make the arbitrary choice to expand along the second column.

We compute the cofactors as:

$$A_{12} = (-1)^{1+2} \det \begin{pmatrix} -1 & 1 \\ -2 & 3 \end{pmatrix} = 1$$

$$A_{22} = (-1)^{2+2} \det \begin{pmatrix} 2 & 3 \\ -2 & 3 \end{pmatrix} = 12$$

$$A_{32} = (-1)^{3+2} \det \begin{pmatrix} 2 & 3 \\ -1 & 1 \end{pmatrix} = -5$$

# Laplace Expansion

## Example (2)

We make the arbitrary choice to expand along the second column.

We compute the cofactors as:

$$A_{12} = (-1)^{1+2} \det \begin{pmatrix} -1 & 1 \\ -2 & 3 \end{pmatrix} = 1$$

$$A_{22} = (-1)^{2+2} \det \begin{pmatrix} 2 & 3 \\ -2 & 3 \end{pmatrix} = 12$$

$$A_{32} = (-1)^{3+2} \det \begin{pmatrix} 2 & 3 \\ -1 & 1 \end{pmatrix} = -5$$

Thus,

$$\det(A) = a_{12}A_{12} + a_{22}A_{22} + a_{32}A_{32} = 1(1) + 2(12) + 2(-5) = 15$$

## Example (2-cont.)

$$A_{12} = (-1)^{1+2} \det \begin{pmatrix} -1 & 1 \\ -2 & 3 \end{pmatrix} = 1$$



# Laplace Expansion

## Example (2-cont.)

$$A_{12} = (-1)^{1+2} \det \begin{pmatrix} -1 & 1 \\ -2 & 3 \end{pmatrix} = 1$$

$$A_{22} = (-1)^{2+2} \det \begin{pmatrix} 2 & 3 \\ -2 & 3 \end{pmatrix} = 12$$

# Laplace Expansion

## Example (2-cont.)

$$A_{12} = (-1)^{1+2} \det \begin{pmatrix} -1 & 1 \\ -2 & 3 \end{pmatrix} = 1$$

$$A_{22} = (-1)^{2+2} \det \begin{pmatrix} 2 & 3 \\ -2 & 3 \end{pmatrix} = 12$$

$$A_{32} = (-1)^{3+2} \det \begin{pmatrix} 2 & 3 \\ -1 & 1 \end{pmatrix} = -5$$

# Laplace Expansion

## Example (2-cont.)

$$A_{12} = (-1)^{1+2} \det \begin{pmatrix} -1 & 1 \\ -2 & 3 \end{pmatrix} = 1$$

$$A_{22} = (-1)^{2+2} \det \begin{pmatrix} 2 & 3 \\ -2 & 3 \end{pmatrix} = 12$$

$$A_{32} = (-1)^{3+2} \det \begin{pmatrix} 2 & 3 \\ -1 & 1 \end{pmatrix} = -5$$

Note that

$$a_{11}A_{12} + a_{21}A_{22} + a_{31}A_{32} = 2(1) + (-1)(12) + (-2)(-5) = 0$$

# Laplace Expansion

## Example (2-cont.)

$$A_{12} = (-1)^{1+2} \det \begin{pmatrix} -1 & 1 \\ -2 & 3 \end{pmatrix} = 1$$

$$A_{22} = (-1)^{2+2} \det \begin{pmatrix} 2 & 3 \\ -2 & 3 \end{pmatrix} = 12$$

$$A_{32} = (-1)^{3+2} \det \begin{pmatrix} 2 & 3 \\ -1 & 1 \end{pmatrix} = -5$$

Note that

$$a_{11}A_{12} + a_{21}A_{22} + a_{31}A_{32} = 2(1) + (-1)(12) + (-2)(-5) = 0$$

$$a_{13}A_{12} + a_{23}A_{22} + a_{33}A_{32} = 3(1) + 1(12) + 3(-5) = 0$$

# Properties of Determinants

---

# Properties of Determinants

- $\det I = 1$

# Properties of Determinants

- $\det I = 1$
- If you exchange two rows or columns of a matrix, you reverse the sign of its determinant from positive to negative or from negative to positive.

# Properties of Determinants

- $\det I = 1$
- If you exchange two rows or columns of a matrix, you reverse the sign of its determinant from positive to negative or from negative to positive. Property 1 tells us that  $\begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = 1$ .



# Properties of Determinants

- $\det I = 1$
- If you exchange two rows or columns of a matrix, you reverse the sign of its determinant from positive to negative or from negative to positive. Property 1 tells us that  $\begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = 1$ .

Property 2 tells us that  $\begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} = -1$

# Properties of Determinants

- $\det I = 1$
- If you exchange two rows or columns of a matrix, you reverse the sign of its determinant from positive to negative or from negative to positive. Property 1 tells us that  $\begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = 1$ .

Property 2 tells us that  $\begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} = -1$

- If we multiply one row of a matrix by  $t$ , the determinant is multiplied by  $t$ .

# Properties of Determinants

- $\det I = 1$
- If you exchange two rows or columns of a matrix, you reverse the sign of its determinant from positive to negative or from negative to positive. Property 1 tells us that  $\begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = 1$ .

Property 2 tells us that  $\begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} = -1$

- If we multiply one row of a matrix by  $t$ , the determinant is multiplied by  $t$ . For example;  $\begin{vmatrix} ta & tb \\ c & d \end{vmatrix} = t \begin{vmatrix} a & b \\ c & d \end{vmatrix}$ .

# Properties of Determinants

- $\det I = 1$
- If you exchange two rows or columns of a matrix, you reverse the sign of its determinant from positive to negative or from negative to positive. Property 1 tells us that  $\begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = 1$ .

Property 2 tells us that  $\begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} = -1$

- If we multiply one row of a matrix by  $t$ , the determinant is multiplied by  $t$ . For example;  $\begin{vmatrix} ta & tb \\ c & d \end{vmatrix} = t \begin{vmatrix} a & b \\ c & d \end{vmatrix}$ .
- The determinant behaves like a linear function on the rows of the matrix:

# Properties of Determinants

- $\det I = 1$
- If you exchange two rows or columns of a matrix, you reverse the sign of its determinant from positive to negative or from negative to positive. Property 1 tells us that  $\begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = 1$ .

Property 2 tells us that  $\begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} = -1$

- If we multiply one row of a matrix by  $t$ , the determinant is multiplied by  $t$ . For example;  $\begin{vmatrix} ta & tb \\ c & d \end{vmatrix} = t \begin{vmatrix} a & b \\ c & d \end{vmatrix}$ .
- The determinant behaves like a linear function on the rows of the matrix:

$$\begin{vmatrix} a + a' & b + b' \\ c & d \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix} + \begin{vmatrix} a' & b' \\ c & d \end{vmatrix}.$$

# Properties of Determinants

- If two rows or columns of a matrix are equal, its determinant is zero.

# Properties of Determinants

- If two rows or columns of a matrix are equal, its determinant is zero.
- If  $i \neq j$ , subtracting  $t$  times row  $i$  from row  $j$  doesn't change the determinant.

# Properties of Determinants

- If two rows or columns of a matrix are equal, its determinant is zero.
- If  $i \neq j$ , subtracting  $t$  times row  $i$  from row  $j$  doesn't change the determinant. In two dimensions, this argument looks like:

$$\begin{vmatrix} a & b \\ c - ta & d - tb \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix} - \begin{vmatrix} a & b \\ ta & tb \end{vmatrix}$$



# Properties of Determinants

- If two rows or columns of a matrix are equal, its determinant is zero.
- If  $i \neq j$ , subtracting  $t$  times row  $i$  from row  $j$  doesn't change the determinant. In two dimensions, this argument looks like:

$$\begin{vmatrix} a & b \\ c - ta & d - tb \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix} - \begin{vmatrix} a & b \\ ta & tb \end{vmatrix}$$
$$= \begin{vmatrix} a & b \\ c & d \end{vmatrix} - t \begin{vmatrix} a & b \\ a & b \end{vmatrix}$$

# Properties of Determinants

- If two rows or columns of a matrix are equal, its determinant is zero.
- If  $i \neq j$ , subtracting  $t$  times row  $i$  from row  $j$  doesn't change the determinant. In two dimensions, this argument looks like:

$$\begin{vmatrix} a & b \\ c - ta & d - tb \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix} - \begin{vmatrix} a & b \\ ta & tb \end{vmatrix} \\ = \begin{vmatrix} a & b \\ c & d \end{vmatrix} - t \begin{vmatrix} a & b \\ a & b \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix}$$

# Properties of Determinants

- If two rows or columns of a matrix are equal, its determinant is zero.
- If  $i \neq j$ , subtracting  $t$  times row  $i$  from row  $j$  doesn't change the determinant. In two dimensions, this argument looks like:

$$\begin{vmatrix} a & b \\ c - ta & d - tb \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix} - \begin{vmatrix} a & b \\ ta & tb \end{vmatrix} \\ = \begin{vmatrix} a & b \\ c & d \end{vmatrix} - t \begin{vmatrix} a & b \\ a & b \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix}$$

- If  $A$  has a row or column that is all zeros, then  $\det(A) = 0$ .

# Properties of Determinants

- If two rows or columns of a matrix are equal, its determinant is zero.
- If  $i \neq j$ , subtracting  $t$  times row  $i$  from row  $j$  doesn't change the determinant. In two dimensions, this argument looks like:

$$\begin{vmatrix} a & b \\ c - ta & d - tb \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix} - \begin{vmatrix} a & b \\ ta & tb \end{vmatrix} \\ = \begin{vmatrix} a & b \\ c & d \end{vmatrix} - t \begin{vmatrix} a & b \\ a & b \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix}$$

- If  $A$  has a row or column that is all zeros, then  $\det(A) = 0$ .
- The determinant of a triangular matrix is the product of the diagonal entries (pivots)  $d_{11}, d_{22}, \dots, d_{nn}$ .

# Properties of Determinants

- $\det (AB) = \det (A) \det (B) .$

# Properties of Determinants

- $\det (AB) = \det (A) \det (B) .$
- $\det (A^{-1}) = \frac{1}{\det(A)} .$

# Properties of Determinants

- $\det (AB) = \det (A) \det (B)$ .
- $\det (A^{-1}) = \frac{1}{\det(A)}$ . Because  $A^{-1}A = I$ .

# Properties of Determinants

- $\det(AB) = \det(A) \det(B)$ .
- $\det(A^{-1}) = \frac{1}{\det(A)}$ . Because  $A^{-1}A = I$ . Note that if  $A$  is singular then  $A^{-1}$  does not exist and  $\det(A^{-1})$  is undefined.



# Properties of Determinants

- $\det (AB) = \det (A) \det (B)$ .
- $\det (A^{-1}) = \frac{1}{\det(A)}$ . Because  $A^{-1}A = I$ . Note that if  $A$  is singular then  $A^{-1}$  does not exist and  $\det (A^{-1})$  is undefined.
- $\det (A^2) = (\det (A))^2$

# Properties of Determinants

- $\det (AB) = \det (A) \det (B)$ .
- $\det (A^{-1}) = \frac{1}{\det(A)}$ . Because  $A^{-1}A = I$ . Note that if  $A$  is singular then  $A^{-1}$  does not exist and  $\det (A^{-1})$  is undefined.
- $\det (A^2) = (\det (A))^2$  and  $\det (2A) = 2^n \det (A)$ .

# Properties of Determinants

- $\det (AB) = \det (A) \det (B)$ .
- $\det (A^{-1}) = \frac{1}{\det(A)}$ . Because  $A^{-1}A = I$ . Note that if  $A$  is singular then  $A^{-1}$  does not exist and  $\det (A^{-1})$  is undefined.
- $\det (A^2) = (\det (A))^2$  and  $\det (2A) = 2^n \det (A)$ . We assume  $A$  is an  $n \times n$  square matrix.

# Properties of Determinants

- $\det (AB) = \det (A) \det (B)$ .
- $\det (A^{-1}) = \frac{1}{\det(A)}$ . Because  $A^{-1}A = I$ . Note that if  $A$  is singular then  $A^{-1}$  does not exist and  $\det (A^{-1})$  is undefined.
- $\det (A^2) = (\det (A))^2$  and  $\det (2A) = 2^n \det (A)$ . We assume  $A$  is an  $n \times n$  square matrix.
- $\det (A^T) = \det (A)$ .

# Properties of Determinants

- $\det (AB) = \det (A) \det (B)$ .
- $\det (A^{-1}) = \frac{1}{\det(A)}$ . Because  $A^{-1}A = I$ . Note that if  $A$  is singular then  $A^{-1}$  does not exist and  $\det (A^{-1})$  is undefined.
- $\det (A^2) = (\det (A))^2$  and  $\det (2A) = 2^n \det (A)$ . We assume  $A$  is an  $n \times n$  square matrix.
- $\det (A^T) = \det (A)$ .

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = \begin{vmatrix} a & c \\ b & d \end{vmatrix} = ad - bc$$

# Properties of Determinants

## Example

By using properties of determinants factorize the determinant

$$\begin{vmatrix} x+y & z & t \\ z & x+y & t \\ z & t & x+y \end{vmatrix}.$$

# Properties of Determinants

## Example

By using properties of determinants factorize the determinant

$$\begin{vmatrix} x+y & z & t \\ z & x+y & t \\ z & t & x+y \end{vmatrix}.$$

$$\Rightarrow \begin{vmatrix} x+y & z & t \\ z & x+y & t \\ z & t & x+y \end{vmatrix}$$

# Properties of Determinants

## Example

By using properties of determinants factorize the determinant

$$\begin{vmatrix} x+y & z & t \\ z & x+y & t \\ z & t & x+y \end{vmatrix}.$$

$$\Rightarrow \begin{vmatrix} x+y & z & t \\ z & x+y & t \\ z & t & x+y \end{vmatrix} = \begin{vmatrix} x+y+z+t & z & t \\ x+y+z+t & x+y & t \\ x+y+z+t & t & x+y \end{vmatrix}$$



# Properties of Determinants

## Example

By using properties of determinants factorize the determinant

$$\begin{vmatrix} x+y & z & t \\ z & x+y & t \\ z & t & x+y \end{vmatrix}.$$

$$\Rightarrow \begin{vmatrix} x+y & z & t \\ z & x+y & t \\ z & t & x+y \end{vmatrix} = \begin{vmatrix} x+y+z+t & z & t \\ x+y+z+t & x+y & t \\ x+y+z+t & t & x+y \end{vmatrix}$$

$$= \underbrace{(x+y+z+t)}_r \begin{vmatrix} 1 & z & t \\ 1 & x+y & t \\ 1 & t & x+y \end{vmatrix}$$

# Properties of Determinants

## Example

By using properties of determinants factorize the determinant

$$\begin{vmatrix} x+y & z & t \\ z & x+y & t \\ z & t & x+y \end{vmatrix}.$$

$$\Rightarrow \begin{vmatrix} x+y & z & t \\ z & x+y & t \\ z & t & x+y \end{vmatrix} = \begin{vmatrix} x+y+z+t & z & t \\ x+y+z+t & x+y & t \\ x+y+z+t & t & x+y \end{vmatrix}$$

$$= \underbrace{(x+y+z+t)}_r \begin{vmatrix} 1 & z & t \\ 1 & x+y & t \\ 1 & t & x+y \end{vmatrix} = r \begin{vmatrix} 1 & z & t \\ 0 & x+y-z & 0 \\ 0 & t-z & x+y-t \end{vmatrix}$$

# Properties of Determinants

## Example

By using properties of determinants factorize the determinant

$$\begin{vmatrix} x+y & z & t \\ z & x+y & t \\ z & t & x+y \end{vmatrix}.$$

$$\Rightarrow \begin{vmatrix} x+y & z & t \\ z & x+y & t \\ z & t & x+y \end{vmatrix} = \begin{vmatrix} x+y+z+t & z & t \\ x+y+z+t & x+y & t \\ x+y+z+t & t & x+y \end{vmatrix}$$

$$= \underbrace{(x+y+z+t)}_r \begin{vmatrix} 1 & z & t \\ 1 & x+y & t \\ 1 & t & x+y \end{vmatrix} = r \begin{vmatrix} 1 & z & t \\ 0 & x+y-z & 0 \\ 0 & t-z & x+y-t \end{vmatrix}$$

$$= r \begin{vmatrix} x+y-z & 0 \\ t-z & x+y-t \end{vmatrix}$$

# Properties of Determinants

## Example

By using properties of determinants factorize the determinant

$$\begin{vmatrix} x+y & z & t \\ z & x+y & t \\ z & t & x+y \end{vmatrix}.$$

$$\Rightarrow \begin{vmatrix} x+y & z & t \\ z & x+y & t \\ z & t & x+y \end{vmatrix} = \begin{vmatrix} x+y+z+t & z & t \\ x+y+z+t & x+y & t \\ x+y+z+t & t & x+y \end{vmatrix}$$

$$= \underbrace{(x+y+z+t)}_r \begin{vmatrix} 1 & z & t \\ 1 & x+y & t \\ 1 & t & x+y \end{vmatrix} = r \begin{vmatrix} 1 & z & t \\ 0 & x+y-z & 0 \\ 0 & t-z & x+y-t \end{vmatrix}$$

$$= r \begin{vmatrix} x+y-z & 0 \\ t-z & x+y-t \end{vmatrix}$$

$$= (x+y+z+t)(x+y-z)(x+y-t).$$

### Example

By using properties of determinants find the determinant of the

matrix  $A = \begin{pmatrix} 2 & 1 & 3 \\ -1 & 2 & 1 \\ -2 & 2 & 3 \end{pmatrix}$ .

### Example

By using properties of determinants find the determinant of the

$$\text{matrix } A = \begin{pmatrix} 2 & 1 & 3 \\ -1 & 2 & 1 \\ -2 & 2 & 3 \end{pmatrix}.$$

$$\left| \begin{array}{ccc} 2 & 1 & 3 \\ -1 & 2 & 1 \\ -2 & 2 & 3 \end{array} \right| \begin{array}{l} r_1 \leftrightarrow r_2 \\ -r_1 \rightarrow r_1 \\ = \end{array}$$

### Example

By using properties of determinants find the determinant of the

$$\text{matrix } A = \begin{pmatrix} 2 & 1 & 3 \\ -1 & 2 & 1 \\ -2 & 2 & 3 \end{pmatrix}.$$

$$\begin{vmatrix} 2 & 1 & 3 \\ -1 & 2 & 1 \\ -2 & 2 & 3 \end{vmatrix} \begin{array}{l} r_1 \leftrightarrow r_2 \\ -r_1 \rightarrow r_1 \\ = \end{array} \begin{vmatrix} 1 & -2 & -1 \\ 2 & 1 & 3 \\ -2 & 2 & 3 \end{vmatrix} \begin{array}{l} -2r_1 + r_2 \rightarrow r_2 \\ 2r_1 + r_3 \rightarrow r_3 \\ = \end{array}$$

### Example

By using properties of determinants find the determinant of the

$$\text{matrix } A = \begin{pmatrix} 2 & 1 & 3 \\ -1 & 2 & 1 \\ -2 & 2 & 3 \end{pmatrix}.$$

$$\begin{array}{ccc} \left| \begin{array}{ccc} 2 & 1 & 3 \\ -1 & 2 & 1 \\ -2 & 2 & 3 \end{array} \right| & \begin{array}{l} r_1 \leftrightarrow r_2 \\ -r_1 \rightarrow r_1 \\ = \end{array} & \left| \begin{array}{ccc} 1 & -2 & -1 \\ 2 & 1 & 3 \\ -2 & 2 & 3 \end{array} \right| \begin{array}{l} -2r_1 + r_2 \rightarrow r_2 \\ 2r_1 + r_3 \rightarrow r_3 \\ = \end{array} \\ \left| \begin{array}{ccc} 1 & -2 & -1 \\ 0 & 5 & 5 \\ 0 & -2 & 1 \end{array} \right| & \begin{array}{l} \frac{1}{5}r_2 \rightarrow r_2 \\ = \end{array} & \end{array}$$



### Example

By using properties of determinants find the determinant of the

$$\text{matrix } A = \begin{pmatrix} 2 & 1 & 3 \\ -1 & 2 & 1 \\ -2 & 2 & 3 \end{pmatrix}.$$

$$\begin{array}{ccc} \left| \begin{array}{ccc} 2 & 1 & 3 \\ -1 & 2 & 1 \\ -2 & 2 & 3 \end{array} \right| & \begin{array}{l} r_1 \leftrightarrow r_2 \\ -r_1 \rightarrow r_1 \\ = \end{array} & \left| \begin{array}{ccc} 1 & -2 & -1 \\ 2 & 1 & 3 \\ -2 & 2 & 3 \end{array} \right| \begin{array}{l} -2r_1 + r_2 \rightarrow r_2 \\ 2r_1 + r_3 \rightarrow r_3 \\ = \end{array} \\ \left| \begin{array}{ccc} 1 & -2 & -1 \\ 0 & 5 & 5 \\ 0 & -2 & 1 \end{array} \right| & \begin{array}{l} \frac{1}{5}r_2 \rightarrow r_2 \\ = \end{array} & 5 \left| \begin{array}{ccc} 1 & -2 & -1 \\ 0 & 1 & 1 \\ 0 & -2 & 1 \end{array} \right| \begin{array}{l} 2r_2 + r_3 \rightarrow r_3 \\ = \end{array} \end{array}$$

### Example

By using properties of determinants find the determinant of the

$$\text{matrix } A = \begin{pmatrix} 2 & 1 & 3 \\ -1 & 2 & 1 \\ -2 & 2 & 3 \end{pmatrix}.$$

$$\begin{array}{l} \left| \begin{array}{ccc} 2 & 1 & 3 \\ -1 & 2 & 1 \\ -2 & 2 & 3 \end{array} \right| \begin{array}{l} r_1 \leftrightarrow r_2 \\ -r_1 \rightarrow r_1 \\ = \end{array} \left| \begin{array}{ccc} 1 & -2 & -1 \\ 2 & 1 & 3 \\ -2 & 2 & 3 \end{array} \right| \begin{array}{l} -2r_1 + r_2 \rightarrow r_2 \\ 2r_1 + r_3 \rightarrow r_3 \\ = \end{array} \\ \left| \begin{array}{ccc} 1 & -2 & -1 \\ 0 & 5 & 5 \\ 0 & -2 & 1 \end{array} \right| \begin{array}{l} \frac{1}{5}r_2 \rightarrow r_2 \\ = \end{array} 5 \left| \begin{array}{ccc} 1 & -2 & -1 \\ 0 & 1 & 1 \\ 0 & -2 & 1 \end{array} \right| \begin{array}{l} 2r_2 + r_3 \rightarrow r_3 \\ = \end{array} \\ \Rightarrow 5 \left| \begin{array}{ccc} 1 & -2 & -1 \\ 0 & 1 & 1 \\ 0 & 0 & 3 \end{array} \right| \Rightarrow |A| = 5 \cdot (1 \cdot 1 \cdot 3) = 15. \end{array}$$

?