



# MAT1320-Linear Algebra

## Lecture Notes

Echelon Form of a Matrix

---

Mehmet E. KÖROĞLU  
Summer 2020

YILDIZ TECHNICAL UNIVERSITY, DEPARTMENT OF MATHEMATICS  
*mkoroglu@yildiz.edu.tr*

# Table of contents

1. Echelon Form of a Matrix
2. Elementary Row (Column) Operations
3. Finding an Inverse using Elementary Row Operations

# Echelon Form of a Matrix

---

# Echelon Form of a Matrix

An  $m \times n$  matrix  $A$  is said to be in **reduced row echelon form (RREF)** if it satisfies the following properties:

# Echelon Form of a Matrix

An  $m \times n$  matrix  $A$  is said to be in **reduced row echelon form (RREF)** if it satisfies the following properties:

- a) All zero rows, if there are any, appear at the bottom of the matrix.

# Echelon Form of a Matrix

An  $m \times n$  matrix  $A$  is said to be in **reduced row echelon form (RREF)** if it satisfies the following properties:

- a) All zero rows, if there are any, appear at the bottom of the matrix.
- b) The first nonzero entry from the left of a nonzero row is a 1.

# Echelon Form of a Matrix

An  $m \times n$  matrix  $A$  is said to be in **reduced row echelon form (RREF)** if it satisfies the following properties:

- a) All zero rows, if there are any, appear at the bottom of the matrix.
- b) The first nonzero entry from the left of a nonzero row is a 1. This entry is called a **leading one** of its row.

# Echelon Form of a Matrix

An  $m \times n$  matrix  $A$  is said to be in **reduced row echelon form (RREF)** if it satisfies the following properties:

- a) All zero rows, if there are any, appear at the bottom of the matrix.
- b) The first nonzero entry from the left of a nonzero row is a 1. This entry is called a **leading one** of its row.
- c) For each nonzero row, the leading one appears to the right and below any leading ones in preceding rows.



# Echelon Form of a Matrix

An  $m \times n$  matrix  $A$  is said to be in **reduced row echelon form (RREF)** if it satisfies the following properties:

- a) All zero rows, if there are any, appear at the bottom of the matrix.
- b) The first nonzero entry from the left of a nonzero row is a 1. This entry is called a **leading one** of its row.
- c) For each nonzero row, the leading one appears to the right and below any leading ones in preceding rows.
- d) If a column contains a leading one, then all other entries in that column are zero.

# Echelon Form of a Matrix

- A matrix in reduced row echelon form appears as a **staircase** ("echelon") pattern of leading ones descending from the upper left corner of the matrix.

# Echelon Form of a Matrix

- A matrix in reduced row echelon form appears as a **staircase ("echelon")** pattern of leading ones descending from the upper left corner of the matrix.
- An  $m \times n$  matrix satisfying properties a), b), and c) is said to be in **row echelon form (REF)**. There may be no zero rows.

# Echelon Form of a Matrix

- A matrix in reduced row echelon form appears as a staircase ("echelon") pattern of leading ones descending from the upper left corner of the matrix.
- An  $m \times n$  matrix satisfying properties a), b), and c) is said to be in row echelon form (REF). There may be no zero rows.
- A similar definition can be formulated in the obvious manner for reduced column echelon form and column echelon form.

# Echelon Form of a Matrix

## Example

The following are matrices in reduced row echelon form, since they satisfy properties *a)*, *b)*, and *d)*:

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

# Echelon Form of a Matrix

## Example

The following are matrices in reduced row echelon form, since they satisfy properties *a)*, *b)*, and *d)*:

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 0 & 0 & 0 & -2 & 4 \\ 0 & 1 & 0 & 0 & 4 & 8 \\ 0 & 0 & 0 & 1 & 7 & -2 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

# Echelon Form of a Matrix

## Example

The following are matrices in reduced row echelon form, since they satisfy properties *a)*, *b)*, and *d)*:

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 0 & 0 & 0 & -2 & 4 \\ 0 & 1 & 0 & 0 & 4 & 8 \\ 0 & 0 & 0 & 1 & 7 & -2 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

and

$$C = \begin{pmatrix} 1 & 2 & 0 & 0 & 1 \\ 0 & 0 & 1 & 2 & 3 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

# Echelon Form of a Matrix

## Example

The matrices that follow are not in reduced row echelon form.

(Why not?)

$$D = \begin{pmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -3 \end{pmatrix},$$



# Echelon Form of a Matrix

## Example

The matrices that follow are not in reduced row echelon form.

(Why not?)

$$D = \begin{pmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -3 \end{pmatrix}, \quad E = \begin{pmatrix} 1 & 0 & 3 & 4 \\ 0 & 2 & -2 & 5 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

# Echelon Form of a Matrix

## Example

The matrices that follow are not in reduced row echelon form.  
(Why not?)

$$D = \begin{pmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -3 \end{pmatrix}, \quad E = \begin{pmatrix} 1 & 0 & 3 & 4 \\ 0 & 2 & -2 & 5 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

$$F = \begin{pmatrix} 1 & 0 & 3 & 4 \\ 0 & 1 & -2 & 5 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

# Echelon Form of a Matrix

## Example

The matrices that follow are not in reduced row echelon form.  
(Why not?)

$$D = \begin{pmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -3 \end{pmatrix}, \quad E = \begin{pmatrix} 1 & 0 & 3 & 4 \\ 0 & 2 & -2 & 5 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

$$F = \begin{pmatrix} 1 & 0 & 3 & 4 \\ 0 & 1 & -2 & 5 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad G = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 5 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

# Echelon Form of a Matrix

## Example

The following are matrices in row echelon form.

$$H = \begin{pmatrix} 1 & 5 & 0 & 2 & -2 & 4 \\ 0 & 1 & 0 & 3 & 4 & 8 \\ 0 & 0 & 0 & 1 & 7 & -2 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

# Echelon Form of a Matrix

## Example

The following are matrices in row echelon form.

$$H = \begin{pmatrix} 1 & 5 & 0 & 2 & -2 & 4 \\ 0 & 1 & 0 & 3 & 4 & 8 \\ 0 & 0 & 0 & 1 & 7 & -2 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad I = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

# Echelon Form of a Matrix

## Example

The following are matrices in row echelon form.

$$H = \begin{pmatrix} 1 & 5 & 0 & 2 & -2 & 4 \\ 0 & 1 & 0 & 3 & 4 & 8 \\ 0 & 0 & 0 & 1 & 7 & -2 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad I = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

**Note:** We shall now show that every matrix can be put into row (column) echelon form, or into reduced row (column) echelon form, by means of certain row (column) operations.

# Elementary Row (Column) Operations

---

# Elementary Row (Column) Operations

An elementary row (column) operation on a matrix  $A$  is any one of the following operations:

1. **Type I:** Interchange any two rows (columns).



# Elementary Row (Column) Operations

An elementary row (column) operation on a matrix  $A$  is any one of the following operations:

1. **Type I:** Interchange any two rows (columns).
2. **Type II:** Multiply a row (column) by a nonzero number.

# Elementary Row (Column) Operations

An elementary row (column) operation on a matrix  $A$  is any one of the following operations:

1. **Type I:** Interchange any two rows (columns).
2. **Type II:** Multiply a row (column) by a nonzero number.
3. **Type III:** Add a multiple of one row (column) to another.

# Elementary Row (Column) Operations

An elementary row (column) operation on a matrix  $A$  is any one of the following operations:

1. **Type I:** Interchange any two rows (columns).
2. **Type II:** Multiply a row (column) by a nonzero number.
3. **Type III:** Add a multiple of one row (column) to another.

We now introduce the following notation for elementary row and elementary column operations on matrices:

# Elementary Row (Column) Operations

An elementary row (column) operation on a matrix  $A$  is any one of the following operations:

1. **Type I:** Interchange any two rows (columns).
2. **Type II:** Multiply a row (column) by a nonzero number.
3. **Type III:** Add a multiple of one row (column) to another.

We now introduce the following notation for elementary row and elementary column operations on matrices:

- Interchange rows (columns)  $i$  and  $j$ , **Type I:**  $\mathbf{r}_i \leftrightarrow \mathbf{r}_j$  ( $\mathbf{c}_i \leftrightarrow \mathbf{c}_j$ )

# Elementary Row (Column) Operations

An elementary row (column) operation on a matrix  $A$  is any one of the following operations:

1. **Type I:** Interchange any two rows (columns).
2. **Type II:** Multiply a row (column) by a nonzero number.
3. **Type III:** Add a multiple of one row (column) to another.

We now introduce the following notation for elementary row and elementary column operations on matrices:

- Interchange rows (columns)  $i$  and  $j$ , **Type I:**  $\mathbf{r}_i \leftrightarrow \mathbf{r}_j$  ( $\mathbf{c}_i \leftrightarrow \mathbf{c}_j$ )
- Replace row (column)  $i$  by  $k$  times row (column)  $i$ , **Type II:**  
 $k\mathbf{r}_i \rightarrow \mathbf{r}_i$  ( $k\mathbf{c}_i \rightarrow \mathbf{c}_i$ )

# Elementary Row (Column) Operations

An elementary row (column) operation on a matrix  $A$  is any one of the following operations:

1. **Type I:** Interchange any two rows (columns).
2. **Type II:** Multiply a row (column) by a nonzero number.
3. **Type III:** Add a multiple of one row (column) to another.

We now introduce the following notation for elementary row and elementary column operations on matrices:

- Interchange rows (columns)  $i$  and  $j$ , **Type I:**  $\mathbf{r}_i \leftrightarrow \mathbf{r}_j$  ( $\mathbf{c}_i \leftrightarrow \mathbf{c}_j$ )
- Replace row (column)  $i$  by  $k$  times row (column)  $i$ , **Type II:**  
 $k\mathbf{r}_i \rightarrow \mathbf{r}_i$  ( $k\mathbf{c}_i \rightarrow \mathbf{c}_i$ )
- Replace row (column)  $j$  by  $k$  times row (column)  $i$  + row (column)  $j$ , **Type III:**  $k\mathbf{r}_i + \mathbf{r}_j \rightarrow \mathbf{r}_j$  ( $k\mathbf{c}_i + \mathbf{c}_j \rightarrow \mathbf{c}_j$ )

# Echelon Form of a Matrix

## Example

$$\text{Let } A = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ 3 & 3 & 6 & -9 \end{pmatrix}.$$

# Echelon Form of a Matrix

## Example

Let  $A = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ 3 & 3 & 6 & -9 \end{pmatrix}$ .

Interchanging rows 1 and 3 of  $A$ , we obtain

$$B = A_{r_1 \leftrightarrow r_3} = \begin{pmatrix} 3 & 3 & 6 & -9 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$



# Echelon Form of a Matrix

## Example

$$\text{Let } A = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ 3 & 3 & 6 & -9 \end{pmatrix}.$$

Interchanging rows 1 and 3 of  $A$ , we obtain

$$B = A_{r_1 \leftrightarrow r_3} = \begin{pmatrix} 3 & 3 & 6 & -9 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Multiplying the third row of  $A$  by  $\frac{1}{3}$ , we obtain

$$C = A_{\frac{1}{3}r_3 \rightarrow r_3} = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ 1 & 1 & 2 & -3 \end{pmatrix}$$

## Echelon Form of a Matrix

Adding  $(-2)$  times row 2 of  $A$  to row 3 of  $A$ , we obtain

$$D = A_{-2r_2+r_3 \rightarrow r_3} = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ -1 & -3 & 6 & -5 \end{pmatrix}$$

## Echelon Form of a Matrix

Adding  $(-2)$  times row 2 of  $A$  to row 3 of  $A$ , we obtain

$$D = A_{-2r_2+r_3 \rightarrow r_3} = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ -1 & -3 & 6 & -5 \end{pmatrix}$$

Observe that in obtaining  $D$  from  $A$ , row 2 of  $A$  did not change.

## Echelon Form of a Matrix

Adding  $(-2)$  times row 2 of  $A$  to row 3 of  $A$ , we obtain

$$D = A_{-2r_2+r_3 \rightarrow r_3} = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ -1 & -3 & 6 & -5 \end{pmatrix}$$

Observe that in obtaining  $D$  from  $A$ , row 2 of  $A$  did not change.

**Note:** An  $m \times n$  matrix  $B$  is said to be row (column) equivalent to an  $m \times n$  matrix  $A$  if  $B$  can be produced by applying a finite sequence of elementary row (column) operations to  $A$ .

### Example

$$\text{Let } A = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 2 & 1 & 3 & 2 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

### Example

$$\text{Let } A = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 2 & 1 & 3 & 2 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

If we add 2 times row 3 of  $A$  to its second row, we obtain

$$B = A_{2r_3+r_2 \rightarrow r_2} = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 4 & -3 & 7 & 8 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

### Example

$$\text{Let } A = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 2 & 1 & 3 & 2 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

If we add 2 times row 3 of  $A$  to its second row, we obtain

$$B = A_{2r_3+r_2 \rightarrow r_2} = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 4 & -3 & 7 & 8 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

so  $B$  is row equivalent to  $A$ .

### Example

$$\text{Let } A = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 2 & 1 & 3 & 2 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

If we add 2 times row 3 of  $A$  to its second row, we obtain

$$B = A_{2r_3+r_2 \rightarrow r_2} = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 4 & -3 & 7 & 8 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

so  $B$  is row equivalent to  $A$ . Multiplying row 1 of  $B$  by 2, we obtain

$$C = B_{2r_1 \rightarrow r_1} = \begin{pmatrix} 2 & 4 & 8 & 6 \\ 4 & -3 & 7 & 8 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$



### Example

$$\text{Let } A = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 2 & 1 & 3 & 2 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

If we add 2 times row 3 of  $A$  to its second row, we obtain

$$B = A_{2r_3+r_2 \rightarrow r_2} = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 4 & -3 & 7 & 8 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

so  $B$  is row equivalent to  $A$ . Multiplying row 1 of  $B$  by 2, we obtain

$$C = B_{2r_1 \rightarrow r_1} = \begin{pmatrix} 2 & 4 & 8 & 6 \\ 4 & -3 & 7 & 8 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

so  $C$  is row equivalent to  $B$ .

### Example

$$\text{Let } A = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 2 & 1 & 3 & 2 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

If we add 2 times row 3 of  $A$  to its second row, we obtain

$$B = A_{2r_3+r_2 \rightarrow r_2} = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 4 & -3 & 7 & 8 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

so  $B$  is row equivalent to  $A$ . Multiplying row 1 of  $B$  by 2, we obtain

$$C = B_{2r_1 \rightarrow r_1} = \begin{pmatrix} 2 & 4 & 8 & 6 \\ 4 & -3 & 7 & 8 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

so  $C$  is row equivalent to  $B$ . It then follows that  $C$  is row equivalent to  $A$ , since we obtained  $C$  by applying two successive elementary row operations to  $A$ .

# Echelon Form of a Matrix

## Theorem

*Every nonzero  $m \times n$  matrix  $\mathbf{A} = [a_{ij}]$  is row (column) equivalent to a matrix in row (column) echelon form.*

# Echelon Form of a Matrix

## Theorem

*Every nonzero  $m \times n$  matrix  $\mathbf{A} = [a_{ij}]$  is row (column) equivalent to a matrix in row (column) echelon form.*

## Theorem

*The RREF of a matrix is unique.*

# Echelon Form of a Matrix

## Theorem

*Every nonzero  $m \times n$  matrix  $\mathbf{A} = [a_{ij}]$  is row (column) equivalent to a matrix in row (column) echelon form.*

## Theorem

*The RREF of a matrix is unique.*

**Note:** In any matrix, the first column with a nonzero entry is called the **pivot column**; the first nonzero entry in the pivot column is called the **pivot**.

# Echelon Form of a Matrix

## Theorem

Every nonzero  $m \times n$  matrix  $\mathbf{A} = [a_{ij}]$  is row (column) equivalent to a matrix in row (column) echelon form.

## Theorem

The RREF of a matrix is unique.

**Note:** In any matrix, the first column with a nonzero entry is called the **pivot column**; the first nonzero entry in the pivot column is called the **pivot**.

## Example

Let  $A = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ 3 & 3 & 6 & -9 \end{pmatrix}$ . Find REF and RREF of  $A$ .

# Echelon Form of a Matrix

## Theorem

Every nonzero  $m \times n$  matrix  $\mathbf{A} = [a_{ij}]$  is row (column) equivalent to a matrix in row (column) echelon form.

## Theorem

The RREF of a matrix is unique.

**Note:** In any matrix, the first column with a nonzero entry is called the **pivot column**; the first nonzero entry in the pivot column is called the **pivot**.

## Example

Let  $A = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ 3 & 3 & 6 & -9 \end{pmatrix}$ . Find REF and RREF of  $A$ .

### Example (cont.)

Interchanging rows 1 and 3 of  $A$ , we obtain

$$B = A_{r_1 \longleftrightarrow r_3} = \begin{pmatrix} 3 & 3 & 6 & -9 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$



### Example (cont.)

Interchanging rows 1 and 3 of  $A$ , we obtain

$$B = A_{r_1 \leftrightarrow r_3} = \begin{pmatrix} 3 & 3 & 6 & -9 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Multiplying the first row of  $B$  by  $\frac{1}{3}$ , we obtain

$$C = B_{\frac{1}{3}r_1 \rightarrow r_1} = \begin{pmatrix} 1 & 1 & 2 & -3 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

### Example (cont.)

Interchanging rows 1 and 3 of  $A$ , we obtain

$$B = A_{r_1 \leftrightarrow r_3} = \begin{pmatrix} 3 & 3 & 6 & -9 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Multiplying the first row of  $B$  by  $\frac{1}{3}$ , we obtain

$$C = B_{\frac{1}{3}r_1 \rightarrow r_1} = \begin{pmatrix} 1 & 1 & 2 & -3 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Adding  $(-2)$  times row 1 of  $C$  to row 2 of  $C$ , we obtain

$$D = C_{-2r_1 + r_2 \rightarrow r_2} = \underbrace{\begin{pmatrix} 1 & 1 & 2 & -3 \\ 0 & 1 & -4 & 4 \\ 0 & 0 & 1 & 2 \end{pmatrix}}_{\text{REF}}$$

### Example (cont.)

Interchanging rows 1 and 3 of  $A$ , we obtain

$$B = A_{r_1 \leftrightarrow r_3} = \begin{pmatrix} 3 & 3 & 6 & -9 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Multiplying the first row of  $B$  by  $\frac{1}{3}$ , we obtain

$$C = B_{\frac{1}{3}r_1 \rightarrow r_1} = \begin{pmatrix} 1 & 1 & 2 & -3 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Adding  $(-2)$  times row 1 of  $C$  to row 2 of  $C$ , we obtain

$$D = C_{-2r_1 + r_2 \rightarrow r_2} = \underbrace{\begin{pmatrix} 1 & 1 & 2 & -3 \\ 0 & 1 & -4 & 4 \\ 0 & 0 & 1 & 2 \end{pmatrix}}_{\text{REF}}$$

### Example (cont.)

Adding  $(-2)$  times row 3 of  $D$  to row 1 of  $D$  and 4 times row 3 of  $D$  to row 2 of  $D$ , we obtain

$$E = D_{\substack{4r_3+r_2 \rightarrow r_2 \\ -2r_3+r_1 \rightarrow r_1}} = \begin{pmatrix} 1 & 1 & 0 & -7 \\ 0 & 1 & 0 & 12 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

### Example (cont.)

Adding  $(-2)$  times row 3 of  $D$  to row 1 of  $D$  and 4 times row 3 of  $D$  to row 2 of  $D$ , we obtain

$$E = D_{\substack{4r_3+r_2 \rightarrow r_2 \\ -2r_3+r_1 \rightarrow r_1}} = \begin{pmatrix} 1 & 1 & 0 & -7 \\ 0 & 1 & 0 & 12 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Adding  $(-1)$  times row 2 of  $E$  to row 1 of  $E$ , we obtain

$$F = E_{-r_2+r_1 \rightarrow r_1} = \underbrace{\begin{pmatrix} 1 & 0 & 0 & -19 \\ 0 & 1 & 0 & 12 \\ 0 & 0 & 1 & 2 \end{pmatrix}}_{\text{RREF}}$$

### Example (cont.)

Adding  $(-2)$  times row 3 of  $D$  to row 1 of  $D$  and 4 times row 3 of  $D$  to row 2 of  $D$ , we obtain

$$E = D_{\substack{4r_3+r_2 \rightarrow r_2 \\ -2r_3+r_1 \rightarrow r_1}} = \begin{pmatrix} 1 & 1 & 0 & -7 \\ 0 & 1 & 0 & 12 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Adding  $(-1)$  times row 2 of  $E$  to row 1 of  $E$ , we obtain

$$F = E_{-r_2+r_1 \rightarrow r_1} = \underbrace{\begin{pmatrix} 1 & 0 & 0 & -19 \\ 0 & 1 & 0 & 12 \\ 0 & 0 & 1 & 2 \end{pmatrix}}_{\text{RREF}}$$

**Note:** The number of nonzero rows in the REF or RREF of a matrix  $\mathbf{A}$  is called the **rank** of  $\mathbf{A}$  and denoted by  $\text{rank}(\mathbf{A})$ .

### Example (cont.)

Adding  $(-2)$  times row 3 of  $D$  to row 1 of  $D$  and 4 times row 3 of  $D$  to row 2 of  $D$ , we obtain

$$E = D_{\substack{4r_3+r_2 \rightarrow r_2 \\ -2r_3+r_1 \rightarrow r_1}} = \begin{pmatrix} 1 & 1 & 0 & -7 \\ 0 & 1 & 0 & 12 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Adding  $(-1)$  times row 2 of  $E$  to row 1 of  $E$ , we obtain

$$F = E_{-r_2+r_1 \rightarrow r_1} = \underbrace{\begin{pmatrix} 1 & 0 & 0 & -19 \\ 0 & 1 & 0 & 12 \\ 0 & 0 & 1 & 2 \end{pmatrix}}_{\text{RREF}}$$

**Note:** The number of nonzero rows in the REF or RREF of a matrix  $\mathbf{A}$  is called the **rank** of  $\mathbf{A}$  and denoted by **rank** ( $\mathbf{A}$ ). The

rank of matrix  $\mathbf{A}$  given in the previous example is 3.

## **Finding an Inverse using Elementary Row Operations**

---



## Finding an Inverse using Elementary Row Operations

- If the inverse of a matrix  $\mathbf{A}$  exists, then there is a series of elementary row operations that reduces  $\mathbf{A}$  to the identity matrix.

## Finding an Inverse using Elementary Row Operations

- If the inverse of a matrix  $\mathbf{A}$  exists, then there is a series of elementary row operations that reduces  $\mathbf{A}$  to the identity matrix.
- Suppose the required elementary row operations are (in order)  $\mathbf{E}_1, \mathbf{E}_2, \dots, \mathbf{E}_n$ , then

$$\mathbf{E}_n \dots \mathbf{E}_2 \mathbf{E}_1 \mathbf{A} = \mathbf{I}$$

which means that  $\mathbf{E}_n \dots \mathbf{E}_2 \mathbf{E}_1 = \mathbf{A}^{-1}$ .

# Finding an Inverse using Elementary Row Operations

- If the inverse of a matrix  $\mathbf{A}$  exists, then there is a series of elementary row operations that reduces  $\mathbf{A}$  to the identity matrix.
- Suppose the required elementary row operations are (in order)  $\mathbf{E}_1, \mathbf{E}_2, \dots, \mathbf{E}_n$ , then

$$\mathbf{E}_n \dots \mathbf{E}_2 \mathbf{E}_1 \mathbf{A} = \mathbf{I}$$

which means that  $\mathbf{E}_n \dots \mathbf{E}_2 \mathbf{E}_1 = \mathbf{A}^{-1}$ .

- Furthermore, because

$$\mathbf{E}_n \dots \mathbf{E}_2 \mathbf{E}_1 \mathbf{I} = \mathbf{E}_n \dots \mathbf{E}_2 \mathbf{E}_1$$

we can use the following technique:

## Finding an Inverse using Elementary Row Operations

- Write  $\mathbf{A}$  and  $\mathbf{I}$  side-by-side.

## Finding an Inverse using Elementary Row Operations

- Write  $\mathbf{A}$  and  $\mathbf{I}$  side-by-side.
- Then apply the same row operations to both  $\mathbf{A}$  and  $\mathbf{I}$  until  $\mathbf{A}$  is reduced to the identity matrix.

# Finding an Inverse using Elementary Row Operations

- Write **A** and **I** side-by-side.
- Then apply the same row operations to both **A** and **I** until **A** is reduced to the identity matrix.
- At the same time, the identity matrix will be "reduced" to the inverse matrix.

$$\begin{array}{c|c} \mathbf{A} & \mathbf{I} \\ \hline \mathbf{E}_1 \mathbf{A} & \mathbf{E}_1 \mathbf{I} \\ \mathbf{E}_2 \mathbf{E}_1 \mathbf{A} & \mathbf{E}_2 \mathbf{E}_1 \mathbf{I} \\ \mathbf{E}_n \dots \mathbf{E}_2 \mathbf{E}_1 \mathbf{A} & \mathbf{E}_n \dots \mathbf{E}_2 \mathbf{E}_1 \mathbf{I} \\ \hline \underbrace{\hspace{10em}}_{\text{reduced to } \mathbf{I}} & \underbrace{\hspace{10em}}_{\mathbf{A}^{-1}} \end{array}$$

# Finding an Inverse using Elementary Row Operations

- Write **A** and **I** side-by-side.
- Then apply the same row operations to both **A** and **I** until **A** is reduced to the identity matrix.
- At the same time, the identity matrix will be "reduced" to the inverse matrix.

$$\begin{array}{c|c} \mathbf{A} & \mathbf{I} \\ \mathbf{E}_1\mathbf{A} & \mathbf{E}_1\mathbf{I} \\ \mathbf{E}_2\mathbf{E}_1\mathbf{A} & \mathbf{E}_2\mathbf{E}_1\mathbf{I} \\ \underbrace{\mathbf{E}_n \dots \mathbf{E}_2\mathbf{E}_1\mathbf{A}}_{\text{reduced to I}} & \underbrace{\mathbf{E}_n \dots \mathbf{E}_2\mathbf{E}_1\mathbf{I}}_{\mathbf{A}^{-1}} \end{array}$$

Here is the fully worked out example:

### Example

Let  $A = \begin{pmatrix} 3 & 3 & 6 \\ 2 & 3 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ . By using elementary row operations, find the inverse of the  $A$ .



### Example

Let  $A = \begin{pmatrix} 3 & 3 & 6 \\ 2 & 3 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ . By using elementary row operations, find the inverse of the  $A$ .

$$B = [A|I_3] = \left( \begin{array}{ccc|ccc} 3 & 3 & 6 & 1 & 0 & 0 \\ 2 & 3 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

### Example

Let  $A = \begin{pmatrix} 3 & 3 & 6 \\ 2 & 3 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ . By using elementary row operations, find the inverse of the  $A$ .

$$B = [A|I_3] = \left( \begin{array}{ccc|ccc} 3 & 3 & 6 & 1 & 0 & 0 \\ 2 & 3 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

Multiplying the first row of  $B$  by  $\frac{1}{3}$ , we obtain

$$C = B_{\frac{1}{3}r_1 \rightarrow r_1} = \left( \begin{array}{ccc|ccc} 1 & 1 & 2 & \frac{1}{3} & 0 & 0 \\ 2 & 3 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

### Example (cont.)

Adding  $(-2)$  times row 1 of  $C$  to row 2 of  $C$ , we obtain

$$D = C_{-2r_1+r_2 \rightarrow r_2} \left( \begin{array}{ccc|ccc} 1 & 1 & 2 & \frac{1}{3} & 0 & 0 \\ 0 & 1 & -4 & \frac{-2}{3} & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

### Example (cont.)

Adding  $(-2)$  times row 1 of  $C$  to row 2 of  $C$ , we obtain

$$D = C_{-2r_1+r_2 \rightarrow r_2} \left( \begin{array}{ccc|ccc} 1 & 1 & 2 & \frac{1}{3} & 0 & 0 \\ 0 & 1 & -4 & \frac{-2}{3} & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

Adding  $(-2)$  times row 3 of  $D$  to row 1 of  $D$  and 4 times row 3 of  $D$  to row 2 of  $D$ , we obtain

$$E = D_{\substack{4r_3+r_2 \rightarrow r_2 \\ -2r_3+r_1 \rightarrow r_1}} = \left( \begin{array}{ccc|ccc} 1 & 1 & 0 & \frac{1}{3} & 0 & -2 \\ 0 & 1 & 0 & \frac{-2}{3} & 1 & 4 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

### Example (cont.)

Adding  $(-2)$  times row 1 of  $C$  to row 2 of  $C$ , we obtain

$$D = C_{-2r_1+r_2 \rightarrow r_2} \left( \begin{array}{ccc|ccc} 1 & 1 & 2 & \frac{1}{3} & 0 & 0 \\ 0 & 1 & -4 & \frac{-2}{3} & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

Adding  $(-2)$  times row 3 of  $D$  to row 1 of  $D$  and 4 times row 3 of  $D$  to row 2 of  $D$ , we obtain

$$E = D_{\substack{4r_3+r_2 \rightarrow r_2 \\ -2r_3+r_1 \rightarrow r_1}} = \left( \begin{array}{ccc|ccc} 1 & 1 & 0 & \frac{1}{3} & 0 & -2 \\ 0 & 1 & 0 & \frac{-2}{3} & 1 & 4 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

Adding  $(-1)$  times row 2 of  $E$  to row 1 of  $E$ , we obtain

$$E = \left( \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & -1 & -6 \\ 0 & 1 & 0 & \frac{-2}{3} & 1 & 4 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

### Example (cont.)

Adding  $(-2)$  times row 1 of  $C$  to row 2 of  $C$ , we obtain

$$D = C_{-2r_1+r_2 \rightarrow r_2} \left( \begin{array}{ccc|ccc} 1 & 1 & 2 & \frac{1}{3} & 0 & 0 \\ 0 & 1 & -4 & \frac{-2}{3} & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

Adding  $(-2)$  times row 3 of  $D$  to row 1 of  $D$  and 4 times row 3 of  $D$  to row 2 of  $D$ , we obtain

$$E = D_{\substack{4r_3+r_2 \rightarrow r_2 \\ -2r_3+r_1 \rightarrow r_1}} = \left( \begin{array}{ccc|ccc} 1 & 1 & 0 & \frac{1}{3} & 0 & -2 \\ 0 & 1 & 0 & \frac{-2}{3} & 1 & 4 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

Adding  $(-1)$  times row 2 of  $E$  to row 1 of  $E$ , we obtain

$$E = \left( \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & -1 & -6 \\ 0 & 1 & 0 & \frac{-2}{3} & 1 & 4 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right), \quad A^{-1} = \left( \begin{array}{ccc} 1 & -1 & -6 \\ \frac{-2}{3} & 1 & 4 \\ 0 & 0 & 1 \end{array} \right).$$

?