



# MAT1320-Linear Algebra Lecture Notes

## Vector Spaces, Subspaces

---

Mehmet E. KÖROĞLU

Fall 2020

YILDIZ TECHNICAL UNIVERSITY, DEPARTMENT OF MATHEMATICS

*mkoroglu@yildiz.edu.tr*

# Table of contents

1. Vector Spaces

2. Subspaces

# Vector Spaces

---

**Definition**

Let  $\emptyset \neq V$  be nonempty set and  $\mathbb{R}$  (in general a field  $F$ ) be the field of real numbers.

# Vector Spaces

## Definition

Let  $\emptyset \neq V$  be nonempty set and  $\mathbb{R}$  (in general a field  $F$ ) be the field of real numbers. Let

$$+ : V \times V \longrightarrow V,$$

# Vector Spaces

## Definition

Let  $\emptyset \neq V$  be nonempty set and  $\mathbb{R}$  (in general a field  $F$ ) be the field of real numbers. Let

$$\begin{aligned} + & : V \times V \longrightarrow V, \\ (\vec{u}, \vec{v}) & \longmapsto \vec{u} + \vec{v} \quad (\text{Addition}) \end{aligned}$$

# Vector Spaces

## Definition

Let  $\emptyset \neq V$  be nonempty set and  $\mathbb{R}$  (in general a field  $F$ ) be the field of real numbers. Let

$$\begin{aligned} + & : V \times V \longrightarrow V, \\ (\vec{u}, \vec{v}) & \longmapsto \vec{u} + \vec{v} \text{ (Addition)} \\ \cdot & : \mathbb{R} \times V \longrightarrow V, \end{aligned}$$

# Vector Spaces

## Definition

Let  $\emptyset \neq V$  be nonempty set and  $\mathbb{R}$  (in general a field  $F$ ) be the field of real numbers. Let

$$\begin{aligned} + & : V \times V \longrightarrow V, \\ (\vec{u}, \vec{v}) & \longmapsto \vec{u} + \vec{v} \text{ (Addition)} \\ \cdot & : \mathbb{R} \times V \longrightarrow V, \\ (\lambda, \vec{u}) & \longmapsto \lambda \vec{u} \text{ (Scalar Multiplication)} \end{aligned}$$



# Vector Spaces

## Definition

Let  $\emptyset \neq V$  be nonempty set and  $\mathbb{R}$  (in general a field  $F$ ) be the field of real numbers. Let

$$\begin{aligned} + & : V \times V \longrightarrow V, \\ (\vec{u}, \vec{v}) & \longmapsto \vec{u} + \vec{v} \text{ (Addition)} \\ \cdot & : \mathbb{R} \times V \longrightarrow V, \\ (\lambda, \vec{u}) & \longmapsto \lambda \vec{u} \text{ (Scalar Multiplication)} \end{aligned}$$

be two binary operations on  $V$ . The triple- $(V, +, \cdot)$  is called a **vector space** over  $\mathbb{R}$  if the following axioms satisfied.

# Vector Spaces

## Definition

Let  $\emptyset \neq V$  be nonempty set and  $\mathbb{R}$  (in general a field  $F$ ) be the field of real numbers. Let

$$\begin{aligned} + & : V \times V \longrightarrow V, \\ (\vec{u}, \vec{v}) & \longmapsto \vec{u} + \vec{v} \text{ (Addition)} \\ \cdot & : \mathbb{R} \times V \longrightarrow V, \\ (\lambda, \vec{u}) & \longmapsto \lambda \vec{u} \text{ (Scalar Multiplication)} \end{aligned}$$

be two binary operations on  $V$ . The triple- $(V, +, \cdot)$  is called a **vector space** over  $\mathbb{R}$  if the following axioms satisfied. The element of  $V$  are called **vectors**.

# Vector Spaces

1. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} \in V$

# Vector Spaces

1. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} \in V$
2. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} = \vec{w} + \vec{v}$

# Vector Spaces

1. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} \in V$
2. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} = \vec{w} + \vec{v}$
3. For each  $\vec{u}, \vec{v}, \vec{w} \in V : (\vec{v} + \vec{w}) + \vec{u} = \vec{v} + (\vec{w} + \vec{u})$

# Vector Spaces

1. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} \in V$
2. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} = \vec{w} + \vec{v}$
3. For each  $\vec{u}, \vec{v}, \vec{w} \in V : (\vec{v} + \vec{w}) + \vec{u} = \vec{v} + (\vec{w} + \vec{u})$
4. For each  $\vec{v} \in V : \vec{v} + \vec{0} = \vec{v}$  such that  $\vec{0} \in V$  exists.

# Vector Spaces

1. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} \in V$
2. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} = \vec{w} + \vec{v}$
3. For each  $\vec{u}, \vec{v}, \vec{w} \in V : (\vec{v} + \vec{w}) + \vec{u} = \vec{v} + (\vec{w} + \vec{u})$
4. For each  $\vec{v} \in V : \vec{v} + \vec{0} = \vec{v}$  such that  $\vec{0} \in V$  exists.
5. For each  $\vec{v} \in V, \vec{v} + \vec{w} = \vec{0}$  such that  $\vec{w} = -\vec{v} \in V$  exists.

# Vector Spaces

1. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} \in V$
2. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} = \vec{w} + \vec{v}$
3. For each  $\vec{u}, \vec{v}, \vec{w} \in V : (\vec{v} + \vec{w}) + \vec{u} = \vec{v} + (\vec{w} + \vec{u})$
4. For each  $\vec{v} \in V : \vec{v} + \vec{0} = \vec{v}$  such that  $\vec{0} \in V$  exists.
5. For each  $\vec{v} \in V, \vec{v} + \vec{w} = \vec{0}$  such that  $\vec{w} = -\vec{v} \in V$  exists.
6. For each  $r \in \mathbb{R}$  and  $\vec{v} \in V, r\vec{v} \in V$



# Vector Spaces

1. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} \in V$
2. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} = \vec{w} + \vec{v}$
3. For each  $\vec{u}, \vec{v}, \vec{w} \in V : (\vec{v} + \vec{w}) + \vec{u} = \vec{v} + (\vec{w} + \vec{u})$
4. For each  $\vec{v} \in V : \vec{v} + \vec{0} = \vec{v}$  such that  $\vec{0} \in V$  exists.
5. For each  $\vec{v} \in V, \vec{v} + \vec{w} = \vec{0}$  such that  $\vec{w} = -\vec{v} \in V$  exists.
6. For each  $r \in \mathbb{R}$  and  $\vec{v} \in V, r\vec{v} \in V$
7. For each  $r, s \in \mathbb{R}$  and  $\vec{v} \in V, (r + s)\vec{v} = r\vec{v} + s\vec{v}$

# Vector Spaces

1. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} \in V$
2. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} = \vec{w} + \vec{v}$
3. For each  $\vec{u}, \vec{v}, \vec{w} \in V : (\vec{v} + \vec{w}) + \vec{u} = \vec{v} + (\vec{w} + \vec{u})$
4. For each  $\vec{v} \in V : \vec{v} + \vec{0} = \vec{v}$  such that  $\vec{0} \in V$  exists.
5. For each  $\vec{v} \in V, \vec{v} + \vec{w} = \vec{0}$  such that  $\vec{w} = -\vec{v} \in V$  exists.
6. For each  $r \in \mathbb{R}$  and  $\vec{v} \in V, r\vec{v} \in V$
7. For each  $r, s \in \mathbb{R}$  and  $\vec{v} \in V, (r + s)\vec{v} = r\vec{v} + s\vec{v}$
8. For each  $r \in \mathbb{R}$  and  $\vec{v}, \vec{w} \in V, r(\vec{v} + \vec{w}) = r\vec{v} + r\vec{w}$

# Vector Spaces

1. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} \in V$
2. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} = \vec{w} + \vec{v}$
3. For each  $\vec{u}, \vec{v}, \vec{w} \in V : (\vec{v} + \vec{w}) + \vec{u} = \vec{v} + (\vec{w} + \vec{u})$
4. For each  $\vec{v} \in V : \vec{v} + \vec{0} = \vec{v}$  such that  $\vec{0} \in V$  exists.
5. For each  $\vec{v} \in V, \vec{v} + \vec{w} = \vec{0}$  such that  $\vec{w} = -\vec{v} \in V$  exists.
6. For each  $r \in \mathbb{R}$  and  $\vec{v} \in V, r\vec{v} \in V$
7. For each  $r, s \in \mathbb{R}$  and  $\vec{v} \in V, (r + s)\vec{v} = r\vec{v} + s\vec{v}$
8. For each  $r \in \mathbb{R}$  and  $\vec{v}, \vec{w} \in V, r(\vec{v} + \vec{w}) = r\vec{v} + r\vec{w}$
9. For each  $r, s \in \mathbb{R}$  and  $\vec{v} \in V, (rs)\vec{v} = r(s\vec{v})$

# Vector Spaces

1. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} \in V$
2. For each  $\vec{v}, \vec{w} \in V : \vec{v} + \vec{w} = \vec{w} + \vec{v}$
3. For each  $\vec{u}, \vec{v}, \vec{w} \in V : (\vec{v} + \vec{w}) + \vec{u} = \vec{v} + (\vec{w} + \vec{u})$
4. For each  $\vec{v} \in V : \vec{v} + \vec{0} = \vec{v}$  such that  $\vec{0} \in V$  exists.
5. For each  $\vec{v} \in V, \vec{v} + \vec{w} = \vec{0}$  such that  $\vec{w} = -\vec{v} \in V$  exists.
6. For each  $r \in \mathbb{R}$  and  $\vec{v} \in V, r\vec{v} \in V$
7. For each  $r, s \in \mathbb{R}$  and  $\vec{v} \in V, (r + s)\vec{v} = r\vec{v} + s\vec{v}$
8. For each  $r \in \mathbb{R}$  and  $\vec{v}, \vec{w} \in V, r(\vec{v} + \vec{w}) = r\vec{v} + r\vec{w}$
9. For each  $r, s \in \mathbb{R}$  and  $\vec{v} \in V, (rs)\vec{v} = r(s\vec{v})$
10. For each  $\vec{v} \in V, 1\vec{v} = \vec{v}$

## Example

The set of  $n$ -tuples  $V = \mathbb{R}^n = \{ \vec{x} = (x_1, x_2, \dots, x_n) \mid x_i \in \mathbb{R} \}$  together with the following binary operations

$$+ : V \times V \longrightarrow V,$$

## Example

The set of  $n$ -tuples  $V = \mathbb{R}^n = \{ \vec{x} = (x_1, x_2, \dots, x_n) \mid x_i \in \mathbb{R} \}$  together with the following binary operations

$$\begin{aligned} + & : V \times V \longrightarrow V, \\ (\vec{x}, \vec{y}) & \longmapsto \vec{x} + \vec{y} = (x_1 + y_1, x_2 + y_2, \dots, x_n + y_n) \end{aligned}$$

## Example

The set of  $n$ -tuples  $V = \mathbb{R}^n = \{ \vec{x} = (x_1, x_2, \dots, x_n) \mid x_i \in \mathbb{R} \}$  together with the following binary operations

$$\begin{aligned} + & : V \times V \longrightarrow V, \\ (\vec{x}, \vec{y}) & \longmapsto \vec{x} + \vec{y} = (x_1 + y_1, x_2 + y_2, \dots, x_n + y_n) \\ \cdot & : \mathbb{R} \times V \longrightarrow V, \end{aligned}$$

## Example

The set of  $n$ -tuples  $V = \mathbb{R}^n = \{ \vec{x} = (x_1, x_2, \dots, x_n) \mid x_i \in \mathbb{R} \}$  together with the following binary operations

$$\begin{aligned} + & : V \times V \longrightarrow V, \\ (\vec{x}, \vec{y}) & \longmapsto \vec{x} + \vec{y} = (x_1 + y_1, x_2 + y_2, \dots, x_n + y_n) \\ \cdot & : \mathbb{R} \times V \longrightarrow V, \\ (\lambda, \vec{x}) & \longmapsto \lambda \vec{x} = (\lambda x_1, \lambda x_2, \dots, \lambda x_n) \end{aligned}$$



## Example

The set of  $n$ -tuples  $V = \mathbb{R}^n = \{ \vec{x} = (x_1, x_2, \dots, x_n) \mid x_i \in \mathbb{R} \}$  together with the following binary operations

$$\begin{aligned} + & : V \times V \longrightarrow V, \\ (\vec{x}, \vec{y}) & \longmapsto \vec{x} + \vec{y} = (x_1 + y_1, x_2 + y_2, \dots, x_n + y_n) \\ \cdot & : \mathbb{R} \times V \longrightarrow V, \\ (\lambda, \vec{x}) & \longmapsto \lambda \vec{x} = (\lambda x_1, \lambda x_2, \dots, \lambda x_n) \end{aligned}$$

is a vector space over  $\mathbb{R}$ .

# Vector Spaces

## Example

Let  $V = \mathbb{R}^2 = \{ \vec{x} = (a, b) \mid a, b \in \mathbb{R} \}$  and

$$+ : V \times V \longrightarrow V, \quad (a, b) + (c, d) = (a + c, b + d)$$

$$\cdot : \mathbb{R} \times V \longrightarrow V, \quad \lambda (a, b) = (a, \lambda b)$$

be given.

# Vector Spaces

## Example

Let  $V = \mathbb{R}^2 = \{ \vec{x} = (a, b) \mid a, b \in \mathbb{R} \}$  and

$$+ : V \times V \longrightarrow V, \quad (a, b) + (c, d) = (a + c, b + d)$$

$$\cdot : \mathbb{R} \times V \longrightarrow V, \quad \lambda(a, b) = (a, \lambda b)$$

be given. Then  $(V, +, \cdot)$  is not a vector space. Because, the following axiom is not satisfied.

# Vector Spaces

## Example

Let  $V = \mathbb{R}^2 = \{ \vec{x} = (a, b) \mid a, b \in \mathbb{R} \}$  and

$$+ : V \times V \longrightarrow V, \quad (a, b) + (c, d) = (a + c, b + d)$$

$$\cdot : \mathbb{R} \times V \longrightarrow V, \quad \lambda (a, b) = (a, \lambda b)$$

be given. Then  $(V, +, \cdot)$  is not a vector space. Because, the following axiom is not satisfied. For each  $r, s \in \mathbb{R}$  and  $\vec{u} = (a, b) \in V$  we have

$$(r + s)\vec{u} = (r + s)(a, b) = (a, (r + s)b)$$

# Vector Spaces

## Example

Let  $V = \mathbb{R}^2 = \{ \vec{x} = (a, b) \mid a, b \in \mathbb{R} \}$  and

$$+ : V \times V \longrightarrow V, (a, b) + (c, d) = (a + c, b + d)$$

$$\cdot : \mathbb{R} \times V \longrightarrow V, \lambda (a, b) = (a, \lambda b)$$

be given. Then  $(V, +, \cdot)$  is not a vector space. Because, the following axiom is not satisfied. For each  $r, s \in \mathbb{R}$  and  $\vec{u} = (a, b) \in V$  we have

$$(r + s)\vec{u} = (r + s)(a, b) = (a, (r + s)b) \quad (1)$$

$$r\vec{u} + s\vec{u} = (a, rb) + (a, sb) = (2a, (r + s)b)$$

# Vector Spaces

## Example

Let  $V = \mathbb{R}^2 = \{ \vec{x} = (a, b) \mid a, b \in \mathbb{R} \}$  and

$$+ : V \times V \longrightarrow V, \quad (a, b) + (c, d) = (a + c, b + d)$$

$$\cdot : \mathbb{R} \times V \longrightarrow V, \quad \lambda (a, b) = (a, \lambda b)$$

be given. Then  $(V, +, \cdot)$  is not a vector space. Because, the following axiom is not satisfied. For each  $r, s \in \mathbb{R}$  and  $\vec{u} = (a, b) \in V$  we have

$$(r + s)\vec{u} = (r + s)(a, b) = (a, (r + s)b) \quad (1)$$

$$r\vec{u} + s\vec{u} = (a, rb) + (a, sb) = (2a, (r + s)b) \quad (2)$$

Note that  $(1) \neq (2)$ .

# Vector Spaces

## Example

The nonempty set  $V = \mathbb{R}^2 = \{ \vec{x} = (a, b) \mid a, b \in \mathbb{R} \}$  together with the binary operations

$$+ : V \times V \longrightarrow V, \quad (a, b) + (c, d) = (b, d)$$

$$\cdot : \mathbb{R} \times V \longrightarrow V, \quad \lambda (a, b) = (\lambda a, \lambda b)$$

is not a vector space.

# Vector Spaces

## Example

The nonempty set  $V = \mathbb{R}^2 = \{ \vec{x} = (a, b) \mid a, b \in \mathbb{R} \}$  together with the binary operations

$$+ : V \times V \longrightarrow V, \quad (a, b) + (c, d) = (b, d)$$

$$\cdot : \mathbb{R} \times V \longrightarrow V, \quad \lambda(a, b) = (\lambda a, \lambda b)$$

is not a vector space. Because,  $V$  is not associative.



# Vector Spaces

## Example

The nonempty set  $V = \mathbb{R}^2 = \{ \vec{x} = (a, b) \mid a, b \in \mathbb{R} \}$  together with the binary operations

$$+ : V \times V \longrightarrow V, \quad (a, b) + (c, d) = (b, d)$$

$$\cdot : \mathbb{R} \times V \longrightarrow V, \quad \lambda(a, b) = (\lambda a, \lambda b)$$

is not a vector space. Because,  $V$  is not associative. Let  $\vec{x} = (a, b)$ ,  $\vec{y} = (c, d)$ ,  $\vec{z} = (e, f) \in V$ . We need to show that  $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$  is not satisfied.

# Vector Spaces

## Example

The nonempty set  $V = \mathbb{R}^2 = \{ \vec{x} = (a, b) \mid a, b \in \mathbb{R} \}$  together with the binary operations

$$+ : V \times V \longrightarrow V, \quad (a, b) + (c, d) = (b, d)$$

$$\cdot : \mathbb{R} \times V \longrightarrow V, \quad \lambda(a, b) = (\lambda a, \lambda b)$$

is not a vector space. Because,  $V$  is not associative. Let

$\vec{x} = (a, b)$ ,  $\vec{y} = (c, d)$ ,  $\vec{z} = (e, f) \in V$ . We need to show that  $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$  is not satisfied.

$$((a, b) + (c, d)) + (e, f)$$

# Vector Spaces

## Example

The nonempty set  $V = \mathbb{R}^2 = \{ \vec{x} = (a, b) \mid a, b \in \mathbb{R} \}$  together with the binary operations

$$+ : V \times V \longrightarrow V, \quad (a, b) + (c, d) = (b, d)$$

$$\cdot : \mathbb{R} \times V \longrightarrow V, \quad \lambda(a, b) = (\lambda a, \lambda b)$$

is not a vector space. Because,  $V$  is not associative. Let  $\vec{x} = (a, b)$ ,  $\vec{y} = (c, d)$ ,  $\vec{z} = (e, f) \in V$ . We need to show that  $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$  is not satisfied.

$$((a, b) + (c, d)) + (e, f) = (b, d) + (e, f) = (d, f)$$

# Vector Spaces

## Example

The nonempty set  $V = \mathbb{R}^2 = \{ \vec{x} = (a, b) \mid a, b \in \mathbb{R} \}$  together with the binary operations

$$+ : V \times V \longrightarrow V, \quad (a, b) + (c, d) = (b, d)$$

$$\cdot : \mathbb{R} \times V \longrightarrow V, \quad \lambda(a, b) = (\lambda a, \lambda b)$$

is not a vector space. Because,  $V$  is not associative. Let

$\vec{x} = (a, b)$ ,  $\vec{y} = (c, d)$ ,  $\vec{z} = (e, f) \in V$ . We need to show that  $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$  is not satisfied.

$$\begin{aligned} ((a, b) + (c, d)) + (e, f) &= (b, d) + (e, f) = (d, f) \\ (a, b) + ((c, d) + (e, f)) & \end{aligned} \quad (3)$$

# Vector Spaces

## Example

The nonempty set  $V = \mathbb{R}^2 = \{ \vec{x} = (a, b) \mid a, b \in \mathbb{R} \}$  together with the binary operations

$$+ : V \times V \longrightarrow V, \quad (a, b) + (c, d) = (b, d)$$

$$\cdot : \mathbb{R} \times V \longrightarrow V, \quad \lambda(a, b) = (\lambda a, \lambda b)$$

is not a vector space. Because,  $V$  is not associative. Let

$\vec{x} = (a, b)$ ,  $\vec{y} = (c, d)$ ,  $\vec{z} = (e, f) \in V$ . We need to show that  $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$  is not satisfied.

$$((a, b) + (c, d)) + (e, f) = (b, d) + (e, f) = (d, f) \quad (3)$$

$$(a, b) + ((c, d) + (e, f)) = (a, b) + (d, f) = (b, f)$$

# Vector Spaces

## Example

The nonempty set  $V = \mathbb{R}^2 = \{ \vec{x} = (a, b) \mid a, b \in \mathbb{R} \}$  together with the binary operations

$$+ : V \times V \longrightarrow V, \quad (a, b) + (c, d) = (b, d)$$

$$\cdot : \mathbb{R} \times V \longrightarrow V, \quad \lambda(a, b) = (\lambda a, \lambda b)$$

is not a vector space. Because,  $V$  is not associative. Let

$\vec{x} = (a, b)$ ,  $\vec{y} = (c, d)$ ,  $\vec{z} = (e, f) \in V$ . We need to show that  $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$  is not satisfied.

$$((a, b) + (c, d)) + (e, f) = (b, d) + (e, f) = (d, f) \quad (3)$$

$$(a, b) + ((c, d) + (e, f)) = (a, b) + (d, f) = (b, f) \quad (4)$$

Note that  $(3) \neq (4)$ .

# Vector Spaces

## Example

The set of column vectors  $P = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} \mid \begin{array}{l} x + y + z = 0 \\ x, y, z \in \mathbb{R} \end{array} \right\}$   
together with binary operations

# Vector Spaces

## Example

The set of column vectors  $P = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} \mid \begin{array}{l} x + y + z = 0 \\ x, y, z \in \mathbb{R} \end{array} \right\}$

together with binary operations

$$+ : \begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} + \begin{pmatrix} x_2 \\ y_2 \\ z_2 \end{pmatrix} = \begin{pmatrix} x_1 + x_2 \\ y_1 + y_2 \\ z_1 + z_2 \end{pmatrix}$$



# Vector Spaces

## Example

The set of column vectors  $P = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} \mid \begin{array}{l} x + y + z = 0 \\ x, y, z \in \mathbb{R} \end{array} \right\}$

together with binary operations

$$+ : \begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} + \begin{pmatrix} x_2 \\ y_2 \\ z_2 \end{pmatrix} = \begin{pmatrix} x_1 + x_2 \\ y_1 + y_2 \\ z_1 + z_2 \end{pmatrix}$$

$$\cdot : \lambda \begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} = \begin{pmatrix} \lambda x_1 \\ \lambda y_1 \\ \lambda z_1 \end{pmatrix}$$

# Vector Spaces

## Example

The set of column vectors  $P = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} \mid \begin{array}{l} x + y + z = 0 \\ x, y, z \in \mathbb{R} \end{array} \right\}$

together with binary operations

$$+ : \begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} + \begin{pmatrix} x_2 \\ y_2 \\ z_2 \end{pmatrix} = \begin{pmatrix} x_1 + x_2 \\ y_1 + y_2 \\ z_1 + z_2 \end{pmatrix}$$

$$\cdot : \lambda \begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} = \begin{pmatrix} \lambda x_1 \\ \lambda y_1 \\ \lambda z_1 \end{pmatrix}$$

is a vector space.

## Example

Let  $\mathcal{P}_3 = \{a_0 + a_1x + a_2x^2 + a_3x^3 \mid a_0, \dots, a_3 \in \mathbb{R}\}$  be the set of all polynomials of degree at most three in one variable.

# Vector Spaces

## Example

Let  $\mathcal{P}_3 = \{a_0 + a_1x + a_2x^2 + a_3x^3 \mid a_0, \dots, a_3 \in \mathbb{R}\}$  be the set of all polynomials of degree at most three in one variable.  $\mathcal{P}_3$  is a vector space with respect to polynomial addition

$$\begin{aligned} & (a_0 + a_1x + a_2x^2 + a_3x^3) + (b_0 + b_1x + b_2x^2 + b_3x^3) \\ = & (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 + (a_3 + b_3)x^3 \end{aligned}$$

# Vector Spaces

## Example

Let  $\mathcal{P}_3 = \{a_0 + a_1x + a_2x^2 + a_3x^3 \mid a_0, \dots, a_3 \in \mathbb{R}\}$  be the set of all polynomials of degree at most three in one variable.  $\mathcal{P}_3$  is a vector space with respect to polynomial addition

$$\begin{aligned} & (a_0 + a_1x + a_2x^2 + a_3x^3) + (b_0 + b_1x + b_2x^2 + b_3x^3) \\ &= (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 + (a_3 + b_3)x^3 \end{aligned}$$

and

$$r(a_0 + a_1x + a_2x^2 + a_3x^3) = (ra_0) + (ra_1)x + (ra_2)x^2 + (ra_3)x^3$$

# Vector Spaces

## Example

Let  $\mathcal{P}_3 = \{a_0 + a_1x + a_2x^2 + a_3x^3 \mid a_0, \dots, a_3 \in \mathbb{R}\}$  be the set of all polynomials of degree at most three in one variable.  $\mathcal{P}_3$  is a vector space with respect to polynomial addition

$$\begin{aligned} & (a_0 + a_1x + a_2x^2 + a_3x^3) + (b_0 + b_1x + b_2x^2 + b_3x^3) \\ &= (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 + (a_3 + b_3)x^3 \end{aligned}$$

and

$$r(a_0 + a_1x + a_2x^2 + a_3x^3) = (ra_0) + (ra_1)x + (ra_2)x^2 + (ra_3)x^3$$

scalar multiplication.

## Example

The set of  $2 \times 2$  square matrices

$\mathcal{M}_{2 \times 2}(\mathbb{R}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{R} \right\}$  is a vector space with respect to binary operations

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} + \begin{pmatrix} w & x \\ y & z \end{pmatrix} = \begin{pmatrix} a+w & b+x \\ c+y & d+z \end{pmatrix}$$

## Example

The set of  $2 \times 2$  square matrices

$\mathcal{M}_{2 \times 2}(\mathbb{R}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{R} \right\}$  is a vector space with respect to binary operations

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} + \begin{pmatrix} w & x \\ y & z \end{pmatrix} = \begin{pmatrix} a+w & b+x \\ c+y & d+z \end{pmatrix}$$

$$r \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} ra & rb \\ rc & rd \end{pmatrix}.$$



# Subspaces

---

# Subspaces

## Definition

Let  $V$  be a real vector space and  $\emptyset \neq U \subseteq V$ . Then  $U$  is a **subspace** of  $V$ , if  $U$  itself is a vector space over  $\mathbb{R}$  with respect to the operations of vector addition and scalar multiplication on  $V$ .

# Subspaces

## Definition

Let  $V$  be a real vector space and  $\emptyset \neq U \subseteq V$ . Then  $U$  is a **subspace** of  $V$ , if  $U$  itself is a vector space over  $\mathbb{R}$  with respect to the operations of vector addition and scalar multiplication on  $V$ .

## Theorem

*Let  $V$  be a real vector space and  $\emptyset \neq U \subseteq V$ . Then  $U$  is a subspace of  $V$  if the following three conditions hold:*

# Subspaces

## Definition

Let  $V$  be a real vector space and  $\emptyset \neq U \subseteq V$ . Then  $U$  is a **subspace** of  $V$ , if  $U$  itself is a vector space over  $\mathbb{R}$  with respect to the operations of vector addition and scalar multiplication on  $V$ .

## Theorem

*Let  $V$  be a real vector space and  $\emptyset \neq U \subseteq V$ . Then  $U$  is a subspace of  $V$  if the following three conditions hold:*

- 1. The zero vector  $0$  belong to  $U$ .*
- 2. For each  $\vec{v}, \vec{w} \in U$ ,  $\vec{v} + \vec{w} \in U$ .*
- 3. For each  $r \in \mathbb{R}$  and  $\vec{v} \in U$ ,  $r\vec{v} \in U$ .*

# Subspaces

## Definition

Let  $V$  be a real vector space and  $\emptyset \neq U \subseteq V$ . Then  $U$  is a **subspace** of  $V$ , if  $U$  itself is a vector space over  $\mathbb{R}$  with respect to the operations of vector addition and scalar multiplication on  $V$ .

## Theorem

*Let  $V$  be a real vector space and  $\emptyset \neq U \subseteq V$ . Then  $U$  is a subspace of  $V$  if the following three conditions hold:*

1. *The zero vector  $0$  belong to  $U$ .*
2. *For each  $\vec{v}, \vec{w} \in U$ ,  $\vec{v} + \vec{w} \in U$ .*
3. *For each  $r \in \mathbb{R}$  and  $\vec{v} \in U$ ,  $r\vec{v} \in U$ .*

**Note:** The three conditions given in above Theorem can be combined as a single one. That is,  $U$  is a subspace of  $V$  if for each  $\vec{v}, \vec{w} \in U$ , and  $r, s \in \mathbb{R}$ ,  $r\vec{v} + s\vec{w} \in U$ .

# Subspaces

## Example

Let  $\mathcal{B} = \left\{ \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \mid c = a + b \text{ and } a, b \in \mathbb{R} \right\} \subset \mathcal{M}_{2 \times 2}(\mathbb{R})$ . Is  $\mathcal{B}$  a subspace of  $\mathcal{M}_{2 \times 2}(\mathbb{R})$ ?

# Subspaces

## Example

Let  $\mathcal{B} = \left\{ \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \mid c = a + b \text{ and } a, b \in \mathbb{R} \right\} \subset \mathcal{M}_{2 \times 2}(\mathbb{R})$ . Is  $\mathcal{B}$  a subspace of  $\mathcal{M}_{2 \times 2}(\mathbb{R})$ ?

$$\begin{pmatrix} a & b \\ 0 & c \end{pmatrix} + \begin{pmatrix} x & y \\ 0 & z \end{pmatrix} = \underbrace{\begin{pmatrix} a+x & b+y \\ 0 & c+z \end{pmatrix}}_{c+z=a+b+x+y} \in \mathcal{B}$$

# Subspaces

## Example

Let  $\mathcal{B} = \left\{ \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \mid c = a + b \text{ and } a, b \in \mathbb{R} \right\} \subset \mathcal{M}_{2 \times 2}(\mathbb{R})$ . Is  $\mathcal{B}$  a subspace of  $\mathcal{M}_{2 \times 2}(\mathbb{R})$ ?

$$\begin{pmatrix} a & b \\ 0 & c \end{pmatrix} + \begin{pmatrix} x & y \\ 0 & z \end{pmatrix} = \underbrace{\begin{pmatrix} a+x & b+y \\ 0 & c+z \end{pmatrix}}_{c+z=a+b+x+y} \in \mathcal{B}$$

$$r \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} = \underbrace{\begin{pmatrix} ra & rb \\ 0 & rc \end{pmatrix}}_{rc=ra+rb} \in \mathcal{B}$$



# Subspaces

## Example

Let  $\mathcal{B} = \left\{ \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \mid c = a + b \text{ and } a, b \in \mathbb{R} \right\} \subset \mathcal{M}_{2 \times 2}(\mathbb{R})$ . Is  $\mathcal{B}$  a subspace of  $\mathcal{M}_{2 \times 2}(\mathbb{R})$ ?

$$\begin{pmatrix} a & b \\ 0 & c \end{pmatrix} + \begin{pmatrix} x & y \\ 0 & z \end{pmatrix} = \underbrace{\begin{pmatrix} a+x & b+y \\ 0 & c+z \end{pmatrix}}_{c+z=a+b+x+y} \in \mathcal{B}$$

$$r \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} = \underbrace{\begin{pmatrix} ra & rb \\ 0 & rc \end{pmatrix}}_{rc=ra+rb} \in \mathcal{B}$$

Also the zero of  $\mathcal{M}_{2 \times 2}(\mathbb{R})$ ,  $\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \in \mathcal{B}$ . Then  $\mathcal{B}$  is a subspace

## Example

Let  $\mathcal{A} = \{ (x, 0) \mid x \in \mathbb{R} \} \subset \mathbb{R}^2$ . Is  $\mathcal{A}$  a subspace of  $\mathbb{R}^2$ ?

## Example

Let  $\mathcal{A} = \{ (x, 0) \mid x \in \mathbb{R} \} \subset \mathbb{R}^2$ . Is  $\mathcal{A}$  a subspace of  $\mathbb{R}^2$ ? For each  $(a, 0), (b, 0) \in \mathcal{A}$  and  $r \in \mathbb{R}$

$$(a, 0) + (b, 0) = (a + b, 0) \in \mathcal{A}$$

## Example

Let  $\mathcal{A} = \{ (x, 0) \mid x \in \mathbb{R} \} \subset \mathbb{R}^2$ . Is  $\mathcal{A}$  a subspace of  $\mathbb{R}^2$ ? For each  $(a, 0), (b, 0) \in \mathcal{A}$  and  $r \in \mathbb{R}$

$$(a, 0) + (b, 0) = (a + b, 0) \in \mathcal{A}$$

$$r(a, 0) = (ra, 0) \in \mathcal{A}$$

## Example

Let  $\mathcal{A} = \{ (x, 0) \mid x \in \mathbb{R} \} \subset \mathbb{R}^2$ . Is  $\mathcal{A}$  a subspace of  $\mathbb{R}^2$ ? For each  $(a, 0), (b, 0) \in \mathcal{A}$  and  $r \in \mathbb{R}$

$$(a, 0) + (b, 0) = (a + b, 0) \in \mathcal{A}$$

$$r(a, 0) = (ra, 0) \in \mathcal{A}$$

Also the zero of  $\mathbb{R}^2$ ,  $(0, 0) \in \mathcal{A}$ . Then  $\mathcal{A}$  is a subspace of  $\mathbb{R}^2$ .

## Example

$\mathcal{W} = \{(a, b, c) \mid a \geq 0 \text{ and } a, b, c \in \mathbb{R}\} \subset \mathbb{R}^3$  is not a subspace of  $\mathbb{R}^3$ .

## Example

$\mathcal{W} = \{(a, b, c) \mid a \geq 0 \text{ and } a, b, c \in \mathbb{R}\} \subset \mathbb{R}^3$  is not a subspace of  $\mathbb{R}^3$ . Since  $\mathcal{W}$  is not closed under scalar multiplication. That is

$$(a, b, c) + (x, y, z) = (a + x, b + y, c + z) \in \mathcal{W}_{a+x \geq 0}$$

## Example

$\mathcal{W} = \{(a, b, c) \mid a \geq 0 \text{ and } a, b, c \in \mathbb{R}\} \subset \mathbb{R}^3$  is not a subspace of  $\mathbb{R}^3$ . Since  $\mathcal{W}$  is not closed under scalar multiplication. That is

$$(a, b, c) + (x, y, z) = (a + x, b + y, c + z) \in \mathcal{W}_{a+x \geq 0}$$

$$r(a, b, c) = \underbrace{(ra, rb, rc)}_{r < 0 \Rightarrow ra < 0} \notin \mathcal{W}.$$



## Example

$\mathcal{W}' = \{ (a, b, c) \mid a^2 + b^2 + c^2 \leq 1 \text{ and } a, b, c \in \mathbb{R} \} \subset \mathbb{R}^3$  is not a subspace of  $\mathbb{R}^3$ .

## Example

$\mathcal{W}' = \{ (a, b, c) \mid a^2 + b^2 + c^2 \leq 1 \text{ and } a, b, c \in \mathbb{R} \} \subset \mathbb{R}^3$  is not a subspace of  $\mathbb{R}^3$ . Because  $\mathcal{W}'$  is not closed under addition. For example,

$$\vec{u} = (1, 0, 0), \vec{v} = (0, 1, 0) \in \mathcal{W}'$$

## Example

$\mathcal{W}' = \{ (a, b, c) \mid a^2 + b^2 + c^2 \leq 1 \text{ and } a, b, c \in \mathbb{R} \} \subset \mathbb{R}^3$  is not a subspace of  $\mathbb{R}^3$ . Because  $\mathcal{W}'$  is not closed under addition. For example,

$$\begin{aligned}\vec{u} &= (1, 0, 0), \vec{v} = (0, 1, 0) \in \mathcal{W}' \\ \vec{u} + \vec{v} &= \underbrace{(1, 1, 0)}_{1^2+1^2=2>1} \notin \mathcal{W}'.\end{aligned}$$

?