



# MAT1320-Linear Algebra

## Lecture Notes

Eigenvalues and Eigenvectors, Characteristic Polynomial,  
Diagonalization, Cayley-Hamilton Theorem

---

Mehmet E. KÖROĞLU

Summer 2020

YILDIZ TECHNICAL UNIVERSITY, DEPARTMENT OF MATHEMATICS

*mkoroglu@yildiz.edu.tr*

# Table of contents

1. Eigenvalues and Eigenvectors
2. The Characteristic Equation
3. Finding Eigenvalues and Eigenvectors
4. Similar Matrices
5. Diagonalization
6. Cayley-Hamilton Theorem

# Eigenvalues and Eigenvectors

---

# Eigenvalues and Eigenvectors

## Definition

A scalar  $\lambda$  is called an **eigenvalue** of the  $n \times n$  square matrix  $A$  if there is a nontrivial solution  $\vec{x}$  of  $A\vec{x} = \lambda\vec{x}$ .

# Eigenvalues and Eigenvectors

## Definition

A scalar  $\lambda$  is called an **eigenvalue** of the  $n \times n$  square matrix  $A$  if there is a nontrivial solution  $\vec{x}$  of  $A\vec{x} = \lambda\vec{x}$ . Such an  $\vec{x}$  is called an **eigenvector** corresponding to the eigenvalue  $\lambda$ .

# Eigenvalues and Eigenvectors

## Definition

A scalar  $\lambda$  is called an **eigenvalue** of the  $n \times n$  square matrix  $A$  if there is a nontrivial solution  $\vec{x}$  of  $A\vec{x} = \lambda\vec{x}$ . Such an  $\vec{x}$  is called an **eigenvector** corresponding to the eigenvalue  $\lambda$ .

**Note:** Note that an eigenvector cannot be  $\vec{0}$ , but an eigenvalue can be  $0 \in \mathbb{R}$ . If 0 is an eigenvalue of  $A$ , then there must be some nontrivial vector  $\vec{x}$  for which  $A\vec{x} = 0\vec{x} = \vec{0}$  which implies that  $A$  is not invertible.

# Eigenvalues and Eigenvectors

## Definition

A scalar  $\lambda$  is called an **eigenvalue** of the  $n \times n$  square matrix  $A$  if there is a nontrivial solution  $\vec{x}$  of  $A\vec{x} = \lambda\vec{x}$ . Such an  $\vec{x}$  is called an **eigenvector** corresponding to the eigenvalue  $\lambda$ .

**Note:** Note that an eigenvector cannot be  $\vec{0}$ , but an eigenvalue can be  $0 \in \mathbb{R}$ . If 0 is an eigenvalue of  $A$ , then there must be some nontrivial vector  $\vec{x}$  for which  $A\vec{x} = 0\vec{x} = \vec{0}$  which implies that  $A$  is not invertible.

## Theorem

*The  $n \times n$  square matrix  $A$  is invertible if and only if 0 is not an eigenvalue of  $A$ .*

# Eigenvalues and Eigenvectors

## Definition

A scalar  $\lambda$  is called an **eigenvalue** of the  $n \times n$  square matrix  $A$  if there is a nontrivial solution  $\vec{x}$  of  $A\vec{x} = \lambda\vec{x}$ . Such an  $\vec{x}$  is called an **eigenvector** corresponding to the eigenvalue  $\lambda$ .

**Note:** Note that an eigenvector cannot be  $\vec{0}$ , but an eigenvalue can be  $0 \in \mathbb{R}$ . If 0 is an eigenvalue of  $A$ , then there must be some nontrivial vector  $\vec{x}$  for which  $A\vec{x} = 0\vec{x} = \vec{0}$  which implies that  $A$  is not invertible.

## Theorem

*The  $n \times n$  square matrix  $A$  is invertible if and only if 0 is not an eigenvalue of  $A$ .*

**Note:** The **eigenspace** of the  $n \times n$  matrix  $A$  corresponding to the eigenvalue  $\lambda$  of  $A$  is the set of all eigenvectors of  $A$  corresponding to  $\lambda$ .



# The Characteristic Equation

---

# The Characteristic Equation

Let  $A$  be an  $n \times n$  square matrix. Then we have

$$A\vec{x} = \lambda\vec{x}$$

# The Characteristic Equation

Let  $A$  be an  $n \times n$  square matrix. Then we have

$$A\vec{x} = \lambda\vec{x} \Rightarrow A\vec{x} - \lambda\vec{x} = \vec{0}$$

# The Characteristic Equation

Let  $A$  be an  $n \times n$  square matrix. Then we have

$$\begin{aligned} A\vec{x} &= \lambda\vec{x} \Rightarrow A\vec{x} - \lambda\vec{x} = \vec{0} \\ A\vec{x} - \lambda I_n \vec{x} &= \vec{0} \end{aligned}$$

# The Characteristic Equation

Let  $A$  be an  $n \times n$  square matrix. Then we have

$$\begin{aligned} A\vec{x} &= \lambda\vec{x} \Rightarrow A\vec{x} - \lambda\vec{x} = \vec{0} \\ A\vec{x} - \lambda I_n \vec{x} &= \vec{0} \Rightarrow (A - \lambda I_n) \vec{x} = \vec{0} \end{aligned}$$

# The Characteristic Equation

Let  $A$  be an  $n \times n$  square matrix. Then we have

$$\begin{aligned} A\vec{x} &= \lambda\vec{x} \Rightarrow A\vec{x} - \lambda\vec{x} = \vec{0} \\ A\vec{x} - \lambda I_n \vec{x} &= \vec{0} \Rightarrow (A - \lambda I_n) \vec{x} = \vec{0} \end{aligned}$$

**Note:** A scalar  $\lambda$  is an eigenvalue of an  $n \times n$  matrix  $A$  if and only if  $\lambda$  satisfies the equation  $\det(A - \lambda I) = 0$ .

# The Characteristic Equation

Let  $A$  be an  $n \times n$  square matrix. Then we have

$$\begin{aligned} A\vec{x} &= \lambda\vec{x} \Rightarrow A\vec{x} - \lambda\vec{x} = \vec{0} \\ A\vec{x} - \lambda I_n \vec{x} &= \vec{0} \Rightarrow (A - \lambda I_n) \vec{x} = \vec{0} \end{aligned}$$

**Note:** A scalar  $\lambda$  is an eigenvalue of an  $n \times n$  matrix  $A$  if and only if  $\lambda$  satisfies the equation  $\det(A - \lambda I) = 0$ .

## Definition

Let  $A$  be an  $n \times n$  square matrix.

# The Characteristic Equation

Let  $A$  be an  $n \times n$  square matrix. Then we have

$$\begin{aligned} A\vec{x} &= \lambda\vec{x} \Rightarrow A\vec{x} - \lambda\vec{x} = \vec{0} \\ A\vec{x} - \lambda I_n \vec{x} &= \vec{0} \Rightarrow (A - \lambda I_n) \vec{x} = \vec{0} \end{aligned}$$

**Note:** A scalar  $\lambda$  is an eigenvalue of an  $n \times n$  matrix  $A$  if and only if  $\lambda$  satisfies the equation  $\det(A - \lambda I) = 0$ .

## Definition

Let  $A$  be an  $n \times n$  square matrix. Then the equation  $\det(A - \lambda I_n) = 0$  is called the **characteristic equation** of the matrix  $A$  and the result of the determinant  $\det(A - \lambda I_n)$  is polynomial of the form  $P_A(\lambda) = \lambda^n + a_{n-1}\lambda^{n-1} + \dots + a_1\lambda + a_0$  is called **characteristic polynomial** of  $A$ .



# The Characteristic Equation

Let  $A$  be an  $n \times n$  square matrix. Then we have

$$\begin{aligned} A\vec{x} &= \lambda\vec{x} \Rightarrow A\vec{x} - \lambda\vec{x} = \vec{0} \\ A\vec{x} - \lambda I_n \vec{x} &= \vec{0} \Rightarrow (A - \lambda I_n) \vec{x} = \vec{0} \end{aligned}$$

**Note:** A scalar  $\lambda$  is an eigenvalue of an  $n \times n$  matrix  $A$  if and only if  $\lambda$  satisfies the equation  $\det(A - \lambda I) = 0$ .

## Definition

Let  $A$  be an  $n \times n$  square matrix. Then the equation  $\det(A - \lambda I_n) = 0$  is called the **characteristic equation** of the matrix  $A$  and the result of the determinant  $\det(A - \lambda I_n)$  is polynomial of the form  $P_A(\lambda) = \lambda^n + a_{n-1}\lambda^{n-1} + \dots + a_1\lambda + a_0$  is called **characteristic polynomial** of  $A$ .

**Note:**  $P_A(\lambda) = (-1)^n \lambda^n + (-1)^{n-1} \text{Tr}(A) \lambda^{n-1} + \dots + \det(A)$ .

For example, for a  $2 \times 2$  square matrix  $A$

Mehmet E. KÖROĞLU

$$P_A(\lambda) = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

# Finding Eigenvalues and Eigenvectors

---

# Finding Eigenvalues and Eigenvectors

1. Find characteristic polynomial  $|A - \lambda I_n| = P_A(\lambda)$  of  $A$ .

# Finding Eigenvalues and Eigenvectors

1. Find characteristic polynomial  $|A - \lambda I_n| = P_A(\lambda)$  of  $A$ .
2. Find the roots of  $P_A(\lambda)$  to find the eigenvalues of  $A$ .

# Finding Eigenvalues and Eigenvectors

1. Find characteristic polynomial  $|A - \lambda I_n| = P_A(\lambda)$  of  $A$ .
2. Find the roots of  $P_A(\lambda)$  to find the eigenvalues of  $A$ .
3. Repeat (a) and (b) for each eigenvalue  $\lambda$  of  $A$ .

# Finding Eigenvalues and Eigenvectors

1. Find characteristic polynomial  $|A - \lambda I_n| = P_A(\lambda)$  of  $A$ .
2. Find the roots of  $P_A(\lambda)$  to find the eigenvalues of  $A$ .
3. Repeat (a) and (b) for each eigenvalue  $\lambda$  of  $A$ .
  - (a) Form the matrix  $M = A - \lambda I$  by subtracting  $\lambda$  down the diagonal of  $A$ .

# Finding Eigenvalues and Eigenvectors

1. Find characteristic polynomial  $|A - \lambda I_n| = P_A(\lambda)$  of  $A$ .
2. Find the roots of  $P_A(\lambda)$  to find the eigenvalues of  $A$ .
3. Repeat (a) and (b) for each eigenvalue  $\lambda$  of  $A$ .
  - (a) Form the matrix  $M = A - \lambda I$  by subtracting  $\lambda$  down the diagonal of  $A$ .
  - (b) Find a basis for the solution space of the homogeneous system  $M\mathbf{X} = 0$ .

# Finding Eigenvalues and Eigenvectors

1. Find characteristic polynomial  $|A - \lambda I_n| = P_A(\lambda)$  of  $A$ .
2. Find the roots of  $P_A(\lambda)$  to find the eigenvalues of  $A$ .
3. Repeat (a) and (b) for each eigenvalue  $\lambda$  of  $A$ .
  - (a) Form the matrix  $M = A - \lambda I$  by subtracting  $\lambda$  down the diagonal of  $A$ .
  - (b) Find a basis for the solution space of the homogeneous system  $M\mathbf{X} = 0$ .

**Note 1:** The eigenvalues of a triangular matrix are the entries on its main diagonal.



# Finding Eigenvalues and Eigenvectors

1. Find characteristic polynomial  $|A - \lambda I_n| = P_A(\lambda)$  of  $A$ .
2. Find the roots of  $P_A(\lambda)$  to find the eigenvalues of  $A$ .
3. Repeat (a) and (b) for each eigenvalue  $\lambda$  of  $A$ .
  - (a) Form the matrix  $M = A - \lambda I$  by subtracting  $\lambda$  down the diagonal of  $A$ .
  - (b) Find a basis for the solution space of the homogeneous system  $M\mathbf{X} = 0$ .

**Note 1:** The eigenvalues of a triangular matrix are the entries on its main diagonal.

**Note 2:** If  $P_A(\lambda)$  has multiple roots, then there exists multiple eigenvalues.

# Finding Eigenvalues and Eigenvectors

## Example

Let  $A = \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix}$ . Find

1. characteristic polynomial,

# Finding Eigenvalues and Eigenvectors

## Example

Let  $A = \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix}$ . Find

1. characteristic polynomial,
2. eigenvalues,

# Finding Eigenvalues and Eigenvectors

## Example

Let  $A = \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix}$ . Find

1. characteristic polynomial,
2. eigenvalues,
3. eigenvectors.

# Finding Eigenvalues and Eigenvectors

## Solution (1)

$$P_A(\lambda) = \det(A - \lambda I_n)$$

$$A - \lambda I_3 = \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix} - \lambda \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

# Finding Eigenvalues and Eigenvectors

## Solution (1)

$$P_A(\lambda) = \det(A - \lambda I_n)$$

$$\begin{aligned} A - \lambda I_3 &= \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix} - \lambda \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 3 - \lambda & 6 & -8 \\ 0 & -\lambda & 6 \\ 0 & 0 & 2 - \lambda \end{pmatrix} \end{aligned}$$

# Finding Eigenvalues and Eigenvectors

## Solution (1)

$$P_A(\lambda) = \det(A - \lambda I_n)$$

$$\begin{aligned} A - \lambda I_3 &= \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix} - \lambda \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 3 - \lambda & 6 & -8 \\ 0 & -\lambda & 6 \\ 0 & 0 & 2 - \lambda \end{pmatrix} \\ \Rightarrow P_A(\lambda) &= -\lambda(3 - \lambda)(2 - \lambda). \end{aligned}$$

# Finding Eigenvalues and Eigenvectors

## Solution (2)

*Since, eigenvalues are the roots of  $P_A(\lambda)$ , we have*

$$P_A(\lambda) = -\lambda(3 - \lambda)(2 - \lambda) = 0$$



# Finding Eigenvalues and Eigenvectors

## Solution (2)

*Since, eigenvalues are the roots of  $P_A(\lambda)$ , we have*

$$\begin{aligned}P_A(\lambda) &= -\lambda(3-\lambda)(2-\lambda) = 0 \\ \Rightarrow \lambda_1 &= 0, \lambda_2 = 2, \lambda_3 = 3.\end{aligned}$$

## Finding Eigenvalues and Eigenvectors

### Solution (3)

For  $\lambda_1 = 0$ ,

$$(A - 0I_3) = A = \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

## Finding Eigenvalues and Eigenvectors

### Solution (3)

For  $\lambda_1 = 0$ ,

$$(A - 0I_3) = A = \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow A\vec{x} = \vec{0}$$

# Finding Eigenvalues and Eigenvectors

## Solution (3)

For  $\lambda_1 = 0$ ,

$$(A - 0I_3) = A = \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow A\vec{x} = \vec{0} \Rightarrow \begin{cases} x + 2y = 0 \\ z = 0 \end{cases}$$

## Finding Eigenvalues and Eigenvectors

### Solution (3)

For  $\lambda_1 = 0$ ,

$$(A - 0I_3) = A = \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow A\vec{x} = \vec{0} \Rightarrow \begin{cases} x + 2y = 0 \\ z = 0 \end{cases}$$

$$\Rightarrow \begin{pmatrix} x \\ y \\ z \end{pmatrix} = t \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix}, t \in \mathbb{R} \Rightarrow \vec{v}_1 = \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix}.$$

## Finding Eigenvalues and Eigenvectors

### Solution (3)

For  $\lambda_2 = 2$ ,

$$(A - 2I_3) = \begin{pmatrix} 1 & 6 & -8 \\ 0 & -2 & 6 \\ 0 & 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 10 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \end{pmatrix}$$

## Finding Eigenvalues and Eigenvectors

### Solution (3)

For  $\lambda_2 = 2$ ,

$$(A - 2I_3) = \begin{pmatrix} 1 & 6 & -8 \\ 0 & -2 & 6 \\ 0 & 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 10 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow (A - 2I_n) \vec{x} = \vec{0}$$

## Finding Eigenvalues and Eigenvectors

### Solution (3)

For  $\lambda_2 = 2$ ,

$$(A - 2I_3) = \begin{pmatrix} 1 & 6 & -8 \\ 0 & -2 & 6 \\ 0 & 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 10 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow (A - 2I_n) \vec{x} = \vec{0} \Rightarrow \begin{cases} x + 10z = 0 \\ y - 3z = 0 \end{cases}$$



## Finding Eigenvalues and Eigenvectors

### Solution (3)

For  $\lambda_2 = 2$ ,

$$(A - 2I_3) = \begin{pmatrix} 1 & 6 & -8 \\ 0 & -2 & 6 \\ 0 & 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 10 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow (A - 2I_n) \vec{x} = \vec{0} \Rightarrow \begin{cases} x + 10z = 0 \\ y - 3z = 0 \end{cases}$$

$$\Rightarrow \begin{pmatrix} x \\ y \\ z \end{pmatrix} = t \begin{pmatrix} -10 \\ 3 \\ 1 \end{pmatrix}, t \in \mathbb{R} \Rightarrow \vec{v}_2 = \begin{pmatrix} -10 \\ 3 \\ 1 \end{pmatrix}.$$

## Finding Eigenvalues and Eigenvectors

### Solution (3)

For  $\lambda_3 = 3$ ,

$$(A - 3I_3) = A = \begin{pmatrix} 0 & 6 & -8 \\ 0 & -3 & 6 \\ 0 & 0 & -1 \end{pmatrix} \sim \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

## Finding Eigenvalues and Eigenvectors

### Solution (3)

For  $\lambda_3 = 3$ ,

$$(A - 3I_3) = A - \begin{pmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{pmatrix} = \begin{pmatrix} 0 & 6 & -8 \\ 0 & -3 & 6 \\ 0 & 0 & -1 \end{pmatrix} \sim \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow (A - 3I_n) \vec{x} = \vec{0}$$

## Finding Eigenvalues and Eigenvectors

### Solution (3)

For  $\lambda_3 = 3$ ,

$$(A - 3I_3) = A - \begin{pmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{pmatrix} = \begin{pmatrix} 0 & 6 & -8 \\ 0 & -3 & 6 \\ 0 & 0 & -1 \end{pmatrix} \sim \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow (A - 3I_n) \vec{x} = \vec{0} \Rightarrow \begin{cases} y = 0 \\ z = 0 \end{cases}$$

## Finding Eigenvalues and Eigenvectors

### Solution (3)

For  $\lambda_3 = 3$ ,

$$(A - 3I_3) = A - \begin{pmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{pmatrix} = \begin{pmatrix} 0 & 6 & -8 \\ 0 & -3 & 6 \\ 0 & 0 & -1 \end{pmatrix} \sim \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow (A - 3I_n) \vec{x} = \vec{0} \Rightarrow \begin{cases} y = 0 \\ z = 0 \end{cases}$$

$$\Rightarrow \begin{pmatrix} x \\ y \\ z \end{pmatrix} = t \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, t \in \mathbb{R} \Rightarrow \vec{v}_3 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}.$$

# Similar Matrices

---

## Definition

The  $n \times n$  matrices  $A$  and  $B$  are said to be **similar** if there is an invertible  $n \times n$  matrix  $P$  such that  $A = PBP^{-1}$ .

## Definition

The  $n \times n$  matrices  $A$  and  $B$  are said to be **similar** if there is an invertible  $n \times n$  matrix  $P$  such that  $A = PBP^{-1}$ .

**Note:** If  $n \times n$  matrices are similar, then they have the same characteristic polynomial and hence the same eigenvalues (with the same multiplicities).



# Similar Matrices

## Definition

The  $n \times n$  matrices  $A$  and  $B$  are said to be **similar** if there is an invertible  $n \times n$  matrix  $P$  such that  $A = PBP^{-1}$ .

**Note:** If  $n \times n$  matrices are similar, then they have the same characteristic polynomial and hence the same eigenvalues (with the same multiplicities).

## Definition

A square matrix  $A$  is said to be **diagonalizable** if it is similar to a diagonal matrix. In other words, a diagonal matrix  $A$  has the property that there exists an invertible matrix  $P$  and a diagonal matrix  $D$  such that  $A = PDP^{-1}$ .

# Similar Matrices

## Definition

The  $n \times n$  matrices  $A$  and  $B$  are said to be **similar** if there is an invertible  $n \times n$  matrix  $P$  such that  $A = PBP^{-1}$ .

**Note:** If  $n \times n$  matrices are similar, then they have the same characteristic polynomial and hence the same eigenvalues (with the same multiplicities).

## Definition

A square matrix  $A$  is said to be **diagonalizable** if it is similar to a diagonal matrix. In other words, a diagonal matrix  $A$  has the property that there exists an invertible matrix  $P$  and a diagonal matrix  $D$  such that  $A = PDP^{-1}$ .

**Note:**  $A = PDP^{-1} \Rightarrow A^k = PD^kP^{-1}$

# Diagonalization

---

## Theorem (The Diagonalization Theorem)

1. *An  $n \times n$  matrix  $A$  is diagonalizable if and only if  $A$  has  $n$  linearly independent eigenvectors.*

## Theorem (The Diagonalization Theorem)

1. An  $n \times n$  matrix  $A$  is diagonalizable if and only if  $A$  has  $n$  linearly independent eigenvectors.
2. If  $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$  are linearly independent eigenvectors of  $A$  and  $\lambda_1, \lambda_2, \dots, \lambda_n$  are their corresponding eigenvalues, then  $A = PDP^{-1}$ , where

$$P = \begin{pmatrix} \vec{v}_1 & \dots & \vec{v}_n \end{pmatrix} \text{ and } D = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n \end{pmatrix}.$$

# Diagonalization

## Example

We have found the eigenvalues and corresponding eigenvectors of

the matrix  $A = \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix}$  as  $\lambda_1 = 0, \lambda_2 = 2, \lambda_3 = 3$  and

$$\vec{v}_1 = \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix}, \vec{v}_2 = \begin{pmatrix} -10 \\ 3 \\ 1 \end{pmatrix}, \vec{v}_3 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}. \text{ Then}$$

1. show that  $\vec{v}_1, \vec{v}_2, \vec{v}_3$  are linear independent,

# Diagonalization

## Example

We have found the eigenvalues and corresponding eigenvectors of

the matrix  $A = \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix}$  as  $\lambda_1 = 0, \lambda_2 = 2, \lambda_3 = 3$  and

$$\vec{v}_1 = \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix}, \vec{v}_2 = \begin{pmatrix} -10 \\ 3 \\ 1 \end{pmatrix}, \vec{v}_3 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}. \text{ Then}$$

1. show that  $\vec{v}_1, \vec{v}_2, \vec{v}_3$  are linear independent,
2. show that there exists matrices  $P$  and  $D$  such that  $A = PDP^{-1}$ ,

# Diagonalization

## Example

We have found the eigenvalues and corresponding eigenvectors of

the matrix  $A = \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix}$  as  $\lambda_1 = 0, \lambda_2 = 2, \lambda_3 = 3$  and

$$\vec{v}_1 = \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix}, \vec{v}_2 = \begin{pmatrix} -10 \\ 3 \\ 1 \end{pmatrix}, \vec{v}_3 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}. \text{ Then}$$

1. show that  $\vec{v}_1, \vec{v}_2, \vec{v}_3$  are linear independent,
2. show that there exists matrices  $P$  and  $D$  such that  $A = PDP^{-1}$ ,
3. and find  $A^{125}$ .



## Solution

$$1. P = \left( \begin{array}{ccc} \vec{v}_1 & \vec{v}_2 & \vec{v}_3 \end{array} \right) = \begin{pmatrix} -2 & -10 & 1 \\ 1 & 3 & 0 \\ 0 & 1 & 0 \end{pmatrix} \text{ and since}$$
$$\det(P) = 1 \neq 0, \vec{v}_1, \vec{v}_2, \vec{v}_3 \text{ are linearly independent.}$$

## Solution

$$1. P = \begin{pmatrix} \vec{v}_1 & \vec{v}_2 & \vec{v}_3 \end{pmatrix} = \begin{pmatrix} -2 & -10 & 1 \\ 1 & 3 & 0 \\ 0 & 1 & 0 \end{pmatrix} \text{ and since}$$
$$\det(P) = 1 \neq 0, \vec{v}_1, \vec{v}_2, \vec{v}_3 \text{ are linearly independent.}$$

$$2. P = \begin{pmatrix} -2 & -10 & 1 \\ 1 & 3 & 0 \\ 0 & 1 & 0 \end{pmatrix}, P^{-1} = \begin{pmatrix} 0 & 1 & -3 \\ 0 & 0 & 1 \\ 1 & 2 & 4 \end{pmatrix} \text{ and}$$

$$D = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix} \Rightarrow A = PDP^{-1}.$$

## Solution

$$1. P = \begin{pmatrix} \vec{v}_1 & \vec{v}_2 & \vec{v}_3 \end{pmatrix} = \begin{pmatrix} -2 & -10 & 1 \\ 1 & 3 & 0 \\ 0 & 1 & 0 \end{pmatrix} \text{ and since}$$

$\det(P) = 1 \neq 0$ ,  $\vec{v}_1, \vec{v}_2, \vec{v}_3$  are linearly independent.

$$2. P = \begin{pmatrix} -2 & -10 & 1 \\ 1 & 3 & 0 \\ 0 & 1 & 0 \end{pmatrix}, P^{-1} = \begin{pmatrix} 0 & 1 & -3 \\ 0 & 0 & 1 \\ 1 & 2 & 4 \end{pmatrix} \text{ and}$$

$$D = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix} \Rightarrow A = PDP^{-1}.$$

$$3. A^{125} = PD^{125}P^{-1} = P \begin{pmatrix} 0 & 0 & 0 \\ 0 & 2^{125} & 0 \\ 0 & 0 & 3^{125} \end{pmatrix} P^{-1}.$$

# Cayley-Hamilton Theorem

---

# Cayley-Hamilton Theorem

## Theorem (Cayley-Hamilton Theorem)

*Every matrix  $A$  is a root of its characteristic polynomial, i.e.,*

$$P_A(A) = O_{n \times n}.$$

# Cayley-Hamilton Theorem

## Theorem (Cayley-Hamilton Theorem)

*Every matrix  $A$  is a root of its characteristic polynomial, i.e.,*

$$P_A(A) = O_{n \times n}.$$

**Note:** The inverse or any power of a square matrix can be computed by using Cayley-Hamilton Theorem.

# Cayley-Hamilton Theorem

## Theorem (Cayley-Hamilton Theorem)

*Every matrix  $A$  is a root of its characteristic polynomial, i.e.,*

$$P_A(A) = O_{n \times n}.$$

**Note:** The inverse or any power of a square matrix can be computed by using Cayley-Hamilton Theorem.

$$P_A(A) = A^n + a_{n-1}A^{n-1} + \dots + a_1A + a_0I_n = 0$$

# Cayley-Hamilton Theorem

## Theorem (Cayley-Hamilton Theorem)

*Every matrix  $A$  is a root of its characteristic polynomial, i.e.,*

$$P_A(A) = O_{n \times n}.$$

**Note:** The inverse or any power of a square matrix can be computed by using Cayley-Hamilton Theorem.

$$\begin{aligned} P_A(A) &= A^n + a_{n-1}A^{n-1} + \dots + a_1A + a_0I_n = 0 \\ \Rightarrow A^n &= -\left(a_{n-1}A^{n-1} + \dots + a_1A + a_0I_n\right) \end{aligned}$$



# Cayley-Hamilton Theorem

## Theorem (Cayley-Hamilton Theorem)

Every matrix  $A$  is a root of its characteristic polynomial, i.e.,

$$P_A(A) = O_{n \times n}.$$

**Note:** The inverse or any power of a square matrix can be computed by using Cayley-Hamilton Theorem.

$$\begin{aligned} P_A(A) &= A^n + a_{n-1}A^{n-1} + \dots + a_1A + a_0I_n = 0 \\ \Rightarrow A^n &= -\left(a_{n-1}A^{n-1} + \dots + a_1A + a_0I_n\right) \end{aligned}$$

If  $a_0 \neq 0$ , then

$$I_n = A \underbrace{\frac{-1}{a_0} \left( A^{n-1} + a_{n-1}A^{n-2} + \dots + a_1I_n \right)}_{A^{-1}}$$

## Example

### Example

Let  $A = \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}$ . Find

1. characteristic polynomial,

# Example

## Example

Let  $A = \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}$ . Find

1. characteristic polynomial,
2. eigenvalues,

# Example

## Example

Let  $A = \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}$ . Find

1. characteristic polynomial,
2. eigenvalues,
3. eigenvectors,

# Example

## Example

Let  $A = \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}$ . Find

1. characteristic polynomial,
2. eigenvalues,
3. eigenvectors,
4. matrices  $P$  and  $D$  such that  $A = PDP^{-1}$ , if any,

## Example

### Example

Let  $A = \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}$ . Find

1. characteristic polynomial,
2. eigenvalues,
3. eigenvectors,
4. matrices  $P$  and  $D$  such that  $A = PDP^{-1}$ , if any,
5.  $A^{-1}$  and  $A^5$  (by using Cayley-Hamilton Theorem).

# Finding Eigenvalues and Eigenvectors

## **Solution (1)**

$$P_A(\lambda) = \det(A - \lambda I_n)$$

$$A - \lambda I_2 = \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} - \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

# Finding Eigenvalues and Eigenvectors

## Solution (1)

$$P_A(\lambda) = \det(A - \lambda I_n)$$

$$\begin{aligned} A - \lambda I_2 &= \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} - \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 - \lambda & 4 \\ 2 & 3 - \lambda \end{pmatrix} \end{aligned}$$



# Finding Eigenvalues and Eigenvectors

## Solution (1)

$$P_A(\lambda) = \det(A - \lambda I_n)$$

$$\begin{aligned} A - \lambda I_2 &= \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} - \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 - \lambda & 4 \\ 2 & 3 - \lambda \end{pmatrix} \\ \Rightarrow P_A(\lambda) &= (1 - \lambda)(3 - \lambda) - 8 \\ &= \lambda^2 - 4\lambda - 5. \end{aligned}$$

# Finding Eigenvalues and Eigenvectors

## Solution (2)

*The eigenvalues are the roots of  $P_A(\lambda)$  such that*

$$P_A(\lambda) = \lambda^2 - 4\lambda - 5 = 0$$

# Finding Eigenvalues and Eigenvectors

## Solution (2)

*The eigenvalues are the roots of  $P_A(\lambda)$  such that*

$$\begin{aligned}P_A(\lambda) &= \lambda^2 - 4\lambda - 5 = 0 \\ \Rightarrow \lambda_1 &= -1, \lambda_2 = 5.\end{aligned}$$

## Finding Eigenvalues and Eigenvectors

**Solution (3)**

For  $\lambda_1 = -1$ ,

$$(A + I_2) = \begin{pmatrix} 2 & 4 \\ 2 & 4 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 \\ 0 & 0 \end{pmatrix}$$

## Finding Eigenvalues and Eigenvectors

### Solution (3)

For  $\lambda_1 = -1$ ,

$$(A + I_2) = \begin{pmatrix} 2 & 4 \\ 2 & 4 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 \\ 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \end{pmatrix} \Rightarrow (A + I_2) \vec{x} = \vec{0}$$

## Finding Eigenvalues and Eigenvectors

### Solution (3)

For  $\lambda_1 = -1$ ,

$$(A + I_2) = \begin{pmatrix} 2 & 4 \\ 2 & 4 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 \\ 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \end{pmatrix} \Rightarrow (A + I_2) \vec{x} = \vec{0} \Rightarrow x + 2y = 0$$

## Finding Eigenvalues and Eigenvectors

### Solution (3)

For  $\lambda_1 = -1$ ,

$$(A + I_2) = \begin{pmatrix} 2 & 4 \\ 2 & 4 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 \\ 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \end{pmatrix} \Rightarrow (A + I_2) \vec{x} = \vec{0} \Rightarrow x + 2y = 0$$

$$\Rightarrow \begin{pmatrix} x \\ y \end{pmatrix} = t \begin{pmatrix} -2 \\ 1 \end{pmatrix}, t \in \mathbb{R} \Rightarrow \vec{v}_1 = \begin{pmatrix} -2 \\ 1 \end{pmatrix}.$$

## Finding Eigenvalues and Eigenvectors

### **Solution (3)**

For  $\lambda_2 = 5$ ,

$$(A - 5I_2) = \begin{pmatrix} -4 & 4 \\ 2 & -2 \end{pmatrix} \sim \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix}$$



## Finding Eigenvalues and Eigenvectors

### Solution (3)

For  $\lambda_2 = 5$ ,

$$(A - 5I_2) = \begin{pmatrix} -4 & 4 \\ 2 & -2 \end{pmatrix} \sim \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix}$$
$$\vec{x} = \begin{pmatrix} x \\ y \end{pmatrix} \Rightarrow (A - 5I_2) \vec{x} = \vec{0}$$

## Finding Eigenvalues and Eigenvectors

### Solution (3)

For  $\lambda_2 = 5$ ,

$$(A - 5I_2) = \begin{pmatrix} -4 & 4 \\ 2 & -2 \end{pmatrix} \sim \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \end{pmatrix} \Rightarrow (A - 5I_2) \vec{x} = \vec{0} \Rightarrow x - y = 0$$

## Finding Eigenvalues and Eigenvectors

### Solution (3)

For  $\lambda_2 = 5$ ,

$$(A - 5I_2) = \begin{pmatrix} -4 & 4 \\ 2 & -2 \end{pmatrix} \sim \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \end{pmatrix} \Rightarrow (A - 5I_2) \vec{x} = \vec{0} \Rightarrow x - y = 0$$

$$\Rightarrow \begin{pmatrix} x \\ y \end{pmatrix} = t \begin{pmatrix} 1 \\ 1 \end{pmatrix}, t \in \mathbb{R} \Rightarrow \vec{v}_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

## Finding Eigenvalues and Eigenvectors

**Solution (4)**

$$P = \begin{pmatrix} -2 & 1 \\ 1 & 1 \end{pmatrix}, P^{-1} = \begin{pmatrix} -\frac{1}{3} & \frac{1}{3} \\ \frac{1}{3} & \frac{2}{3} \end{pmatrix} \text{ and } D = \begin{pmatrix} -1 & 0 \\ 0 & 5 \end{pmatrix} \Rightarrow$$

$$A = PDP^{-1}.$$

# Cayley-Hamilton Theorem

## Solution (5)

$$P_A(A) = A^2 - 4A - 5I_2 = 0$$

# Cayley-Hamilton Theorem

## Solution (5)

$$\begin{aligned}P_A(A) &= A^2 - 4A - 5I_2 = 0 \\ \Rightarrow \underbrace{A \frac{1}{5} (A - 4I_2)}_{A^{-1}} &= I_2\end{aligned}$$

# Cayley-Hamilton Theorem

## Solution (5)

$$P_A(A) = A^2 - 4A - 5I_2 = 0$$

$$\Rightarrow \underbrace{A \frac{1}{5} (A - 4I_2)}_{A^{-1}} = I_2$$

$$A^{-1} = \frac{1}{5} \left( \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} - \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix} \right) = \frac{1}{5} \begin{pmatrix} -3 & 4 \\ 2 & -1 \end{pmatrix}$$

# Cayley-Hamilton Theorem

## Solution (5)

$$P_A(A) = A^2 - 4A - 5I_2 = 0$$

$$\Rightarrow \underbrace{A \frac{1}{5} (A - 4I_2)}_{A^{-1}} = I_2$$

$$A^{-1} = \frac{1}{5} \left( \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} - \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix} \right) = \frac{1}{5} \begin{pmatrix} -3 & 4 \\ 2 & -1 \end{pmatrix}$$

$$A^2 = 4A + 5I_2$$



# Cayley-Hamilton Theorem

## Solution (5)

$$P_A(A) = A^2 - 4A - 5I_2 = 0$$

$$\Rightarrow \underbrace{A \frac{1}{5} (A - 4I_2)}_{A^{-1}} = I_2$$

$$A^{-1} = \frac{1}{5} \left( \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} - \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix} \right) = \frac{1}{5} \begin{pmatrix} -3 & 4 \\ 2 & -1 \end{pmatrix}$$

$$A^2 = 4A + 5I_2$$

$$A^4 = (4A + 5I_2)(4A + 5I_2)$$

# Cayley-Hamilton Theorem

## Solution (5)

$$P_A(A) = A^2 - 4A - 5I_2 = 0$$

$$\Rightarrow \underbrace{A \frac{1}{5} (A - 4I_2)}_{A^{-1}} = I_2$$

$$A^{-1} = \frac{1}{5} \left( \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} - \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix} \right) = \frac{1}{5} \begin{pmatrix} -3 & 4 \\ 2 & -1 \end{pmatrix}$$

$$A^2 = 4A + 5I_2$$

$$A^4 = (4A + 5I_2)(4A + 5I_2)$$

$$A^5 = A(4A + 5I_2)(4A + 5I_2)$$

?