



MAT1320-Linear Algebra

Lecture Notes

Matrices

Mehmet E. KÖROĞLU
Summer 2020

YILDIZ TECHNICAL UNIVERSITY, DEPARTMENT OF MATHEMATICS
mkoroglu@yildiz.edu.tr

Table of contents

1. Matrices
2. Matrix Addition and Scalar Multiplication
3. Summation Symbol
4. Matrix Multiplication
5. Transpose of a Matrix
6. Square Matrices
7. Diagonal and Trace
8. Identity Matrix, Scalar Matrices
9. Powers of Matrices, Polynomials in Matrices
10. Invertible (Nonsingular) Matrices

Matrices

Matrices

A **matrix** A over a field K or, simply, a matrix A (when K is implicit) is a rectangular array of scalars usually presented in the following form:

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}$$

Matrices

A **matrix** A over a field K or, simply, a matrix A (when K is implicit) is a rectangular array of scalars usually presented in the following form:

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}$$

The rows of such a matrix A are the m horizontal lists of scalars:

$$(a_{11} \ a_{12} \ \dots \ a_{1n}), \dots, (a_{m1} \ a_{m2} \ \dots \ a_{mn})$$

Matrices

A **matrix** A over a field K or, simply, a matrix A (when K is implicit) is a rectangular array of scalars usually presented in the following form:

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}$$

The rows of such a matrix A are the m horizontal lists of scalars:

$$(a_{11} \ a_{12} \ \dots \ a_{1n}), \dots, (a_{m1} \ a_{m2} \ \dots \ a_{mn})$$

and the columns of A are the n vertical lists of scalars:

$$\begin{pmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{pmatrix}, \dots, \begin{pmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{pmatrix}$$

- Note that the element a_{ij} , called the ij -entry or ij -element, appears in row i and column j . We frequently denote such a matrix by simply writing $A = [a_{ij}]$.

- Note that the element a_{ij} , called the *ij*-entry or *ij*-element, appears in row i and column j . We frequently denote such a matrix by simply writing $A = [a_{ij}]$.
- A matrix with m rows and n columns is called an m by n matrix, written $m \times n$. The pair of numbers m and n is called the *size* of the matrix.

Matrices

- Note that the element a_{ij} , called the *ij*-entry or *ij*-element, appears in row i and column j . We frequently denote such a matrix by simply writing $A = [a_{ij}]$.
- A matrix with m rows and n columns is called an m by n matrix, written $m \times n$. The pair of numbers m and n is called the *size* of the matrix.
- Two matrices A and B are *equal*, written $A = B$, if they have the same size and if corresponding elements are equal. Thus, the equality of two $m \times n$ matrices is equivalent to a system of mn equalities, one for each corresponding pair of elements.

- A matrix with only one row is called a row matrix or row vector, and a matrix with only one column is called a column matrix or column vector.

Matrices

- A matrix with only one row is called a **row matrix** or **row vector**, and a matrix with only one column is called a **column matrix** or **column vector**.
- A matrix whose entries are all zero is called a **zero matrix** and will usually be denoted by 0 .

Matrices

- A matrix with only one row is called a **row matrix** or **row vector**, and a matrix with only one column is called a **column matrix** or **column vector**.
- A matrix whose entries are all zero is called a **zero matrix** and will usually be denoted by 0 .
- **Matrices whose entries are all real numbers are called real matrices and are said to be matrices over \mathbb{R} .**

Matrices

- A matrix with only one row is called a **row matrix** or **row vector**, and a matrix with only one column is called a **column matrix** or **column vector**.
- A matrix whose entries are all zero is called a **zero matrix** and will usually be denoted by 0 .
- Matrices whose entries are all real numbers are called **real matrices** and are said to be matrices over \mathbb{R} .
- Analogously, matrices whose entries are all complex numbers are called **complex matrices** and are said to be matrices over \mathbb{C} . This text will be mainly concerned with such real matrices.

Example

The rectangular array $A = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix}$ is a 2×3 matrix.

Example

The rectangular array $A = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix}$ is a 2×3 matrix. Its rows are

$$(1 \ 2 \ 3) \text{ and } (3 \ 1 \ 2),$$

Example

The rectangular array $A = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix}$ is a 2×3 matrix. Its rows are

$$(1 \ 2 \ 3) \text{ and } (3 \ 1 \ 2),$$

and its columns are

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 3 \\ 2 \end{pmatrix}.$$

Example

The rectangular array $A = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix}$ is a 2×3 matrix. Its rows are

$$(1 \ 2 \ 3) \text{ and } (3 \ 1 \ 2),$$

and its columns are

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 3 \\ 2 \end{pmatrix}.$$

Example

The 2×4 zero matrix is the matrix $0 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$.

Example

Find x, y, z, t such that
$$\begin{pmatrix} x + y & 3z + t \\ x - y & z - t \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 4 & 3 \end{pmatrix}.$$

Example

Find x, y, z, t such that $\begin{pmatrix} x+y & 3z+t \\ x-y & z-t \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 4 & 3 \end{pmatrix}$. By definition of equality of matrices, the four corresponding entries must be equal.

Example

Find x, y, z, t such that $\begin{pmatrix} x+y & 3z+t \\ x-y & z-t \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 4 & 3 \end{pmatrix}$. By definition of equality of matrices, the four corresponding entries must be equal. Thus,

$$x + y = 2, \quad 3z + t = 1,$$

$$x - y = 4, \quad z - t = 3.$$

Example

Find x, y, z, t such that $\begin{pmatrix} x+y & 3z+t \\ x-y & z-t \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 4 & 3 \end{pmatrix}$. By definition of equality of matrices, the four corresponding entries must be equal. Thus,

$$x+y = 2, \quad 3z+t = 1,$$

$$x-y = 4, \quad z-t = 3.$$

Solving the above system of equations yields

$$x = 3, y = -1, z = 1 \text{ and } t = -2.$$

Matrix Addition and Scalar Multiplication

Matrix Addition and Scalar Multiplication

Let $A = [a_{ij}]$ and $B = [b_{ij}]$ be two matrices with the same size, say $m \times n$ matrices.

Matrix Addition and Scalar Multiplication

Let $A = [a_{ij}]$ and $B = [b_{ij}]$ be two matrices with the same size, say $m \times n$ matrices. The **sum** of A and B , written $A + B$, is the matrix obtained by adding corresponding elements from A and B .

Matrix Addition and Scalar Multiplication

Let $A = [a_{ij}]$ and $B = [b_{ij}]$ be two matrices with the same size, say $m \times n$ matrices. The **sum** of A and B , written $A + B$, is the matrix obtained by adding corresponding elements from A and B . That is,

$$A + B = \begin{pmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \dots & a_{1n} + b_{1n} \\ a_{21} + b_{21} & a_{22} + b_{22} & \dots & a_{2n} + b_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} + b_{m1} & a_{m2} + b_{m2} & \dots & a_{mn} + b_{mn} \end{pmatrix}.$$

Matrix Addition and Scalar Multiplication

The product of the matrix A by a scalar k , written $k.A$ or simply kA , is the matrix obtained by multiplying each element of A by k .

Matrix Addition and Scalar Multiplication

The **product of the matrix A by a scalar k** , written $k.A$ or simply kA , is the matrix obtained by multiplying each element of A by k . That is,

$$kA = \begin{pmatrix} ka_{11} & ka_{12} & \dots & ka_{1n} \\ ka_{21} & ka_{22} & \dots & ka_{2n} \\ \dots & \dots & \dots & \dots \\ ka_{m1} & ka_{m2} & \dots & ka_{mn} \end{pmatrix}.$$

Matrix Addition and Scalar Multiplication

The **product of the matrix A by a scalar k** , written $k.A$ or simply kA , is the matrix obtained by multiplying each element of A by k . That is,

$$kA = \begin{pmatrix} ka_{11} & ka_{12} & \dots & ka_{1n} \\ ka_{21} & ka_{22} & \dots & ka_{2n} \\ \dots & \dots & \dots & \dots \\ ka_{m1} & ka_{m2} & \dots & ka_{mn} \end{pmatrix}.$$

Notes:

- Observe that $A + B$ and kA are also $m \times n$ matrices.

Matrix Addition and Scalar Multiplication

The **product of the matrix A by a scalar k** , written $k.A$ or simply kA , is the matrix obtained by multiplying each element of A by k . That is,

$$kA = \begin{pmatrix} ka_{11} & ka_{12} & \dots & ka_{1n} \\ ka_{21} & ka_{22} & \dots & ka_{2n} \\ \dots & \dots & \dots & \dots \\ ka_{m1} & ka_{m2} & \dots & ka_{mn} \end{pmatrix}.$$

Notes:

- Observe that $A + B$ and kA are also $m \times n$ matrices.
- We also define $-A = (-1)A$ and $A - B = A + (-1)B$. The matrix $-A$ is called the negative of the matrix A , and the matrix $A - B$ is called the **difference** of A and B .

Matrix Addition and Scalar Multiplication

The **product of the matrix A by a scalar k** , written $k.A$ or simply kA , is the matrix obtained by multiplying each element of A by k . That is,

$$kA = \begin{pmatrix} ka_{11} & ka_{12} & \dots & ka_{1n} \\ ka_{21} & ka_{22} & \dots & ka_{2n} \\ \dots & \dots & \dots & \dots \\ ka_{m1} & ka_{m2} & \dots & ka_{mn} \end{pmatrix}.$$

Notes:

- Observe that $A + B$ and kA are also $m \times n$ matrices.
- We also define $-A = (-1)A$ and $A - B = A + (-1)B$. The matrix $-A$ is called the negative of the matrix A , and the matrix $A - B$ is called the **difference** of A and B .
- The sum of matrices with different sizes is not defined.

Matrix Addition and Scalar Multiplication

Example

Let $A = \begin{pmatrix} 1 & -2 & 3 \\ 0 & 4 & 5 \end{pmatrix}$ and $B = \begin{pmatrix} 4 & 6 & 8 \\ 1 & -3 & -7 \end{pmatrix}$.

Matrix Addition and Scalar Multiplication

Example

Let $A = \begin{pmatrix} 1 & -2 & 3 \\ 0 & 4 & 5 \end{pmatrix}$ and $B = \begin{pmatrix} 4 & 6 & 8 \\ 1 & -3 & -7 \end{pmatrix}$. Then

$$A + B$$

Matrix Addition and Scalar Multiplication

Example

Let $A = \begin{pmatrix} 1 & -2 & 3 \\ 0 & 4 & 5 \end{pmatrix}$ and $B = \begin{pmatrix} 4 & 6 & 8 \\ 1 & -3 & -7 \end{pmatrix}$. Then

$$A + B = \begin{pmatrix} 1+4 & -2+6 & 3+8 \\ 0+1 & 4+(-3) & 5+(-7) \end{pmatrix}$$

Matrix Addition and Scalar Multiplication

Example

Let $A = \begin{pmatrix} 1 & -2 & 3 \\ 0 & 4 & 5 \end{pmatrix}$ and $B = \begin{pmatrix} 4 & 6 & 8 \\ 1 & -3 & -7 \end{pmatrix}$. Then

$$A + B = \begin{pmatrix} 1+4 & -2+6 & 3+8 \\ 0+1 & 4+(-3) & 5+(-7) \end{pmatrix} = \begin{pmatrix} 5 & 4 & 11 \\ 1 & 1 & -2 \end{pmatrix}$$

Matrix Addition and Scalar Multiplication

Example

Let $A = \begin{pmatrix} 1 & -2 & 3 \\ 0 & 4 & 5 \end{pmatrix}$ and $B = \begin{pmatrix} 4 & 6 & 8 \\ 1 & -3 & -7 \end{pmatrix}$. Then

$$A + B = \begin{pmatrix} 1+4 & -2+6 & 3+8 \\ 0+1 & 4+(-3) & 5+(-7) \end{pmatrix} = \begin{pmatrix} 5 & 4 & 11 \\ 1 & 1 & -2 \end{pmatrix}$$

$$3A$$

Matrix Addition and Scalar Multiplication

Example

Let $A = \begin{pmatrix} 1 & -2 & 3 \\ 0 & 4 & 5 \end{pmatrix}$ and $B = \begin{pmatrix} 4 & 6 & 8 \\ 1 & -3 & -7 \end{pmatrix}$. Then

$$A + B = \begin{pmatrix} 1+4 & -2+6 & 3+8 \\ 0+1 & 4+(-3) & 5+(-7) \end{pmatrix} = \begin{pmatrix} 5 & 4 & 11 \\ 1 & 1 & -2 \end{pmatrix}$$

$$3A = \begin{pmatrix} 3(1) & 3(-2) & 3(3) \\ 3(0) & 3(4) & 3(5) \end{pmatrix}$$

Matrix Addition and Scalar Multiplication

Example

Let $A = \begin{pmatrix} 1 & -2 & 3 \\ 0 & 4 & 5 \end{pmatrix}$ and $B = \begin{pmatrix} 4 & 6 & 8 \\ 1 & -3 & -7 \end{pmatrix}$. Then

$$A + B = \begin{pmatrix} 1+4 & -2+6 & 3+8 \\ 0+1 & 4+(-3) & 5+(-7) \end{pmatrix} = \begin{pmatrix} 5 & 4 & 11 \\ 1 & 1 & -2 \end{pmatrix}$$

$$3A = \begin{pmatrix} 3(1) & 3(-2) & 3(3) \\ 3(0) & 3(4) & 3(5) \end{pmatrix} = \begin{pmatrix} 3 & -6 & 9 \\ 0 & 12 & 15 \end{pmatrix}$$

Matrix Addition and Scalar Multiplication

Example

Let $A = \begin{pmatrix} 1 & -2 & 3 \\ 0 & 4 & 5 \end{pmatrix}$ and $B = \begin{pmatrix} 4 & 6 & 8 \\ 1 & -3 & -7 \end{pmatrix}$. Then

$$A + B = \begin{pmatrix} 1+4 & -2+6 & 3+8 \\ 0+1 & 4+(-3) & 5+(-7) \end{pmatrix} = \begin{pmatrix} 5 & 4 & 11 \\ 1 & 1 & -2 \end{pmatrix}$$

$$3A = \begin{pmatrix} 3(1) & 3(-2) & 3(3) \\ 3(0) & 3(4) & 3(5) \end{pmatrix} = \begin{pmatrix} 3 & -6 & 9 \\ 0 & 12 & 15 \end{pmatrix}$$

$$2A - 3B$$

Matrix Addition and Scalar Multiplication

Example

Let $A = \begin{pmatrix} 1 & -2 & 3 \\ 0 & 4 & 5 \end{pmatrix}$ and $B = \begin{pmatrix} 4 & 6 & 8 \\ 1 & -3 & -7 \end{pmatrix}$. Then

$$A + B = \begin{pmatrix} 1+4 & -2+6 & 3+8 \\ 0+1 & 4+(-3) & 5+(-7) \end{pmatrix} = \begin{pmatrix} 5 & 4 & 11 \\ 1 & 1 & -2 \end{pmatrix}$$

$$3A = \begin{pmatrix} 3(1) & 3(-2) & 3(3) \\ 3(0) & 3(4) & 3(5) \end{pmatrix} = \begin{pmatrix} 3 & -6 & 9 \\ 0 & 12 & 15 \end{pmatrix}$$

$$2A - 3B = \begin{pmatrix} 2 & -4 & 6 \\ 0 & 8 & 10 \end{pmatrix} + \begin{pmatrix} -12 & -18 & -24 \\ -3 & 9 & 21 \end{pmatrix}$$

Matrix Addition and Scalar Multiplication

Example

Let $A = \begin{pmatrix} 1 & -2 & 3 \\ 0 & 4 & 5 \end{pmatrix}$ and $B = \begin{pmatrix} 4 & 6 & 8 \\ 1 & -3 & -7 \end{pmatrix}$. Then

$$A + B = \begin{pmatrix} 1+4 & -2+6 & 3+8 \\ 0+1 & 4+(-3) & 5+(-7) \end{pmatrix} = \begin{pmatrix} 5 & 4 & 11 \\ 1 & 1 & -2 \end{pmatrix}$$

$$3A = \begin{pmatrix} 3(1) & 3(-2) & 3(3) \\ 3(0) & 3(4) & 3(5) \end{pmatrix} = \begin{pmatrix} 3 & -6 & 9 \\ 0 & 12 & 15 \end{pmatrix}$$

$$\begin{aligned} 2A - 3B &= \begin{pmatrix} 2 & -4 & 6 \\ 0 & 8 & 10 \end{pmatrix} + \begin{pmatrix} -12 & -18 & -24 \\ -3 & 9 & 21 \end{pmatrix} \\ &= \begin{pmatrix} -10 & -22 & -18 \\ -3 & 17 & 31 \end{pmatrix} \end{aligned}$$

Matrix Addition and Scalar Multiplication

Basic properties of matrices under the operations of matrix addition and scalar multiplication follow.

Matrix Addition and Scalar Multiplication

Basic properties of matrices under the operations of matrix addition and scalar multiplication follow.

Theorem

Consider any matrices A, B, C (with the same size) and any scalars k and k' . Then

Matrix Addition and Scalar Multiplication

Basic properties of matrices under the operations of matrix addition and scalar multiplication follow.

Theorem

Consider any matrices A, B, C (with the same size) and any scalars k and k' . Then

- $(A + B) + C = A + (B + C)$

Matrix Addition and Scalar Multiplication

Basic properties of matrices under the operations of matrix addition and scalar multiplication follow.

Theorem

Consider any matrices A, B, C (with the same size) and any scalars k and k' . Then

- $(A + B) + C = A + (B + C)$
- $A + 0 = 0 + A = A$

Matrix Addition and Scalar Multiplication

Basic properties of matrices under the operations of matrix addition and scalar multiplication follow.

Theorem

Consider any matrices A, B, C (with the same size) and any scalars k and k' . Then

- $(A + B) + C = A + (B + C)$
- $A + 0 = 0 + A = A$
- $A + (-A) = (-A) + A = 0$

Matrix Addition and Scalar Multiplication

Basic properties of matrices under the operations of matrix addition and scalar multiplication follow.

Theorem

Consider any matrices A, B, C (with the same size) and any scalars k and k' . Then

- $(A + B) + C = A + (B + C)$
- $A + 0 = 0 + A = A$
- $A + (-A) = (-A) + A = 0$
- $A + B = B + A$

Matrix Addition and Scalar Multiplication

Basic properties of matrices under the operations of matrix addition and scalar multiplication follow.

Theorem

Consider any matrices A, B, C (with the same size) and any scalars k and k' . Then

- $(A + B) + C = A + (B + C)$
- $A + 0 = 0 + A = A$
- $A + (-A) = (-A) + A = 0$
- $A + B = B + A$
- $k(A + B) = kA + kB$

Matrix Addition and Scalar Multiplication

Basic properties of matrices under the operations of matrix addition and scalar multiplication follow.

Theorem

Consider any matrices A, B, C (with the same size) and any scalars k and k' . Then

- $(A + B) + C = A + (B + C)$
- $A + 0 = 0 + A = A$
- $A + (-A) = (-A) + A = 0$
- $A + B = B + A$
- $k(A + B) = kA + kB$
- $(k + k')A = kA + k'A$

Matrix Addition and Scalar Multiplication

Basic properties of matrices under the operations of matrix addition and scalar multiplication follow.

Theorem

Consider any matrices A, B, C (with the same size) and any scalars k and k' . Then

- $(A + B) + C = A + (B + C)$
- $A + 0 = 0 + A = A$
- $A + (-A) = (-A) + A = 0$
- $A + B = B + A$
- $k(A + B) = kA + kB$
- $(k + k')A = kA + k'A$
- $(kk')A = k(k'A)$

Matrix Addition and Scalar Multiplication

Basic properties of matrices under the operations of matrix addition and scalar multiplication follow.

Theorem

Consider any matrices A, B, C (with the same size) and any scalars k and k' . Then

- $(A + B) + C = A + (B + C)$
- $A + 0 = 0 + A = A$
- $A + (-A) = (-A) + A = 0$
- $A + B = B + A$
- $k(A + B) = kA + kB$
- $(k + k')A = kA + k'A$
- $(kk')A = k(k'A)$
- $1 \cdot A = A$

Summation Symbol

Summation Symbol

Before we define matrix multiplication, it will be instructive to first introduce the summation symbol \sum (the Greek capital letter sigma).

Summation Symbol

Before we define matrix multiplication, it will be instructive to first introduce the summation symbol \sum (the Greek capital letter sigma).

- Suppose $f(k)$ is an algebraic expression involving the letter k .

Then

$$\sum_{k=1}^n f(k) = f(1) + f(2) + \dots + f(n).$$

Summation Symbol

Before we define matrix multiplication, it will be instructive to first introduce the summation symbol \sum (the Greek capital letter sigma).

- Suppose $f(k)$ is an algebraic expression involving the letter k .

Then

$$\sum_{k=1}^n f(k) = f(1) + f(2) + \dots + f(n).$$

- The letter k is called the index, and 1 and n are called, respectively, the lower and upper limits.

Summation Symbol

Before we define matrix multiplication, it will be instructive to first introduce the summation symbol \sum (the Greek capital letter sigma).

- Suppose $f(k)$ is an algebraic expression involving the letter k .

Then

$$\sum_{k=1}^n f(k) = f(1) + f(2) + \dots + f(n).$$

- The letter k is called the index, and 1 and n are called, respectively, the lower and upper limits.
- We also generalize our definition by allowing the sum to range from any integer n_1 to any integer n_2 . That is, we define

$$\sum_{k=n_1}^{n_2} f(k) = f(n_1) + f(n_1 + 1) + f(n_1 + 2) + \dots + f(n_2)$$

Example

$$1. \sum_{k=1}^5 x_k = x_1 + x_2 + x_3 + x_4 + x_5$$

Summation Symbol

Example

$$1. \sum_{k=1}^5 x_k = x_1 + x_2 + x_3 + x_4 + x_5$$

$$2. \sum_{i=1}^n a_i b_i = a_1 b_1 + a_2 b_2 + \cdots + a_n b_n$$

Summation Symbol

Example

$$1. \sum_{k=1}^5 x_k = x_1 + x_2 + x_3 + x_4 + x_5$$

$$2. \sum_{i=1}^n a_i b_i = a_1 b_1 + a_2 b_2 + \cdots + a_n b_n$$

$$3. \sum_{j=2}^5 j^2 = 2^2 + 3^2 + 4^2 + 5^2 = 54$$

Summation Symbol

Example

$$1. \sum_{k=1}^5 x_k = x_1 + x_2 + x_3 + x_4 + x_5$$

$$2. \sum_{i=1}^n a_i b_i = a_1 b_1 + a_2 b_2 + \cdots + a_n b_n$$

$$3. \sum_{j=2}^5 j^2 = 2^2 + 3^2 + 4^2 + 5^2 = 54$$

$$4. \sum_{i=0}^n a_i x^i = a_0 + a_1 x + a_2 x^2 + \cdots + a_n x^n$$

Summation Symbol

Example

$$1. \sum_{k=1}^5 x_k = x_1 + x_2 + x_3 + x_4 + x_5$$

$$2. \sum_{i=1}^n a_i b_i = a_1 b_1 + a_2 b_2 + \cdots + a_n b_n$$

$$3. \sum_{j=2}^5 j^2 = 2^2 + 3^2 + 4^2 + 5^2 = 54$$

$$4. \sum_{i=0}^n a_i x^i = a_0 + a_1 x + a_2 x^2 + \cdots + a_n x^n$$

$$5. \sum_{k=1}^p a_{ik} b_{kj} = a_{i1} b_{1j} + a_{i2} b_{2j} + a_{i3} b_{3j} + \cdots + a_{ip} b_{pj}$$

Matrix Multiplication

Matrix Multiplication

- The product of matrices A and B , written AB , is somewhat complicated. For this reason, we first begin with a special case.

Matrix Multiplication

- The product of matrices A and B , written AB , is somewhat complicated. For this reason, we first begin with a special case.
- The product AB of a row matrix $A = [a_i]$ and a column matrix $B = [b_i]$ with the same number of elements is defined to be the scalar (or 1×1 matrix) obtained by multiplying corresponding entries and adding; that is,

$$AB = \begin{pmatrix} a_1 & a_2 & \dots & a_n \end{pmatrix} \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix}$$

Matrix Multiplication

- The product of matrices A and B , written AB , is somewhat complicated. For this reason, we first begin with a special case.
- The product AB of a row matrix $A = [a_i]$ and a column matrix $B = [b_i]$ with the same number of elements is defined to be the scalar (or 1×1 matrix) obtained by multiplying corresponding entries and adding; that is,

$$AB = \begin{pmatrix} a_1 & a_2 & \dots & a_n \end{pmatrix} \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix}$$

Matrix Multiplication

- The product of matrices A and B , written AB , is somewhat complicated. For this reason, we first begin with a special case.
- The product AB of a row matrix $A = [a_i]$ and a column matrix $B = [b_i]$ with the same number of elements is defined to be the scalar (or 1×1 matrix) obtained by multiplying corresponding entries and adding; that is,

$$AB = \begin{pmatrix} a_1 & a_2 & \dots & a_n \end{pmatrix} \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix} = a_1b_1 + a_2b_2 + \dots + a_nb_n$$

Matrix Multiplication

- The product of matrices A and B , written AB , is somewhat complicated. For this reason, we first begin with a special case.
- The product AB of a row matrix $A = [a_i]$ and a column matrix $B = [b_i]$ with the same number of elements is defined to be the scalar (or 1×1 matrix) obtained by multiplying corresponding entries and adding; that is,

$$AB = \begin{pmatrix} a_1 & a_2 & \dots & a_n \end{pmatrix} \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix} = a_1b_1 + a_2b_2 + \dots + a_nb_n = \sum_{k=1}^n a_k b_k$$

Matrix Multiplication

- The product of matrices A and B , written AB , is somewhat complicated. For this reason, we first begin with a special case.
- The product AB of a row matrix $A = [a_i]$ and a column matrix $B = [b_i]$ with the same number of elements is defined to be the scalar (or 1×1 matrix) obtained by multiplying corresponding entries and adding; that is,

$$AB = \begin{pmatrix} a_1 & a_2 & \dots & a_n \end{pmatrix} \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix} = a_1b_1 + a_2b_2 + \dots + a_nb_n = \sum_{k=1}^n a_k b_k$$

- The product AB is not defined when A and B have different numbers of elements.

Matrix Multiplication

Example

$$1. \begin{pmatrix} 7 & -4 & 5 \end{pmatrix} \begin{pmatrix} 3 \\ 2 \\ -1 \end{pmatrix} = 7(3) + (-4)(2) + 5(-1) =$$
$$21 - 8 - 5 = 8.$$

Matrix Multiplication

Example

$$1. \begin{pmatrix} 7 & -4 & 5 \end{pmatrix} \begin{pmatrix} 3 \\ 2 \\ -1 \end{pmatrix} = 7(3) + (-4)(2) + 5(-1) = 21 - 8 - 5 = 8.$$

$$2. \begin{pmatrix} 6 & -1 & 8 & 3 \end{pmatrix} \begin{pmatrix} 4 \\ -9 \\ -2 \\ 5 \end{pmatrix} = 24 + 9 - 16 + 15 = 32.$$

Matrix Multiplication

Example

$$1. \begin{pmatrix} 7 & -4 & 5 \end{pmatrix} \begin{pmatrix} 3 \\ 2 \\ -1 \end{pmatrix} = 7(3) + (-4)(2) + 5(-1) = 21 - 8 - 5 = 8.$$

$$2. \begin{pmatrix} 6 & -1 & 8 & 3 \end{pmatrix} \begin{pmatrix} 4 \\ -9 \\ -2 \\ 5 \end{pmatrix} = 24 + 9 - 16 + 15 = 32.$$

We are now ready to define matrix multiplication in general.

Matrix Multiplication

Suppose $A = [a_{ik}]$ and $B = [b_{kj}]$ are matrices such that the number of columns of A is equal to the number of rows of B ; say, A is an $m \times p$ matrix and B is a $p \times n$ matrix.

Matrix Multiplication

Suppose $A = [a_{ik}]$ and $B = [b_{kj}]$ are matrices such that the number of columns of A is equal to the number of rows of B ; say, A is an $m \times p$ matrix and B is a $p \times n$ matrix. Then the product AB is the $m \times n$ matrix whose ij -entry is obtained by multiplying the i^{th} row of A by the j^{th} column of B .

Matrix Multiplication

Suppose $A = [a_{ik}]$ and $B = [b_{kj}]$ are matrices such that the number of columns of A is equal to the number of rows of B ; say, A is an $m \times p$ matrix and B is a $p \times n$ matrix. Then the product AB is the $m \times n$ matrix whose ij -entry is obtained by multiplying the i^{th} row of A by the j^{th} column of B . That is,

$$\begin{pmatrix} a_{11} & \dots & a_{1p} \\ \vdots & \dots & \vdots \\ a_{i1} & \dots & a_{ip} \\ \vdots & \dots & \vdots \\ a_{m1} & \dots & a_{mp} \end{pmatrix} \begin{pmatrix} b_{11} & \dots & b_{1j} & \dots & b_{1n} \\ \vdots & \dots & \vdots & \dots & \vdots \\ \vdots & \dots & \vdots & \dots & \vdots \\ \vdots & \dots & \vdots & \dots & \vdots \\ b_{p1} & \dots & b_{pj} & \dots & b_{pn} \end{pmatrix} = \begin{pmatrix} c_{11} & \dots & c_{1n} \\ \vdots & \dots & \vdots \\ \vdots & c_{ij} & \vdots \\ \vdots & \dots & \vdots \\ c_{m1} & \dots & c_{mn} \end{pmatrix}$$

Matrix Multiplication

Suppose $A = [a_{ik}]$ and $B = [b_{kj}]$ are matrices such that the number of columns of A is equal to the number of rows of B ; say, A is an $m \times p$ matrix and B is a $p \times n$ matrix. Then the product AB is the $m \times n$ matrix whose ij -entry is obtained by multiplying the i^{th} row of A by the j^{th} column of B . That is,

$$\begin{pmatrix} a_{11} & \cdots & a_{1p} \\ \vdots & \ddots & \vdots \\ a_{i1} & \cdots & a_{ip} \\ \vdots & \ddots & \vdots \\ a_{m1} & \cdots & a_{mp} \end{pmatrix} \begin{pmatrix} b_{11} & \cdots & b_{1j} & \cdots & b_{1n} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ b_{p1} & \cdots & b_{pj} & \cdots & b_{pn} \end{pmatrix} = \begin{pmatrix} c_{11} & \cdots & c_{1n} \\ \vdots & \ddots & \vdots \\ \vdots & c_{ij} & \vdots \\ \vdots & \ddots & \vdots \\ c_{m1} & \cdots & c_{mn} \end{pmatrix}$$

$$c_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{ip}b_{pj} = \sum_{k=1}^p a_{ik}b_{kj}$$

Matrix Multiplication

Suppose $A = [a_{ik}]$ and $B = [b_{kj}]$ are matrices such that the number of columns of A is equal to the number of rows of B ; say, A is an $m \times p$ matrix and B is a $p \times n$ matrix. Then the product AB is the $m \times n$ matrix whose ij -entry is obtained by multiplying the i^{th} row of A by the j^{th} column of B . That is,

$$\begin{pmatrix} a_{11} & \cdots & a_{1p} \\ \vdots & \ddots & \vdots \\ a_{i1} & \cdots & a_{ip} \\ \vdots & \ddots & \vdots \\ a_{m1} & \cdots & a_{mp} \end{pmatrix} \begin{pmatrix} b_{11} & \cdots & b_{1j} & \cdots & b_{1n} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ b_{p1} & \cdots & b_{pj} & \cdots & b_{pn} \end{pmatrix} = \begin{pmatrix} c_{11} & \cdots & c_{1n} \\ \vdots & \ddots & \vdots \\ \vdots & c_{ij} & \vdots \\ \vdots & \ddots & \vdots \\ c_{m1} & \cdots & c_{mn} \end{pmatrix}$$

$$c_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{ip}b_{pj} = \sum_{k=1}^p a_{ik}b_{kj}$$

- The product AB is not defined if A is an $m \times p$ matrix and B is a $q \times n$ matrix, where $p \neq q$.

Example

Find AB where $A = \begin{pmatrix} 1 & 3 \\ 2 & -1 \end{pmatrix}$ and $B = \begin{pmatrix} 2 & 0 & -4 \\ 5 & -2 & 6 \end{pmatrix}$.

Example

Find AB where $A = \begin{pmatrix} 1 & 3 \\ 2 & -1 \end{pmatrix}$ and $B = \begin{pmatrix} 2 & 0 & -4 \\ 5 & -2 & 6 \end{pmatrix}$.

Because A is 2×2 and B is 2×3 , the product AB is defined and AB is a 2×3 matrix.

Example

Find AB where $A = \begin{pmatrix} 1 & 3 \\ 2 & -1 \end{pmatrix}$ and $B = \begin{pmatrix} 2 & 0 & -4 \\ 5 & -2 & 6 \end{pmatrix}$.

Because A is 2×2 and B is 2×3 , the product AB is defined and AB is a 2×3 matrix. To obtain the first row of the product matrix AB , multiply the first row $\begin{pmatrix} 1 & 3 \end{pmatrix}$ of A by each column of B , $\begin{pmatrix} 2 \\ 5 \end{pmatrix}$, $\begin{pmatrix} 0 \\ -2 \end{pmatrix}$, $\begin{pmatrix} -4 \\ 6 \end{pmatrix}$ respectively.

Example

Find AB where $A = \begin{pmatrix} 1 & 3 \\ 2 & -1 \end{pmatrix}$ and $B = \begin{pmatrix} 2 & 0 & -4 \\ 5 & -2 & 6 \end{pmatrix}$.

Because A is 2×2 and B is 2×3 , the product AB is defined and AB is a 2×3 matrix. To obtain the first row of the product matrix AB , multiply the first row $\begin{pmatrix} 1 & 3 \end{pmatrix}$ of A by each column of B ,

$\begin{pmatrix} 2 \\ 5 \end{pmatrix}$, $\begin{pmatrix} 0 \\ -2 \end{pmatrix}$, $\begin{pmatrix} -4 \\ 6 \end{pmatrix}$ respectively. That is,

$$AB = \begin{pmatrix} 2 + 15 & 0 - 6 & -4 + 18 \\ . & . & . \end{pmatrix} = \begin{pmatrix} 17 & -6 & 14 \\ . & . & . \end{pmatrix}$$

Example

Find AB where $A = \begin{pmatrix} 1 & 3 \\ 2 & -1 \end{pmatrix}$ and $B = \begin{pmatrix} 2 & 0 & -4 \\ 5 & -2 & 6 \end{pmatrix}$.

Because A is 2×2 and B is 2×3 , the product AB is defined and AB is a 2×3 matrix. To obtain the first row of the product matrix AB , multiply the first row $\begin{pmatrix} 1 & 3 \end{pmatrix}$ of A by each column of B ,

$\begin{pmatrix} 2 \\ 5 \end{pmatrix}$, $\begin{pmatrix} 0 \\ -2 \end{pmatrix}$, $\begin{pmatrix} -4 \\ 6 \end{pmatrix}$ respectively. That is,

$$AB = \begin{pmatrix} 2 + 15 & 0 - 6 & -4 + 18 \\ . & . & . \end{pmatrix} = \begin{pmatrix} 17 & -6 & 14 \\ . & . & . \end{pmatrix}$$

To obtain the second row of AB , multiply the second row $\begin{pmatrix} 2 & -1 \end{pmatrix}$ of A by each column of B .

Example

Find AB where $A = \begin{pmatrix} 1 & 3 \\ 2 & -1 \end{pmatrix}$ and $B = \begin{pmatrix} 2 & 0 & -4 \\ 5 & -2 & 6 \end{pmatrix}$.

Because A is 2×2 and B is 2×3 , the product AB is defined and AB is a 2×3 matrix. To obtain the first row of the product matrix AB , multiply the first row $\begin{pmatrix} 1 & 3 \end{pmatrix}$ of A by each column of B ,

$\begin{pmatrix} 2 \\ 5 \end{pmatrix}$, $\begin{pmatrix} 0 \\ -2 \end{pmatrix}$, $\begin{pmatrix} -4 \\ 6 \end{pmatrix}$ respectively. That is,

$$AB = \begin{pmatrix} 2 + 15 & 0 - 6 & -4 + 18 \\ . & . & . \end{pmatrix} = \begin{pmatrix} 17 & -6 & 14 \\ . & . & . \end{pmatrix}$$

To obtain the second row of AB , multiply the second row $\begin{pmatrix} 2 & -1 \end{pmatrix}$ of A by each column of B . Thus,

$$AB = \begin{pmatrix} 17 & -6 & 14 \\ 4 - 5 & 0 + 2 & -8 - 6 \end{pmatrix} = \begin{pmatrix} 17 & -6 & 14 \\ -1 & 2 & -14 \end{pmatrix}$$

Matrix Multiplication

Example

Suppose $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$ and $B = \begin{pmatrix} 5 & 6 \\ 0 & -2 \end{pmatrix}$.

Matrix Multiplication

Example

Suppose $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$ and $B = \begin{pmatrix} 5 & 6 \\ 0 & -2 \end{pmatrix}$. Then

$$AB = \begin{pmatrix} 5+0 & 6-4 \\ 15+0 & 18-8 \end{pmatrix} = \begin{pmatrix} 5 & 2 \\ 15 & 10 \end{pmatrix}$$

Matrix Multiplication

Example

Suppose $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$ and $B = \begin{pmatrix} 5 & 6 \\ 0 & -2 \end{pmatrix}$. Then

$$AB = \begin{pmatrix} 5+0 & 6-4 \\ 15+0 & 18-8 \end{pmatrix} = \begin{pmatrix} 5 & 2 \\ 15 & 10 \end{pmatrix}$$

and

$$BA = \begin{pmatrix} 5+18 & 10+24 \\ 0-6 & 0-8 \end{pmatrix} = \begin{pmatrix} 23 & 34 \\ -6 & -8 \end{pmatrix}$$

Matrix Multiplication

Example

Suppose $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$ and $B = \begin{pmatrix} 5 & 6 \\ 0 & -2 \end{pmatrix}$. Then

$$AB = \begin{pmatrix} 5+0 & 6-4 \\ 15+0 & 18-8 \end{pmatrix} = \begin{pmatrix} 5 & 2 \\ 15 & 10 \end{pmatrix}$$

and

$$BA = \begin{pmatrix} 5+18 & 10+24 \\ 0-6 & 0-8 \end{pmatrix} = \begin{pmatrix} 23 & 34 \\ -6 & -8 \end{pmatrix}$$

Note: The above example shows that matrix multiplication is not commutative that is, in general, $AB \neq BA$. However, matrix multiplication does satisfy the following properties.

Theorem

Let A, B, C be matrices. Then, whenever the products and sums are defined,

Theorem

Let A, B, C be matrices. Then, whenever the products and sums are defined,

1. $(AB)C = A(BC)$ (*associative law*),

Theorem

Let A, B, C be matrices. Then, whenever the products and sums are defined,

1. $(AB)C = A(BC)$ (*associative law*),
2. $A(B + C) = AB + AC$ (*left distributive law*),

Theorem

Let A, B, C be matrices. Then, whenever the products and sums are defined,

1. $(AB)C = A(BC)$ (*associative law*),
2. $A(B + C) = AB + AC$ (*left distributive law*),
3. $(B + C)A = BA + CA$ (*right distributive law*),

Theorem

Let A, B, C be matrices. Then, whenever the products and sums are defined,

1. $(AB)C = A(BC)$ (*associative law*),
2. $A(B + C) = AB + AC$ (*left distributive law*),
3. $(B + C)A = BA + CA$ (*right distributive law*),
4. $k(AB) = (kA)B = A(kB)$, *where k is a scalar.*

Theorem

Let A, B, C be matrices. Then, whenever the products and sums are defined,

1. $(AB)C = A(BC)$ (*associative law*),
2. $A(B + C) = AB + AC$ (*left distributive law*),
3. $(B + C)A = BA + CA$ (*right distributive law*),
4. $k(AB) = (kA)B = A(kB)$, *where k is a scalar.*
5. *We note that $0A = 0$ and $B0 = 0$, where 0 is the zero matrix.*

Transpose of a Matrix

Transpose of a Matrix

The transpose of a matrix A , written A^T , is the matrix obtained by writing the columns of A , in order, as rows.

Transpose of a Matrix

The transpose of a matrix A , written A^T , is the matrix obtained by writing the columns of A , in order, as rows. For example,

$$\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}^T = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix}$$

Transpose of a Matrix

The transpose of a matrix A , written A^T , is the matrix obtained by writing the columns of A , in order, as rows. For example,

$$\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}^T = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} \quad \text{and} \quad (1 \quad -3 \quad -5)^T = \begin{pmatrix} 1 \\ -3 \\ -5 \end{pmatrix}$$

Transpose of a Matrix

The transpose of a matrix A , written A^T , is the matrix obtained by writing the columns of A , in order, as rows. For example,

$$\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}^T = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} \quad \text{and} \quad (1 \quad -3 \quad -5)^T = \begin{pmatrix} 1 \\ -3 \\ -5 \end{pmatrix}$$

- In other words, if $A = [a_{ij}]$ is an $m \times n$ matrix, then $A^T = [b_{ij}]$ is the $n \times m$ matrix where $b_{ij} = a_{ji}$.

Transpose of a Matrix

The transpose of a matrix A , written A^T , is the matrix obtained by writing the columns of A , in order, as rows. For example,

$$\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}^T = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} \quad \text{and} \quad (1 \quad -3 \quad -5)^T = \begin{pmatrix} 1 \\ -3 \\ -5 \end{pmatrix}$$

- In other words, if $A = [a_{ij}]$ is an $m \times n$ matrix, then $A^T = [b_{ij}]$ is the $n \times m$ matrix where $b_{ij} = a_{ji}$.
- Observe that the transpose of a row vector is a column vector. Similarly, the transpose of a column vector is a row vector.

Transpose of a Matrix

The transpose of a matrix A , written A^T , is the matrix obtained by writing the columns of A , in order, as rows. For example,

$$\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}^T = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} \quad \text{and} \quad (1 \quad -3 \quad -5)^T = \begin{pmatrix} 1 \\ -3 \\ -5 \end{pmatrix}$$

- In other words, if $A = [a_{ij}]$ is an $m \times n$ matrix, then $A^T = [b_{ij}]$ is the $n \times m$ matrix where $b_{ij} = a_{ji}$.
- Observe that the transpose of a row vector is a column vector. Similarly, the transpose of a column vector is a row vector.
- The next theorem lists basic properties of the transpose operation.

Transpose of a Matrix

Theorem

Let A and B be matrices and let k be a scalar. Then, whenever the sum and product are defined,

Transpose of a Matrix

Theorem

Let A and B be matrices and let k be a scalar. Then, whenever the sum and product are defined,

1. $(A + B)^T = A^T + B^T$

Transpose of a Matrix

Theorem

Let A and B be matrices and let k be a scalar. Then, whenever the sum and product are defined,

$$1. (A + B)^T = A^T + B^T$$

$$2. (kA)^T = kA^T$$

Transpose of a Matrix

Theorem

Let A and B be matrices and let k be a scalar. Then, whenever the sum and product are defined,

$$1. (A + B)^T = A^T + B^T$$

$$2. (kA)^T = kA^T$$

$$3. (A^T)^T = A$$

Transpose of a Matrix

Theorem

Let A and B be matrices and let k be a scalar. Then, whenever the sum and product are defined,

$$1. (A + B)^T = A^T + B^T$$

$$2. (kA)^T = kA^T$$

$$3. (A^T)^T = A$$

$$4. (AB)^T = B^T A^T$$

Transpose of a Matrix

Theorem

Let A and B be matrices and let k be a scalar. Then, whenever the sum and product are defined,

$$1. (A + B)^T = A^T + B^T$$

$$2. (kA)^T = kA^T$$

$$3. (A^T)^T = A$$

$$4. (AB)^T = B^T A^T$$

We emphasize that, by (4), the transpose of a product is the product of the transposes, but in the reverse order.

Square Matrices

Square Matrices

A square matrix is a matrix with the same number of rows as columns. An $n \times n$ square matrix is said to be of order n and is sometimes called an n -square matrix.

Square Matrices

A square matrix is a matrix with the same number of rows as columns. An $n \times n$ square matrix is said to be of order n and is sometimes called an n -square matrix.

- Recall that not every two matrices can be added or multiplied. However, if we only consider square matrices of some given order n , then this inconvenience disappears.

Square Matrices

A square matrix is a matrix with the same number of rows as columns. An $n \times n$ square matrix is said to be of order n and is sometimes called an n -square matrix.

- Recall that not every two matrices can be added or multiplied. However, if we only consider square matrices of some given order n , then this inconvenience disappears.
- Specifically, the operations of addition, multiplication, scalar multiplication, and transpose can be performed on any $n \times n$ matrices, and the result is again an $n \times n$ matrix.

Square Matrices

Example

The following are square matrices of order 3:

Square Matrices

Example

The following are square matrices of order 3:

$$A = \begin{pmatrix} 1 & 2 & 3 \\ -4 & -4 & -4 \\ 5 & 6 & 7 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 2 & -5 & 1 \\ 0 & 3 & -2 \\ 1 & 2 & -4 \end{pmatrix}$$

Square Matrices

Example

The following are square matrices of order 3:

$$A = \begin{pmatrix} 1 & 2 & 3 \\ -4 & -4 & -4 \\ 5 & 6 & 7 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 2 & -5 & 1 \\ 0 & 3 & -2 \\ 1 & 2 & -4 \end{pmatrix}$$

The following are also matrices of order 3:

Square Matrices

Example

The following are square matrices of order 3:

$$A = \begin{pmatrix} 1 & 2 & 3 \\ -4 & -4 & -4 \\ 5 & 6 & 7 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 2 & -5 & 1 \\ 0 & 3 & -2 \\ 1 & 2 & -4 \end{pmatrix}$$

The following are also matrices of order 3:

$$A + B = \begin{pmatrix} 3 & -3 & 4 \\ -4 & -1 & -6 \\ 6 & 8 & 3 \end{pmatrix},$$

Square Matrices

Example

The following are square matrices of order 3:

$$A = \begin{pmatrix} 1 & 2 & 3 \\ -4 & -4 & -4 \\ 5 & 6 & 7 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 2 & -5 & 1 \\ 0 & 3 & -2 \\ 1 & 2 & -4 \end{pmatrix}$$

The following are also matrices of order 3:

$$A + B = \begin{pmatrix} 3 & -3 & 4 \\ -4 & -1 & -6 \\ 6 & 8 & 3 \end{pmatrix}, \quad 2A = \begin{pmatrix} 2 & 4 & 6 \\ -8 & -8 & -8 \\ 10 & 12 & 14 \end{pmatrix},$$

Square Matrices

Example

The following are square matrices of order 3:

$$A = \begin{pmatrix} 1 & 2 & 3 \\ -4 & -4 & -4 \\ 5 & 6 & 7 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 2 & -5 & 1 \\ 0 & 3 & -2 \\ 1 & 2 & -4 \end{pmatrix}$$

The following are also matrices of order 3:

$$A + B = \begin{pmatrix} 3 & -3 & 4 \\ -4 & -1 & -6 \\ 6 & 8 & 3 \end{pmatrix}, \quad 2A = \begin{pmatrix} 2 & 4 & 6 \\ -8 & -8 & -8 \\ 10 & 12 & 14 \end{pmatrix}, \quad A^T = \begin{pmatrix} 1 & -4 & 5 \\ 2 & -4 & 6 \\ 3 & -4 & 7 \end{pmatrix}$$

Square Matrices

Example

The following are square matrices of order 3:

$$A = \begin{pmatrix} 1 & 2 & 3 \\ -4 & -4 & -4 \\ 5 & 6 & 7 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 2 & -5 & 1 \\ 0 & 3 & -2 \\ 1 & 2 & -4 \end{pmatrix}$$

The following are also matrices of order 3:

$$A + B = \begin{pmatrix} 3 & -3 & 4 \\ -4 & -1 & -6 \\ 6 & 8 & 3 \end{pmatrix}, \quad 2A = \begin{pmatrix} 2 & 4 & 6 \\ -8 & -8 & -8 \\ 10 & 12 & 14 \end{pmatrix}, \quad A^T = \begin{pmatrix} 1 & -4 & 5 \\ 2 & -4 & 6 \\ 3 & -4 & 7 \end{pmatrix}$$

$$AB = \begin{pmatrix} 5 & 7 & -15 \\ -12 & 0 & 20 \\ 17 & 7 & -35 \end{pmatrix},$$

Square Matrices

Example

The following are square matrices of order 3:

$$A = \begin{pmatrix} 1 & 2 & 3 \\ -4 & -4 & -4 \\ 5 & 6 & 7 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 2 & -5 & 1 \\ 0 & 3 & -2 \\ 1 & 2 & -4 \end{pmatrix}$$

The following are also matrices of order 3:

$$A + B = \begin{pmatrix} 3 & -3 & 4 \\ -4 & -1 & -6 \\ 6 & 8 & 3 \end{pmatrix}, \quad 2A = \begin{pmatrix} 2 & 4 & 6 \\ -8 & -8 & -8 \\ 10 & 12 & 14 \end{pmatrix}, \quad A^T = \begin{pmatrix} 1 & -4 & 5 \\ 2 & -4 & 6 \\ 3 & -4 & 7 \end{pmatrix}$$

$$AB = \begin{pmatrix} 5 & 7 & -15 \\ -12 & 0 & 20 \\ 17 & 7 & -35 \end{pmatrix}, \quad BA = \begin{pmatrix} 27 & 30 & 33 \\ -22 & -24 & -26 \\ -27 & -30 & -33 \end{pmatrix}$$

Diagonal and Trace

Diagonal and Trace

Let $A = [a_{ij}]$ be an n -square matrix. The diagonal or main diagonal of A consists of the elements with the same subscripts - that is,

$$a_{11}, \quad a_{22}, \quad a_{33}, \quad \dots, \quad a_{nn}$$

Diagonal and Trace

Let $A = [a_{ij}]$ be an n -square matrix. The diagonal or main diagonal of A consists of the elements with the same subscripts - that is,

$$a_{11}, \quad a_{22}, \quad a_{33}, \quad \dots, \quad a_{nn}$$

The trace of A , written $Tr(A)$, is the sum of the diagonal elements. Namely,

$$Tr(A) = a_{11} + a_{22} + a_{33} + \dots + a_{nn}$$

Diagonal and Trace

Let $A = [a_{ij}]$ be an n -square matrix. The diagonal or main diagonal of A consists of the elements with the same subscripts - that is,

$$a_{11}, \quad a_{22}, \quad a_{33}, \quad \dots, \quad a_{nn}$$

The trace of A , written $Tr(A)$, is the sum of the diagonal elements. Namely,

$$Tr(A) = a_{11} + a_{22} + a_{33} + \dots + a_{nn}$$

The following theorem applies.

Theorem

Suppose $A = [a_{ij}]$ and $B = [b_{ij}]$ are n -square matrices and k is a scalar. Then

Theorem

Suppose $A = [a_{ij}]$ and $B = [b_{ij}]$ are n -square matrices and k is a scalar. Then

1. $Tr(A + B) = Tr(A) + Tr(B)$

Theorem

Suppose $A = [a_{ij}]$ and $B = [b_{ij}]$ are n -square matrices and k is a scalar. Then

1. $Tr(A + B) = Tr(A) + Tr(B)$
2. $Tr(A^T) = Tr(A)$

Theorem

Suppose $A = [a_{ij}]$ and $B = [b_{ij}]$ are n -square matrices and k is a scalar. Then

1. $Tr(A + B) = Tr(A) + Tr(B)$
2. $Tr(A^T) = Tr(A)$
3. $Tr(kA) = kTr(A)$

Theorem

Suppose $A = [a_{ij}]$ and $B = [b_{ij}]$ are n -square matrices and k is a scalar. Then

1. $Tr(A + B) = Tr(A) + Tr(B)$
2. $Tr(A^T) = Tr(A)$
3. $Tr(kA) = kTr(A)$
4. $Tr(AB) = Tr(BA)$

Example

Let A and B be square matrices of order 3:

$$A = \begin{pmatrix} 1 & 2 & 3 \\ -4 & -4 & -4 \\ 5 & 6 & 7 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 2 & -5 & 1 \\ 0 & 3 & -2 \\ 1 & 2 & -4 \end{pmatrix}$$

Then diagonal of $A = \{1, -4, 7\}$ and $Tr(A) = 1 - 4 + 7 = 4$
diagonal of $B = \{2, 3, -4\}$ and $Tr(B) = 2 + 3 - 4 = 1$

$$Tr(A+B) = 3 - 1 + 3 = 5, \quad Tr(2A) = 2 - 8 + 14 = 8, \quad Tr(A^T) = 4$$
$$Tr(AB) = 5 + 0 - 35 = -30, \quad Tr(BA) = 27 - 24 - 33 = -30$$

As expected from previous Theorem,

$$Tr(A+B) = Tr(A) + Tr(B), \quad Tr(A^T) = Tr(A), \quad Tr(2A) = 2Tr(A)$$

Furthermore, although $AB \neq BA$, the traces are equal.

Example

Let A and B be square matrices of order 3:

$$A = \begin{pmatrix} 1 & 2 & 3 \\ -4 & -4 & -4 \\ 5 & 6 & 7 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 2 & -5 & 1 \\ 0 & 3 & -2 \\ 1 & 2 & -4 \end{pmatrix}$$

Then diagonal of $A = \{1, -4, 7\}$ and $Tr(A) = 1 - 4 + 7 = 4$
diagonal of $B = \{2, 3, -4\}$ and $Tr(B) = 2 + 3 - 4 = 1$

$$Tr(A+B) = 3 - 1 + 3 = 5, \quad Tr(2A) = 2 - 8 + 14 = 8, \quad Tr(A^T) = 4$$
$$Tr(AB) = 5 + 0 - 35 = -30, \quad Tr(BA) = 27 - 24 - 33 = -30$$

As expected from previous Theorem,

$$Tr(A+B) = Tr(A) + Tr(B), \quad Tr(A^T) = Tr(A), \quad Tr(2A) = 2Tr(A)$$

Furthermore, although $AB \neq BA$, the traces are equal.

Identity Matrix, Scalar Matrices

Identity Matrix, Scalar Matrices

The n -square identity or unit matrix, denoted by I_n , or simply I , is the n -square matrix with 1's on the diagonal and 0's elsewhere.

The identity matrix I is similar to the scalar 1 in that, for any n -square matrix A

$$AI = IA = A$$

Identity Matrix, Scalar Matrices

The n -square identity or unit matrix, denoted by I_n , or simply I , is the n -square matrix with 1's on the diagonal and 0's elsewhere.

The identity matrix I is similar to the scalar 1 in that, for any n -square matrix A

$$AI = IA = A$$

More generally, if B is an $m \times n$ matrix, then $BI_n = I_m B = B$.

Identity Matrix, Scalar Matrices

The n -square identity or unit matrix, denoted by I_n , or simply I , is the n -square matrix with 1's on the diagonal and 0's elsewhere. The identity matrix I is similar to the scalar 1 in that, for any n -square matrix A

$$AI = IA = A$$

More generally, if B is an $m \times n$ matrix, then $BI_n = I_m B = B$.

For any scalar k , the matrix kI that contains k 's on the diagonal and 0's elsewhere is called the scalar matrix corresponding to the scalar k .

Identity Matrix, Scalar Matrices

The n -square identity or unit matrix, denoted by I_n , or simply I , is the n -square matrix with 1's on the diagonal and 0's elsewhere. The identity matrix I is similar to the scalar 1 in that, for any n -square matrix A

$$AI = IA = A$$

More generally, if B is an $m \times n$ matrix, then $BI_n = I_m B = B$.

For any scalar k , the matrix kI that contains k 's on the diagonal and 0's elsewhere is called the scalar matrix corresponding to the scalar k . Observe that

$$(kI)A = k(IA) = kA$$

That is, multiplying a matrix A by the scalar matrix kI is equivalent to multiplying A by the scalar k

Identity Matrix, Scalar Matrices

Example

The following are the identity matrices of orders 3 and 4 and the corresponding scalar matrices for $k = 5$:

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & 1 \end{pmatrix}, \quad \begin{pmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{pmatrix}, \quad \begin{pmatrix} 5 & & & \\ & 5 & & \\ & & 5 & \\ & & & 5 \end{pmatrix}$$

Powers of Matrices, Polynomials in Matrices

Powers of Matrices, Polynomials in Matrices

Let A be an n -square matrix over a field K . Powers of A are defined as follows:

$$A^2 = AA, \quad A^3 = A^2A, \quad \dots, \quad A^{n+1} = A^nA, \quad \dots, \quad \text{and} \quad A^0 = I$$

Powers of Matrices, Polynomials in Matrices

Let A be an n -square matrix over a field K . Powers of A are defined as follows:

$$A^2 = AA, \quad A^3 = A^2A, \quad \dots, \quad A^{n+1} = A^nA, \quad \dots, \quad \text{and} \quad A^0 = I$$

Polynomials in the matrix A are also defined. Specifically, for any polynomial

$$f(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$$

where the a_i are scalars in K ,

Powers of Matrices, Polynomials in Matrices

Let A be an n -square matrix over a field K . Powers of A are defined as follows:

$$A^2 = AA, \quad A^3 = A^2A, \quad \dots, \quad A^{n+1} = A^nA, \quad \dots, \quad \text{and} \quad A^0 = I$$

Polynomials in the matrix A are also defined. Specifically, for any polynomial

$$f(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$$

where the a_i are scalars in K , $f(A)$ is defined to be the following matrix:

$$f(A) = a_0I + a_1A + a_2A^2 + \dots + a_nA^n$$

Powers of Matrices, Polynomials in Matrices

Let A be an n -square matrix over a field K . Powers of A are defined as follows:

$$A^2 = AA, \quad A^3 = A^2A, \quad \dots, \quad A^{n+1} = A^nA, \quad \dots, \quad \text{and} \quad A^0 = I$$

Polynomials in the matrix A are also defined. Specifically, for any polynomial

$$f(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$$

where the a_i are scalars in K , $f(A)$ is defined to be the following matrix:

$$f(A) = a_0I + a_1A + a_2A^2 + \dots + a_nA^n$$

Note that $f(A)$ is obtained from $f(x)$ by substituting the matrix A for the variable x and substituting the scalar matrix a_0I for the scalar a_0 .

Powers of Matrices, Polynomials in Matrices

Let A be an n -square matrix over a field K . Powers of A are defined as follows:

$$A^2 = AA, \quad A^3 = A^2A, \quad \dots, \quad A^{n+1} = A^nA, \quad \dots, \quad \text{and} \quad A^0 = I$$

Polynomials in the matrix A are also defined. Specifically, for any polynomial

$$f(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$$

where the a_i are scalars in K , $f(A)$ is defined to be the following matrix:

$$f(A) = a_0I + a_1A + a_2A^2 + \dots + a_nA^n$$

Note that $f(A)$ is obtained from $f(x)$ by substituting the matrix A for the variable x and substituting the scalar matrix a_0I for the scalar a_0 . If $f(A)$ is the zero matrix, then A is called a zero or root of $f(x)$.

Powers of Matrices, Polynomials in Matrices

Example

Suppose $A = \begin{pmatrix} 1 & 2 \\ 3 & -4 \end{pmatrix}$. Then

$$A^2 = \begin{pmatrix} 1 & 2 \\ 3 & -4 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 3 & -4 \end{pmatrix} = \begin{pmatrix} 7 & -6 \\ -9 & 22 \end{pmatrix} \quad \text{and}$$

$$A^3 = A^2 A = \begin{pmatrix} 7 & -6 \\ -9 & 22 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 3 & -4 \end{pmatrix} = \begin{pmatrix} -11 & 38 \\ 57 & -106 \end{pmatrix}$$

Suppose $f(x) = 2x^2 - 3x + 5$ and $g(x) = x^2 + 3x - 10$. Then

$$f(A) = 2 \begin{pmatrix} 7 & -6 \\ -9 & 22 \end{pmatrix} - 3 \begin{pmatrix} 1 & 2 \\ 3 & -4 \end{pmatrix} + 5 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 16 & -18 \\ -27 & 61 \end{pmatrix}$$

$$g(A) = \begin{pmatrix} 7 & -6 \\ -9 & 22 \end{pmatrix} + 3 \begin{pmatrix} 1 & 2 \\ 3 & -4 \end{pmatrix} - 10 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$$

Thus, A is a zero of the polynomial $g(x)$

Invertible (Nonsingular) Matrices

Invertible (Nonsingular) Matrices

A square matrix A is said to be invertible or nonsingular if there exists a matrix B such that

$$AB = BA = I$$

where I is the identity matrix. Such a matrix B is unique. That is, if $AB_1 = B_1A = I$ and $AB_2 = B_2A = I$ then

$$B_1 = B_1I = B_1(AB_2) = (B_1A)B_2 = IB_2 = B_2$$

We call such a matrix B the **inverse** of A and denote it by A^{-1} . Observe that the above relation is symmetric; that is, if B is the inverse of A , then A is the inverse of B .

Invertible (Nonsingular) Matrices

Example

Suppose that $A = \begin{pmatrix} 2 & 5 \\ 1 & 3 \end{pmatrix}$ and $B = \begin{pmatrix} 3 & -5 \\ -1 & 2 \end{pmatrix}$. Then

$$AB = \begin{pmatrix} 6-5 & -10+10 \\ 3-3 & -5+6 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$BA = \begin{pmatrix} 6-5 & 15-15 \\ -2+2 & -5+6 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Thus, A and B are inverses. It is known that $AB = I$ if and only if $BA = I$. Thus, it is necessary to test only one product to determine whether or not two given matrices are inverses.

Invertible (Nonsingular) Matrices

Now suppose A and B are invertible. Then AB is invertible and $(AB)^{-1} = B^{-1}A^{-1}$. More generally, if A_1, A_2, \dots, A_k are invertible, then their product is invertible and

$$(A_1 A_2 \dots A_k)^{-1} = A_k^{-1} \dots A_2^{-1} A_1^{-1}$$

the product of the inverses in the reverse order.

?