
Problem Set - 09

MA 1201

Spring Sem, 2025

LINEAR TRANSFORMATION AND MATRIX REPRESENTATION

1. In this exercise, $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a function. For each of the following parts, state why T is *not* linear.

- (a) $T(a_1, a_2) = (1, a_2)$
- (b) $T(a_1, a_2) = (a_1, a_2^2)$
- (c) $T(a_1, a_2) = (\sin a_1, 0)$
- (d) $T(a_1, a_2) = (|a_1|, a_2)$
- (e) $T(a_1, a_2) = (a_1 + 1, a_2)$

For Exercises 2 through 6, find the matrix representation of the linear map with respect to standard basis in each case. Note that the matrices E_{ij} - with ij -th entry 1 and all other entries 0, $1 \leq i \leq m, 1 \leq j \leq n$, forms the standard basis of $M_{m \times n}(\mathbb{F})$ and $\{1, x, \dots, x^k\}$ forms the standard basis of $P_k(\mathbb{R})$.

2. $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ defined by

$$T(a_1, a_2, a_3) = (a_1 - a_2, 2a_3).$$

3. $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ defined by

$$T(a_1, a_2) = (a_1 + a_2, 0, 2a_1 - a_2).$$

4. $T : M_{2 \times 3}(\mathbb{R}) \rightarrow M_{2 \times 2}(\mathbb{R})$ defined by

$$T \left(\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} \right) = \begin{pmatrix} 2a_{11} - a_{12} & a_{13} + 2a_{12} \\ 0 & 0 \end{pmatrix}.$$

5. $T : P_2(\mathbb{R}) \rightarrow P_3(\mathbb{R})$ defined by

$$T(f(x)) = xf(x) + f'(x).$$

6. $T : M_{n \times n}(F) \rightarrow F$ defined by $T(A) = \text{tr}(A)$ (called *Trace* of A), where

$$\text{tr}(A) = \sum_{i=1}^n A_{ii}.$$

7. Suppose that $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is linear, $T(1, 0) = (1, 4)$, and $T(1, 1) = (2, 5)$. What is $T(2, 3)$? Is T one-to-one?

8. Prove that there exists a linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ such that $T(1, 1) = (1, 0, 2)$ and $T(2, 3) = (1, -1, 4)$. What is $T(8, 11)$?

9. For each of the following linear transformations T , determine whether T is invertible and justify your answer.

- (a) $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ defined by $T(a_1, a_2) = (3a_1 - a_2, a_2, 4a_1)$
 (b) $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(a_1, a_2, a_3) = (3a_1 - 2a_3, a_2, 3a_1 + 4a_2)$
 (c) $T : P_3(\mathbb{R}) \rightarrow P_2(\mathbb{R})$ defined by $T(p(x)) = p'(x)$
 (d) $T : M_{2 \times 2}(\mathbb{R}) \rightarrow P_2(\mathbb{R})$ defined by

$$T\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right) = a + 2bx + (c + d)x^2$$

- (e) $T : M_{2 \times 2}(\mathbb{R}) \rightarrow M_{2 \times 2}(\mathbb{R})$ defined by

$$T\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right) = \begin{pmatrix} a+b & a \\ c & c+d \end{pmatrix}$$

10. Which of the following pairs of vector spaces are isomorphic? Justify your answers.

- (a) \mathbb{F}^3 and $P_3(\mathbb{F})$
 (b) \mathbb{F}^4 and $P_3(\mathbb{F})$
 (c) $M_{2 \times 2}(\mathbb{R})$ and $P_3(\mathbb{R})$
 (d) $V = \{A \in M_{2 \times 2}(\mathbb{R}) : \text{tr}(A) = 0\}$ and \mathbb{R}^4
11. Let $g(x) = 3 + x$. Let $T : P_2(\mathbb{R}) \rightarrow P_2(\mathbb{R})$ and $U : P_2(\mathbb{R}) \rightarrow \mathbb{R}^3$ be the linear transformations respectively defined by

$$T(f(x)) = f'(x)g(x) + 2f(x) \quad \text{and} \quad U(a + bx + cx^2) = (a + b, c, a - b)$$

Let β and γ be the standard ordered bases of $P_2(\mathbb{R})$ and \mathbb{R}^3 , respectively.

- (a) Compute $[U]_{\gamma}^{\beta}$, $[T]_{\beta}^{\beta}$, and $[UT]_{\gamma}^{\beta}$ directly and verify that matrix representation of the composition is matrix multiplication of the individual matrix in the same order.
 (b) Let $h(x) = 3 - 2x + x^2$. Compute $[h]_{\beta}$ and $[U(h)]_{\gamma}$. Then verify $[U(h)]_{\gamma} = [U]_{\gamma}^{\beta}[h]_{\beta}$.
12. Find linear transformations $U, T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that $UT = 0$ (the zero transformation) but $TU \neq 0$.
13. Let T be the linear operator on \mathbb{R}^2 defined by

$$T\begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 2a + b \\ a - 3b \end{pmatrix},$$

let β be the standard ordered basis for \mathbb{R}^2 , and let

$$\beta' = \left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \end{pmatrix} \right\}.$$

Use the change of basis matrix to find $[T]_{\beta'}^{\beta'}$.

14. Let T be the linear operator on $P_1(\mathbb{R})$ defined by $T(p(x)) = p'(x)$, the derivative of $p(x)$. Let $\beta = \{1, x\}$ and $\beta' = \{1+x, 1-x\}$. Use the change of basis matrix to find $[T]_{\beta'}^{\beta'}$ and verify by computing independently.
15. If the transformation T is a reflection across the 45° line in the plane, find its matrix with respect to the standard basis $w_1 = (1, 3)$, $w_2 = (2, -1)$, and also with respect to $v_1 = (1, 2)$, $v_2 = (3, -1)$. Show also that those matrices are similar and find the matrix which gives the similarity.

DETERMINANT

16. Let Q be an orthogonal $n \times n$ matrix, that is, $Q^T Q = I$. Prove that $\det Q$ equals $+1$ or -1 .

17. Show that

$$\det \begin{pmatrix} a & b & c \\ b & c & a \\ c & a & b \end{pmatrix} = -(a^3 + b^3 + c^3 - 3abc).$$

18. Without expanding the determinant prove that

$$\det \begin{pmatrix} 0 & a & b \\ a & 0 & c \\ b & -c & 0 \end{pmatrix} = 0.$$

19. Use row operations to verify that the 3 by 3 "Vandermonde determinant" is

$$\det \begin{pmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{pmatrix} = (b-a)(c-a)(c-b).$$

20. Use the problem above to show that if $s_r = a^r + b^r + c^r$, then

$$\det \begin{pmatrix} s_0 & s_1 & s_2 \\ s_1 & s_2 & s_3 \\ s_2 & s_3 & s_4 \end{pmatrix} = (b-a)^2(c-a)^2(c-b)^2.$$

21. True or false, with reason if true and counterexample if false:

- (a) If A and B are identical except that $b_{11} = 2a_{11}$, then $\det B = 2 \det A$.
- (b) The determinant is the product of the pivots.
- (c) If A is invertible and B is singular (*not invertible*), then $A + B$ is invertible.
- (d) If A is invertible and B is singular, then AB is singular.
- (e) The determinant of $AB - BA$ is zero.

22. (a) If every row of A adds to zero, prove that $\det A = 0$.

- (b) If every row of A adds to 1, prove that $\det(A - I) = 0$. Show by example that this does not imply $\det A = 1$.

23. Find these 4 by 4 determinants by Gaussian elimination:

$$\det \begin{pmatrix} 11 & 12 & 13 & 14 \\ 21 & 22 & 23 & 24 \\ 31 & 32 & 33 & 34 \\ 41 & 42 & 43 & 44 \end{pmatrix} \quad \text{and} \quad \det \begin{pmatrix} 1 & t & t^2 & t^3 \\ t & 1 & t & t^2 \\ t^2 & t & 1 & t \\ t^3 & t^2 & t & 1 \end{pmatrix}.$$

24. Given $A = \begin{pmatrix} 4 & 2 \\ 1 & 3 \end{pmatrix}$, find the determinants of $A - \lambda I = \begin{pmatrix} 4 - \lambda & 2 \\ 1 & 3 - \lambda \end{pmatrix}$, for every $\lambda \in \mathbb{R}$.

For which values of λ is $A - \lambda I$ a singular matrix (*not invertible*)?

25. Let $A = (a_{ij}) \in M_n(\mathbb{R})$.

(a) For any $n \geq 3$, if $a_{ij} = i + j$, show that $\det A = 0$.

(b) For any $n \geq 2$, if $a_{ij} = ij$, show that $\det A = 0$.

26. If $A = \begin{pmatrix} 1 & 0 & 1 \\ 3 & 4 & 5 \\ 2 & 3 & 4 \end{pmatrix}$, find A^{-1} .

27. Find the matrix A , if $A^{-1} = \begin{pmatrix} 3 & -1 & 1 \\ 1 & -2 & 3 \\ 3 & -3 & 4 \end{pmatrix}$.

28. Solve the following system of equations by Cramer's rule:

$$x + 2y - 3z = 1$$

$$2x - y + z = 4$$

$$x + 3y = 5.$$

29. Solve by Cramer's rule:

$$(i) \quad x + y + z = 6$$

$$x + 2y + 3z = 14$$

$$x - y + z = 2,$$

$$(ii) \quad x + y + z = 1$$

$$ax + by + cz = 1$$

$$a^2x + b^2y + c^2z = 1, \quad a \neq b \neq c.$$

EIGEN-VALUES AND EIGEN-VECTORS

30. Find the rank and all four eigenvalues for the checker board matrix $C = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix}$. Which

eigenvectors correspond to nonzero eigenvalues?

31. (a) Construct 2 by 2 matrices A and B such that the eigenvalues of AB are not the products of the eigenvalues of A and B , and the eigenvalues of $A + B$ are not the sums of the individual eigenvalues.

(b) Construct 3 by 3 matrices A and B with same properties as in part (a).

32. Prove that the real eigenvalues of A equal the eigenvalues of A^T . Show by an example that the eigenvectors of A and A^T need not be the same.

33. Every permutation matrix leaves $x = (1, 1, \dots, 1)$ unchanged. Thus, $\lambda = 1$ is an eigenvalue for every permutation matrix. Find two more eigenvalues for the following permutations:

$$P_1 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \quad \text{and} \quad P_2 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

34. Find the 2 by 2 matrix A whose eigenvalues are 1 and 4, with the corresponding eigenvectors being $(3, 1)$ and $(2, 1)$ respectively.
35. Which of these matrices cannot be diagonalized? Recall that diagonalizable means that there exists a basis consisting of eigen vectors.

$$A_1 = \begin{pmatrix} 2 & -2 \\ 2 & -2 \end{pmatrix}, \quad A_2 = \begin{pmatrix} 2 & 0 \\ 2 & -2 \end{pmatrix}, \quad A_3 = \begin{pmatrix} 2 & 0 \\ 2 & 2 \end{pmatrix}.$$

36. Describe all matrices S that diagonalize the matrix $A = \begin{pmatrix} 4 & 0 \\ 1 & 2 \end{pmatrix}$. Then describe all matrices that diagonalize A^{-1} .
37. If the eigenvalues of a 3 by 3 matrix A are 1, 1, 2, which of the following are certain to be true? Give a reason if true or a counterexample if false:

- (a) A is invertible.
- (b) A is diagonalizable.
- (c) A is not diagonalizable.

38. Suppose the only eigenvectors of a 3 by 3 matrix A are multiples of $(1, 0, 0)$. True or false:

- (a) A is not invertible.
- (b) A has a repeated eigenvalue.
- (c) A is not diagonalizable.

39. Diagonalize $A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$, that is, find a matrix S such that $S^{-1}AS = D$ is a diagonal matrix. Use it to prove that for any $k \in \mathbb{N}$, we have that $A^k = \frac{1}{2} \begin{pmatrix} 3^k + 1 & 3^k - 1 \\ 3^k - 1 & 3^k + 1 \end{pmatrix}$.

40. Given $A \in M_n(\mathbb{R})$, explain why A is never similar to $A + I$.
41. Let $A, B \in M_n(\mathbb{R})$. If A or B is invertible then show that BA is similar to AB . Give an example of matrices A and B such BA is not similar to AB .
42. Let $A, B \in M_n(\mathbb{R})$. Prove that AB has the same eigenvalues as BA .