

SOLUTIONS OF HARTSHORNE'S ALGEBRAIC GEOMETRY

CONTENTS

1.	Varieties	1
1.1.	Affine Varieties	1
1.2.	Projective Varieties	3
1.3.	Morphisms	3
1.4.	Rational Maps	4
1.5.	Nonsingular Varieties	4
1.6.	Nonsingular Curves	4
1.7.	Intersections in Projective Space	4

1. VARIETIES

1.1. Affine Varieties.

Exercise 1.1.1.

- (a) Let Y be the plane curve $y = x^2$ (i.e., Y is the zero set of the polynomial $f = y - x^2$). Show that $A(Y)$ is isomorphic to a polynomial ring in one variable over k .
- (b) Let Z be the plane curve $xy = 1$. Show that $A(Z)$ is not isomorphic to a polynomial ring in one variable over k .
- (c) Let f be any irreducible quadratic polynomial in $k[x, y]$, and let W be the conic defined by f . Show that $A(W)$ is isomorphic to $A(Y)$ or $A(Z)$. Which one is it when?

Proof.

- (a) Since $y - x^2$ is an irreducible polynomial, $(y - x^2)$ is a prime ideal of $k[x, y]$, so

$$\begin{aligned} A(Y) &= k[x, y]/I(Y) = k[x, y]/I(Z(y - x^2)) \\ &= k[x, y]/\sqrt{(y - x^2)} = k[x, y]/(y - x^2) \end{aligned}$$

Consider the map

$$\varphi : k[x, y]/(y - x^2) \rightarrow k[t]$$

$$f(x, y) + (y - x^2) \mapsto f(t, t^2)$$

If $y - x^2 \mid f(x, y)$, then $f(t, t^2) = 0$ so it is well defined ring homomorphism. And define

$$\psi : k[t] \rightarrow k[x, y]/(y - x^2)$$

$$g(t) \mapsto g(x) + (y - x^2)$$

We want to show that $\varphi \circ \psi = 1_{k[t]}$ and $\psi \circ \varphi = 1_{k[x, y]/(y - x^2)}$ so $k[t] \cong k[x, y]/(y - x^2)$. The first one is easy because

$$\varphi(\psi(g(t))) = \varphi(g(x) + (y - x^2)) = g(t)$$

Take any $f(x, y) = \sum_{i,j} a_{ij} x^i y^j \in k[x, y]$. Then,

$$\psi(\varphi(f(x, y) + (y - x^2))) = \psi(f(t, t^2)) = f(x, x^2) + (y - x^2)$$

so we need to show that $f(x, y) - f(x, x^2) \in (y - x^2)$.

$$\begin{aligned} f(x, y) - f(x, x^2) &= \sum_{i,j} a_{ij} x^i y^j - \sum_{i,j} a_{ij} x^i x^{2j} \\ &= \sum_{i,j} a_{ij} x^i (y^j - (x^2)^j) \\ &= \sum_{i,j} a_{ij} x^i (y - x^2) (y^{j-1} + y^{j-2}(x^2)^1 + \cdots + y^1(x^2)^{j-2} + (x^2)^{j-1}) \end{aligned}$$

Hence, $A(Y) \cong k[t]$

(b) Note that

$$A(Z) = k[x, y]/(xy - 1)$$

and suppose $\theta : A(Z) \rightarrow k[t]$ is a ring isomorphism. Then, $x + (xy - 1), y + (xy - 1) \in A(Z)$ are unit elements, so $\theta(x + (xy - 1)), \theta(y + (xy - 1)) \in k[t]$ are unit elements, which implies they are in k . Then there is no element in $A(Z)$ that goes to t through the map θ , which is a contradiction with the fact that θ is an isomorphism.

(c) a

□

Exercise 1.1.2. *The Twisted Cubic Curve.* Let $Y \subseteq \mathbb{A}^3$ be the set $Y = \{(t, t^2, t^3) \mid t \in k\}$. Show that Y is an affine variety of dimension 1. Find generators for the ideal $I(Y)$. Show that $A(Y)$ is isomorphic to a polynomial ring in one variable over k . We say that Y is given by the *parametric representation* $x = t, y = t^2, z = t^3$.

Proof.

□

Exercise 1.1.3. Let Y be the algebraic set in \mathbb{A}^3 defined by the two polynomials $x^2 - yz$ and $xz - x$. Show that Y is a union of three irreducible components. Describe them and find their prime ideals.

Proof.

□

Exercise 1.1.4. If we identify \mathbb{A}^2 with $\mathbb{A}^1 \times \mathbb{A}^1$ in the natural way, show that the Zariski topology on \mathbb{A}^2 is not the product topology of the Zariski topologies on the two copies of \mathbb{A}^1 .

Proof.

□

Exercise 1.1.5. Show that a k -algebra B is isomorphic to the affine coordinate ring of some algebraic set in \mathbb{A}^n , for some n , if and only if B is a finitely generated k -algebra with no nilpotent elements.

Proof.

□

Exercise 1.1.6. Any nonempty open subset of an irreducible topological space is dense and irreducible. If Y is a subset of a topological space X , which is irreducible in its induced topology, then the closure \bar{Y} is also irreducible.

Proof.

□

Exercise 1.1.7.

- (a) Show that the following conditions are equivalent for a topological space X : (i) X is noetherian; (ii) every nonempty family of closed subsets has a minimal element; (iii) X satisfies the ascending chain condition for open subsets; (iv) every nonempty family of open subsets has a maximal element.
- (b) A noetherian topological space is *quasi-compact*, i.e., every open cover has a finite subcover.
- (c) Any subset of a noetherian topological space is noetherian in its induced topology.
- (d) A noetherian space which is also Hausdorff must be a finite set with the discrete topology.

Proof.

□

Exercise 1.1.8. Let Y be an affine variety of dimension r in \mathbb{A}^n . Let H be a hypersurface in \mathbb{A}^n , and assume that $Y \not\subseteq H$. Then every irreducible component of $Y \cap H$ has dimension $r - 1$. (See (7.1) for a generalization.)

Proof.

□

Exercise 1.1.9. Let $\mathfrak{a} \subseteq A = k[x_1, \dots, x_n]$ be an ideal which can be generated by r elements. Then every irreducible component of $Z(\mathfrak{a})$ has dimension $\geq n - r$.

Proof.

□

Exercise 1.1.10.

- (a) If Y is any subset of a topological space X , then $\dim Y \leq \dim X$.
- (b) If X is a topological space which is covered by a family of open subsets $\{U_i\}$, then $\dim X = \sup \dim U_i$.
- (c) Give an example of a topological space X and a dense open subset U with $\dim U < \dim X$.
- (d) If Y is a closed subset of an irreducible finite-dimensional topological space X , and if $\dim Y = \dim X$, then $Y = X$.
- (e) Give an example of a noetherian topological space of infinite dimension.

Proof.

□

Exercise 1.1.11. Let $Y \subseteq \mathbb{A}^3$ be the curve given parametrically by $x = t^3, y = t^4, z = t^5$. Show that $I(Y)$ is a prime ideal of height 2 in $k[x, y, z]$ which cannot be generated by 2 elements. We say Y is *not a local complete intersection* cf. (Ex. 2.17).

Proof.

□

Exercise 1.1.12. Give an example of an irreducible polynomial $f \in \mathbb{R}[x, y]$, whose zero set $Z(f)$ in $\mathbb{A}_{\mathbb{R}}^2$ is not irreducible (cf. 1.4.2).

Proof.

□

1.2. Projective Varieties.

Exercise 1.2.1.

Proof.

□

1.3. Morphisms.

Exercise 1.3.1.

Proof.

□

1.4. Rational Maps.**Exercise 1.4.1.***Proof.*

□

1.5. Nonsingular Varieties.**Exercise 1.5.1.***Proof.*

□

1.6. Nonsingular Curves.**Exercise 1.6.1.***Proof.*

□

1.7. Intersections in Projective Space.**Exercise 1.7.1.***Proof.*

□