

①

## Test of Significance for Single mean

Let  $x_1, x_2, x_3, \dots, x_n$  be a random sample of size 'n' from a large sample  $x_1, x_2, \dots, x_N$  (of size N) with mean  $\mu$  and variance  $\sigma^2$

Then sample mean ( $\bar{x}$ ) and variance ( $s^2$ )

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i, \quad s^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$$

Also, the standard error (S.E.) of mean of a random sample of size (n) from a population with variance ( $\sigma^2$ )

$$\text{is } \sigma / \sqrt{n}$$

$$\text{i.e., } S.E.(\bar{x}) = \sqrt{\frac{\sigma^2}{n}} = \frac{\sigma}{\sqrt{n}}$$

For large sample, the standard normal variate corresponding  $\bar{x}$  is given by

$$Z = \frac{\bar{x} - \mu}{\sigma / \sqrt{n}}$$

(2)

Under null hypothesis ( $H_0$ ) that the sample has been drawn from a population with mean ' $\mu$ ' and variance ' $\sigma^2$ '

ie there is no significant difference between the sample mean ( $\bar{x}$ ) and population mean ( $\mu$ ) for large sample the test statistic is

$$Z = \frac{\bar{x} - \mu}{\sigma / \sqrt{n}}$$

If the population standard deviation ' $\sigma$ ' is not known

$$Z = \frac{\bar{x} - \mu}{s / \sqrt{n}}$$

where ' $s$ ' is the standard deviation of the sample.



## Confidence limits for $\mu$

① If the LOS is  $\alpha$  and  $Z_\alpha$  is the critical value then

$$-Z_\alpha < \left| \frac{\bar{x} - \mu}{\sigma/\sqrt{n}} \right| < Z_\alpha$$

The limit of population mean ' $\mu$ ' is given by

$$\bar{x} - Z_\alpha \frac{\sigma}{\sqrt{n}} < \mu < \bar{x} + Z_\alpha \frac{\sigma}{\sqrt{n}}$$

② 95% confidence interval for  $\mu$  at 5% level of significance is

$$\bar{x} - 1.96 \frac{\sigma}{\sqrt{n}} < \mu < \bar{x} + 1.96 \frac{\sigma}{\sqrt{n}}$$

③ 99% confidence interval for  $\mu$  at 1% level of significance is

(4)

$$\bar{x} - 2.58 \frac{\sigma}{\sqrt{n}} < \mu < \bar{x} + 2.58 \frac{\sigma}{\sqrt{n}}$$

③ In sampling from a finite population of size  $N$  the corresponding 95% and 99% confidence limit for  $\mu$  are

$$\bar{x} \pm 1.96 \frac{\sigma}{\sqrt{n}} \sqrt{\frac{N-n}{N-1}} \quad \left[ \begin{array}{c} \text{for} \\ 95\% \end{array} \right]$$

$$\bar{x} \pm 2.58 \frac{\sigma}{\sqrt{n}} \sqrt{\frac{N-n}{N-1}} \quad \left( \begin{array}{c} \text{for} \\ 99\% \end{array} \right)$$

Q-① The average marks in Mathematics of a sample of 100 students was 51 with a S.D of 6 marks. Could this have been a random sample from a population with average marks 50?



(5)

Sol Given that Sample size  $n = 100$

$\bar{x}$  = Sample mean = 51

Sample S.D  $s = 6$

population mean  $\mu = 50$

( $\sigma$  is unknown) here

consider null hypothesis

$H_0$ : the sample drawn from  
a population with  $\mu = 50$

$H_1$ :  $\mu \neq 50$

Under  $H_0$ ,

$$Z = \frac{\bar{x} - \mu}{s/\sqrt{n}} = \frac{51 - 50}{6/\sqrt{100}}$$

$$= \frac{1}{6/10} = \frac{10}{6}$$

$$\Rightarrow Z = 1.6666 \approx \text{to } 1.67$$

$$\Rightarrow |Z| = 1.67$$

The ladder of success is never crowded at the top

(6)

Conclusion: Since  $|z| = 1.87 < 1.96$

When  $z$  is the significant value of 2 at 5% L.O.I.

$H_0$  is accepted.

~~Q-2~~ A normal population has a mean of 6.8 and S.D 1.5.

A sample of 400 members gave a mean of 6.75, is the difference significant.

~~Soln~~  $H_0$ : There is no significant difference between  $\bar{x}$  and  $\mu$ .

$H_1$ : There is significant difference between  $\bar{x}$  and  $\mu$ .

Priorities

Success is only another form of failure if we forget what our priorities should be



Given that  $\mu = 6.8$

$$\sigma = 1.5$$

$$\bar{x} = 6.75, n = 400$$

$$\therefore |Z| = \left| \frac{\bar{x} - \mu}{\sigma / \sqrt{n}} \right|$$

$$= \left| \frac{6.75 - 6.8}{1.5 / \sqrt{400}} \right|$$

$$= | -0.67 | = 0.67$$

Conclusion

$$|Z| < Z_{\alpha} = 1.96$$

at 5% level of significance.

$H_0$  is accepted.

Hence there is no significant difference between  $\bar{x}$  and  $\mu$ .

Failure is the condiment that gives success its flavor

10

Friday

Wk 15 - Day 100

③

P  
2  
1513 14 15 16 17 18 19 20 21 22 23 24 25 26  
27 28 29 30

~~Q. ③~~ A random sample of 900 members has mean 3.4 cm.

Can it be reasonably regarded as a sample from a large sample population of mean 3.2 cms and S.D 2.3 cm.

Soln Given that

$$n = \text{Sample size} = 900$$

$$\bar{x} = 3.4 = \text{Sample mean}$$

$$\mu = 3.2 \text{ (Population mean)}$$

$$\sigma = 2.3 \text{ (Population S.D)}$$

Consider null hypothesis

$H_0$ : Assume the sample is

Priorities

drawn from a large

population with mean 3.2

and S.D = 2.3

Men are failures, not because they are stupid, but because they are not sufficiently impassioned



(9)

~~Q~~  $H_1: \mu \neq 3.2$

Under  $H_0$

$$Z = \frac{\bar{x} - \mu}{\sigma / \sqrt{n}} = \frac{3.4 - 3.2}{2.3 / \sqrt{900}} = 0.261$$

$$|Z| = 0.261$$

Conclusion:

Now  $|Z| = 0.261 < 1.96$  (2 $\alpha$ )

The significant value of Z is less than 1.96.

$H_0$  is accepted therefore the sample is drawn from the population with  $\mu = 3.2$ .

Sunday 12

and S.D 2.3.

(15)

Q. The mean weight obtained from a random sample of size 100 is 64 gm. The S.D. is 3. The <sup>mean</sup> 1st distribution of the population is 67 gm. at 5% level. Also set up 99% confidence limits of the mean weight of the population.

Soln Given that

$$n = 100, \mu = 67$$

$$\bar{x} = 64, \sigma = 3$$

H<sub>0</sub>: There is no significant difference between sample and population mean.

A<sub>1</sub>: There is significant

There are men in the world who derive an exaltation from the proximity of disaster and ruin, as other from success



(11)

difference between sample mean  
 and population mean

under  $H_0$

$$|Z| = \left| \frac{\bar{x} - \mu}{\sigma/\sqrt{n}} \right| = \left| \frac{64 - 62}{3/\sqrt{100}} \right| = 10$$

$$|Z| = 10$$

Conclusion =

Since  $|Z| > 1.96 (2\alpha)$

the significant value of 2  
 at 5% L.O.S.

$H_0$  is rejected.

i.e. the sample is not drawn from  
 the population with  $\mu = 62$

⑪ 99% confidence limit is  
 given by

$$\bar{x} \pm 2.58 \frac{\sigma}{\sqrt{n}} = 64 \pm 2.58 \cdot \frac{3}{\sqrt{100}} \\ = 64.774 \quad 63.226$$