


①

Simpson's $\frac{1}{3}$ rd Rule :-

When $n=2$ in Newton-Cotes's quadrature formula we get

Simpson's $\frac{1}{3}$ rd Rule.

Let $f(x)$ be a function which is defined at $(n+1)$ points such as $x_0, x_1, x_2, \dots, x_n$ which are equispaced.



$h \rightarrow$ step size / step length.

Now
$$\int_a^b f(x) dx = \int_{x_0}^{x_n} f(x) dx$$

Then by Newton-Cotes's quadrature formula we get

$$\int_a^b f(x) dx = \lambda_0 f_0 + \lambda_1 f_1 + \lambda_2 f_2$$

where $\lambda_0, \lambda_1, \lambda_2$ can be determined by following formula

$$\lambda_k = \frac{(-1)^{n-k} \cdot h}{k! \cdot (n-k)!} \int_0^n \frac{s(s-1)(s-2) \cdots (s-(k-1))}{(s-(k+1)) \cdots (s-n)} ds$$

for $k=0, 1, 2, \dots, n$

Now taking $k=0, n=2$ we get

$$\lambda_0 = \frac{(-1)^{2-0} \cdot h}{0! \cdot (2-0)!} \int_0^2 (s-1)(s-2) ds$$

$$= \frac{(-1)^2 \cdot h}{1 \cdot 2!} \int_0^2 (s^2 - 3s + 2) ds$$

(3)

$$= \frac{h}{2} \left[\int_0^2 s^2 ds - 3 \int_0^2 s ds + 2 \int_0^2 ds \right]$$

$$= \frac{h}{2} \left[\left[\frac{s^3}{3} \right]_0^2 - 3 \left[\frac{s^2}{2} \right]_0^2 + 2 \left[s \right]_0^2 \right]$$

$$= \frac{h}{2} \left[\left(\frac{2^3}{3} - 0 \right) - 3 \left(\frac{2^2}{2} - 0 \right) + 2(2 - 0) \right]$$

$$= \frac{h}{2} \left[\frac{8}{3} - 6 + 4 \right]$$

$$= \frac{h}{2} \left[\frac{8}{3} - 2 \right] = \frac{h}{2} \cdot \left(\frac{2}{3} \right) = \frac{h}{3}$$

$$\boxed{\lambda_0 = \frac{h}{3}}$$

Next taking $n=2$, $k=1$ we get

$$\lambda_1 = \frac{(-1)^{2-1} \cdot h}{1! (2-1)!} \int_0^2 s(s-2) ds$$

(4)

$$= \frac{(-1) \cdot h}{1 \cdot 1} \int_0^2 (s^2 - 2s) ds$$

$$= -h \left[\int_0^2 s^2 ds - 2 \int_0^2 s ds \right]$$

$$= -h \left[\left[\frac{s^3}{3} \right]_0^2 - 2 \left[\frac{s^2}{2} \right]_0^2 \right]$$

$$= -h \left[\left(\frac{2^3}{3} - 0 \right) - (2^2 - 0) \right]$$

$$= -h \left[\frac{8}{3} - 4 \right]$$

$$= -h \left(-\frac{4}{3} \right) = \frac{4h}{3}$$

$$\therefore \boxed{\lambda_1 = \frac{4h}{3}}$$

A gain taking $k=2$, $n=2$ magnet,

⑤

$$\lambda_2 = \frac{(-1)^{2-2} \cdot h}{2! (2-2)!} \int_0^2 s(s-1) ds$$

$$= \frac{1 \cdot h}{2 \cdot 0!} \int_0^2 (s^2 - s) ds$$

$$= \frac{h}{2} \left[\int_0^2 s^2 ds - \int_0^2 s ds \right]$$

$$= \frac{h}{2} \left[\left[\frac{s^3}{3} \right]_0^2 - \left[\frac{s^2}{2} \right]_0^2 \right]$$

$$= \frac{h}{2} \left[\left(\frac{2^3}{3} - 0 \right) - \left(\frac{2^2}{2} - 0 \right) \right]$$

$$= \frac{h}{2} \left[\frac{8}{3} - 2 \right] = \frac{h}{2} \left(\frac{2}{3} \right) = \frac{h}{3}$$

$$\therefore \boxed{\lambda_2 = \frac{h}{3}}$$

Putting the value of $\lambda_0, \lambda_1, \lambda_2$ in eqn (1) we get,

03 / 2018	01 02 03 04	05 06 07 08 09 10 11	12 13 14 15 16 17 18	19 20 21 22 23 24 25	26 27 28 29 30 31
MARCH / Week	9	10	11	12	13

$$\int_a^b f(x) dx = \lambda_0 f_0 + \lambda_1 f_1 + \lambda_2 f_2$$

$$= \frac{h}{3} f_0 + \frac{4h}{3} f_1 + \frac{h}{3} f_2$$

$$= \frac{h}{3} [f_0 + 4f_1 + f_2]$$

$$= \frac{h}{3} [f(a) + 4f(m) + f(b)]$$

$$\int_a^b f(x) dx = \int_{x_0}^{x_2} f(x) dx = \frac{h}{3} [f(x_0) + 4f(x_1) + f(x_2)]$$

Which is called Simpson's
 $\frac{1}{3}$ rd Rule.

OB

$$\int_a^b f(x) dx = \frac{h}{3} [f(a) + 4f\left(\frac{a+b}{2}\right) + f(b)]$$

$$\int_a^b f(x) dx = \frac{b-a}{6} [f(a) + 4f\left(\frac{a+b}{2}\right) + f(b)]$$

(7)

Note :- It works on even number of intervals. (n is multiple of 2)
i.e. no. of ordinates must be odd.

Composite Simpson's $\frac{1}{3}$ rd Rule :-

The Composite Simpson's $\frac{1}{3}$ rd Rule can be written as

$$\begin{aligned} \int_a^b f(x) dx &= \int_{x_0}^{x_n} f(x) dx \\ &= \int_{x_0}^{x_2} f(x) dx + \int_{x_2}^{x_4} f(x) dx + \int_{x_4}^{x_6} f(x) dx \\ &\quad + \dots + \int_{x_{n-2}}^{x_n} f(x) dx \end{aligned}$$

where n must be even.

03 / 2018	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	
MARCH / Week	9				10				11				12				13															

$$= \frac{h}{3} [f(x_0) + 4f(x_1) + f(x_2)]$$

$$+ \frac{h}{3} [f(x_2) + 4f(x_3) + f(x_4)]$$

$$+ \frac{h}{3} [f(x_4) + 4f(x_5) + f(x_6)]$$

$$+ \dots$$

$$+ \frac{h}{3} [f(x_{n-2}) + 4f(x_{n-1}) + f(x_n)]$$

$$= \frac{h}{3} [f(x_0) + 4f(x_1) + 4f(x_3) + 4f(x_5)$$

$$+ \dots + 4f(x_{n-1}) + 2f(x_2)$$

$$+ 2f(x_4) + 2f(x_6) + \dots$$

$$+ 2f(x_{n-2}) + f(x_n)]$$

⑨

$$= \frac{h}{3} \left[(f(x_0) + f(x_n)) + 4(f(x_1) + f(x_3) + \dots + f(x_{n-1})) + 2(f(x_2) + f(x_4) + \dots + f(x_{n-2})) \right]$$

$$\int_a^b f(x) dx = \int_{x_0}^{x_n} f(x) dx$$

$$= \frac{h}{3} \left[(y_0 + y_n) + 4(y_1 + y_3 + \dots + y_{n-1}) + 2(y_2 + y_4 + \dots + y_{n-2}) \right]$$

which is called ^{Composite} Simpson's $\frac{1}{3}$ rd Rule.

It can be also expressed as

$$\int_a^b f(x) dx = \frac{h}{3} [A + 4B + 2C]$$

09/2018

01 02 03 04

05 06 07 08 09 10 11

12 13 14 15 16 17 18

19 20 21 22 23 24 25

26 27 28 29 30 31

MARCH / Week

9

10

11

12

13

where

A \rightarrow Sum of 1st and last ordinates.

B \rightarrow Sum of odd ordinates

C \rightarrow Sum of even ordinates.

Q Using Simpson's $\frac{1}{3}$ rd Rule, Evaluate $\int_0^6 \frac{1}{1+x^2} dx$ where $h=1$.

Soln Given that $\int_a^b \frac{1}{1+x^2} dx = \int_a^b f(x) dx = \int_{x_0}^{x_n} f(x) dx$

Here $f(x) = \frac{1}{1+x^2}$

$$a = x_0 = 0$$

$$b = x_n = 6, h=1.$$

x	x_0	x_1	x_2	x_3	x_4	x_5	x_6
$y = f(x)$	1	0.5	0.2	0.1	0.05	0.03	0.02

$$y_0 = f(0) = \frac{1}{1+0} = \frac{1}{1} = 1$$

$$y_1 = f(1) = \frac{1}{1+1^2} = \frac{1}{2} = 0.5$$

$$y_2 = f(2) = \frac{1}{1+2^2} = \frac{1}{5} = 0.2$$

$$y_3 = f(3) = \frac{1}{1+3^2} = \frac{1}{10} = 0.1$$

$$y_4 = f(u) = \frac{1}{1+u^2} = \frac{1}{17} = 0.059$$

$$y_5 = f(5) = \frac{1}{1+5^2} = \frac{1}{26} = 0.038$$

$$y_6 = f(6) = \frac{1}{1+6^2} = \frac{1}{37} = 0.027$$

Here $n=6$ which is even.

no. of radii are odd,

04 / 2018	01	02 03 04 05 06 07 08	09 10 11 12 13 14 15	16 17 18 19 20 21 22	23 24 25 26 27 28 29	30
APRIL / Week	13	14	15	16	17	18

W.K.T

$$\int_a^b f(x) dx = \frac{h}{3} \left[(y_0 + y_n) + 4(y_1 + y_3 + \dots + y_{n-1}) + 2(y_2 + y_4 + \dots + y_{n-2}) \right]$$

Here

$$\int_0^6 \frac{1}{1+x^2} dx = \frac{1}{3} \left[(y_0 + y_6) + 4(y_1 + y_3 + y_5) + 2(y_2 + y_4) \right]$$

$$= \frac{1}{3} \left[(1 + 0.027) + 4(0.5 + 0.4 + 0.038) + 2(0.2 + 0.059) \right]$$

04 SUNDAY • 063-302

$$= \frac{1}{3} \left[1.027 + 4(0.638) + 2(0.259) \right]$$

$$= \frac{1}{3} \left[1.027 + 2.552 + 0.518 \right]$$

$$= \frac{1}{3} [4.097] = 1.366$$

$$\int_0^6 \frac{1}{1+x^2} dx = 1.366$$

Q. Using Simpson's $\frac{1}{3}$ rd rule evaluate $\int_0^{\pi} \sin x dx$ taking 10 equal parts.

Soln Given that

$$\int_0^{\pi} \sin x dx = \int_a^b f(x) dx = \int_{x_0}^{x_n} f(x) dx$$

Here $f(x) = \sin x$

$$a = 0 = x_0, \quad b = \pi = x_n$$

$$\therefore h = \frac{x_n - x_0}{n} = \frac{b - a}{n} = \frac{\pi - 0}{10} = \frac{\pi}{10}$$




04 / 2018	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
APRIL / Week	13				14					15						16									17					18

06

TUESDAY
065-300
Week 10

(14)

MARCH 2018



$$18 \text{ Here } f(x) = \sin x$$

$$19 \quad f(0) = \sin 0 = 0$$

$$f\left(\frac{\pi}{10}\right) = \sin \frac{\pi}{10} = 0.309$$

$$2 \quad f\left(\frac{2\pi}{10}\right) = \sin \left(\frac{\pi}{5}\right) = 0.588$$

$$f\left(\frac{3\pi}{10}\right) = \sin \left(\frac{3\pi}{10}\right) = 0.809$$

03 / 2018	01 02 03 04	05 06 07 08 09 10 11	12 13 14 15 16 17 18	19 20 21 22 23 24 25	26 27 28 29 30 31
MARCH / Week	9	10	11	12	13

(15)

$$f\left(\frac{4\pi}{10}\right) = \sin\left(\frac{2\pi}{5}\right) = 0.951$$

$$f\left(\frac{5\pi}{10}\right) = \sin\left(\frac{\pi}{2}\right) = 1$$

$$f\left(\frac{6\pi}{10}\right) = \sin\left(\frac{3\pi}{5}\right) = 0.951$$

$$f\left(\frac{7\pi}{10}\right) = \sin\left(\frac{7\pi}{10}\right) = 0.809$$

$$f\left(\frac{8\pi}{10}\right) = \sin\left(\frac{4\pi}{5}\right) = 0.588$$

$$f\left(\frac{9\pi}{10}\right) = \sin\left(\frac{9\pi}{10}\right) = 0.309$$

$$f(\pi) = \sin(\pi) = 0$$

Since $n=10$ and no. of ordinates are odd.

We apply composite Simpson's $\frac{1}{3}$ rd rule.

$$\int_a^b f(x) dx = \frac{h}{3} \left[(y_0 + y_n) + 4(y_1 + y_3 + \dots + y_{n-1}) + 2(y_2 + y_4 + \dots + y_{n-2}) \right]$$

Here

$$\int_0^{\pi/10} \sin x \, dx = \frac{\pi/10}{3} \left[(y_0 + y_{10}) + 4(y_1 + y_3 + y_5 + y_7 + y_9) + 2(y_2 + y_4 + y_6 + y_8) \right]$$

$$= \frac{\pi}{30} \left[(0 + 0) + 4(0.309 + 0.809 + 1 + 0.809 + 0.309) + 2(0.588 + 0.951 + 0.951 + 0.588) \right]$$

$$= \frac{\pi}{30} \left[0 + 4(3.235) + 2(3.074) \right]$$

$$= \frac{\pi}{30} [12.944 + 6.148]$$

$$= \frac{\pi}{30} [19.1]$$

$$= (0.105) \times (19.1) = 2.006$$

observed value

(17)

But actual integration

$$\int_0^{\pi} \sin x \, dx = [-\cos x]_0^{\pi}$$

$$= -[\cos x]_0^{\pi}$$

$$= -[\cos \pi - \cos 0]$$

$$= -[-1 - 1]$$

$$= -(-2) = 2 \quad (\text{exact value})$$

$$\text{error} = \text{exact value} - \text{observed value}$$

$$= 2 - 2.006$$

$$= -0.006$$

$$\text{Absolute error} = |-0.006| = 0.006.$$

