

①

Factorial:- The product of first n natural numbers is called " n factorial".

It is denoted by (n) or $(n!)$

It is defined as

$$n! = 1 \times 2 \times 3 \times 4 \times \dots \times (n-1) \times n$$

$$n! = n \times (n-1) \times (n-2) \times \dots \times 3 \times 2 \times 1.$$

Also it is expressed as

$$n! = n \times (n-1)!$$

Ex

$$7! = 7 \times 6 \times 5 \times 4 \times 3 \times 2 \times 1 = 5040$$

$$5! = 5 \times 4 \times 3 \times 2 \times 1 = 120$$

$$6! = 1 \times 2 \times 3 \times 4 \times 5 \times 6 = 720$$

Note:-

$$0! = 1$$

$$1! = 1$$

Q. Define permutation.

(2)

Ans An arrangement of objects by taking some at a time or all at a time.

Let n and r two non-negative integers such that $0 \leq r \leq n$.

The permutation of n different objects taken r at a time is denoted by ${}^n P_r$ or $P(n, r)$.

It is defined as

$${}^n P_r \text{ or } P(n, r) = \frac{n!}{(n-r)!}$$

When $r=0$

$${}^n P_0 \text{ or } P(n, 0) = \frac{n!}{(n-0)!} = \frac{n!}{n!} = 1$$

$$\therefore {}^n P_0 = 1$$

(3)

When $r=n$

$${}_nP_n \text{ or } P(n, n) = \frac{n!}{(n-n)!} = \frac{n!}{0!} = \frac{n!}{1}$$

$$\therefore \boxed{{}_nP_n = n!}$$

Note :- We can not evaluate factorial of negative integers and fractions.

ie Ex $(-5)!$ not defined

$(\frac{2}{3})!$ not defined.

Combination :-

The selection of objects by taking some at a time or all at a time is called combination.

It is also called grouping of objects

The combination of n different objects taken r at a time is denoted by $C(n, r)$ or $\binom{n}{r}$ or $\underline{\binom{n}{r}}$ or $\underline{\binom{n}{r}}$.

It is defined as

$$\binom{n}{r} \text{ or } C(n, r) = \frac{n!}{r!(n-r)!}$$

When $0 \leq r \leq n$.

When $r=0$

$$\binom{n}{0} = \frac{n!}{0!(n-0)!} = \frac{n!}{1 \cdot n!} = 1$$

$$\binom{n}{0} = 1$$

When $r=n$

$$\binom{n}{n} = \frac{n!}{n!(n-n)!} = \frac{n!}{n! \cdot 0!} = 1$$

12 / 2018	01 02	03 04 05 06 07 08 09	10 11 12 13 14 15 16	17 18 19 20 21 22 23	24 25 26 27 28 29 30	31
DECEMBER / Week	48	49	50	51	52	53

(5)

$${}^nC_n = 1$$

$${}^nC_0 = {}^nC_n = 1$$

Note If ${}^nC_r = {}^nC_y$ then

$$r = y \text{ or } r + y = n$$

Q A bag consists of 9 red balls, 7 white balls and 4 black balls. If 2 balls are drawn at random what is the probability that

(i) both are red.

(ii) both are white

(iii) both are black

(iv) one is white other is red

(v) one is white other is black

(vi) one is white

(6)

(vi) one is red

(vii) one is black

(ix) both are same colour.

Soln Given that bag consists of
9 red, 7 white and 4 black balls.

$$\therefore \text{Total no. of balls} = 9 + 7 + 4 = 20 \text{ balls.}$$

A/2 2 balls are selected random
out of 20 balls.

$$\therefore |S| = 20C_2$$

(1) First we have to find probability
of getting both are red balls.

There are 9 red balls in the
bag.

(7)

we have to choose 2 ~~red~~ red balls out of 9 red balls.

Let that event is A.

$$\therefore |A| = {}^9C_2$$

$$\therefore P(A) = \frac{|A|}{|S|} = \frac{{}^9C_2}{{}^{20}C_2} = \frac{9!}{2!(9-2)!} \cdot \frac{20!}{2!(20-2)!}$$

$$= \frac{9 \times 8 \times 7!}{7!} \cdot \frac{9 \times 8^2}{20 \times 19} = \frac{18}{95}$$

$$\therefore P(A) = \frac{18}{95}$$

(11) we have to find the probability of getting both one white balls.

There are 7 white balls in the bag.

we choose 2 white balls from 7 white balls.

01 / 2019	01 02 03 04 05 06	07 08 09 10 11 12 13	14 15 16 17 18 19 20	21 22 23 24 25 26 27	28 29 30 31
JANUARY / Week	1	2	3	4	5

(8)

Let that event is B.

$$\therefore |B| = 7C_2.$$

$$\therefore P(B) = \frac{|B|}{|S|} = \frac{7C_2}{20C_2} = \frac{7!}{2!(7-2)!} \cdot \frac{20!}{2!(20-2)!}$$

$$= \frac{7 \times 6 \times 5!}{5!} \cdot \frac{20 \times 19 \times 18!}{18!}$$

$$= \frac{7 \times 6 \times 3}{20 \times 19} = \frac{21}{190}$$

Ⓜ We have to find the probability of both are black balls.

There are 4 black balls.

We choose 2 black balls out of 4 black balls.

Let that event is C

$$|C| = 4C_2$$

$$P(C) = \frac{|C|}{|S|} = \frac{4C_2}{20C_2} = \frac{4!/2!(4-2)!}{20!}$$

$$= \frac{4!/2!}{20!/18!} = \frac{4 \times 3 \times 2! / 2!}{20 \times 19 \times 18! / 18!}$$

$$= \frac{4 \times 3}{20 \times 19} = \frac{3}{95}$$

(IV) $P(\text{one is white and other is red})$

$$= P(W_1, R_1) = \frac{7C_1 \times 9C_1}{20C_2}$$

$$= \frac{7!}{1!(7-1)!} \times \frac{9!}{1!(9-1)!}{20!}$$

$$= \frac{7 \times 6!}{6!} \times \frac{9 \times 8!}{8!} / \frac{10 \times 19 \times 18!}{2 \times 1 \times 18!}$$

$$= \frac{7 \times 9}{10 \times 19} = \frac{63}{190}$$

10

(V) P(one is white and other is black)

$$= P(W_1, B_1) = \frac{7C_1 \times 4C_1}{20C_2}$$

13

14

$$= \frac{7 \times 4}{20 \times 19} = \frac{28}{190} = \frac{14}{95}$$

15

16

$$P(WB) = \frac{14}{95}$$

17

18

(VI) we have to find probability one is white,

19

\therefore other is either black or red.

09

SUNDAY • 343-022

$$P(W_1) = \frac{7C_1 \times 13C_1}{20C_2} = \frac{7 \times 13}{190} = \frac{91}{190}$$

12/2018	01 02	03 04 05 06 07 08 09	10 11 12 13 14 15 16	17 18 19 20 21 22 23	24 25 26 27 28 29 30	31
DECEMBER / Week	48	49	50	51	52	53

(11)

(VII) we have to find probability of one red.

- Other 13 either black or white.

$$P(R_1) = \frac{{}^9C_1 \times {}^{11}C_1}{{}^{20}C_2} = \frac{99}{190}$$

(VIII) we have to find probability of one is black.

- Other 13 either red or white.

$$P(B_1) = \frac{{}^4C_1 \times {}^{16}C_1}{{}^{20}C_2} = \frac{4 \times 16}{190}$$

$$= \frac{32}{95}$$

(12)

we have to find the probability
of ~~find~~ getting both balls on
same ~~to~~ colour.

either both balls are red
or both are white or both are
black.

$$P(2R / 2W / 2B)$$

$$= \frac{{}^9C_2}{{}^{20}C_2} + \frac{{}^7C_2}{{}^{20}C_2} + \frac{{}^4C_2}{{}^{20}C_2}$$

$$= \frac{18}{95} + \frac{21}{190} + \frac{3}{95}$$

$$= \frac{36 + 21 + 6}{190} = \frac{63}{190}$$

12 / 2018	01 02	03 04 05 06 07 08 09	10 11 12 13 14 15 16	17 18 19 20 21 22 23	24 25 26 27 28 29 30	31
DECEMBER / Week	48	49	50	51	52	53

(13)

Q In a single throw of 2 dice find the probability of sum of two faces is either 8 or 11.

Ans Since 2 dice are thrown once

$$S = \{ (1,1) (1,2) (1,3) (1,4) (1,5) (1,6) \\ (2,1) (2,2) (2,3) (2,4) (2,5) (2,6) \\ (3,1) (3,2) (3,3) (3,4) (3,5) (3,6) \\ (4,1) (4,2) (4,3) (4,4) (4,5) (4,6) \\ (5,1) (5,2) (5,3) (5,4) (5,5) (5,6) \\ (6,1) (6,2) (6,3) (6,4) (6,5) (6,6) \}$$

$$|S| = 36$$

Let A be the event sum of two faces is '8'

01 / 2019	01 02 03 04 05 06	07 08 09 10 11 12 13	14 15 16 17 18 19 20	21 22 23 24 25 26 27	28 29 30 31
JANUARY / Week	1	2	3	4	5

(14)

$$A = \{(2,6), (3,5), (4,4), (5,3), (6,2)\}$$

$$|A| = 5$$

$$\therefore P(A) = \frac{|A|}{|S|} = \frac{5}{36}$$

Let B be the event sum of two faces is 11.

$$B = \{(5,6), (6,5)\}, |B| = 2$$

$$P(B) = \frac{|B|}{|S|} = \frac{2}{36} = \frac{1}{18}$$

Here A and B are mutually exclusive events.

$$A \cap B = \emptyset$$

$$\Rightarrow P(A \cap B) = 0$$

15

SATURDAY
349-016
Week 50

DECEMBER 20

(16)

$$|S| = 36$$

Let A be the event first
face is even number

$$A = \{ (2,1) (2,2) (2,3) (2,4) (2,5) (2,6) \\ (4,1) (4,2) (4,3) (4,4) (4,5) (4,6) \\ (6,1) (6,2) (6,3) (6,4) (6,5) (6,6) \}$$

$$|A| = 18$$

$$P(A) = \frac{|A|}{|S|} = \frac{18}{36}$$

Let B be the event sum of
two faces is 8

$$B = \{ (2,6) (3,5) (4,4) (5,3) (6,2) \}$$

$$|B| = 5$$

16 SUNDAY • 350-015

12 / 2018	01 02	03 04 05 06 07 08 09	10 11 12 13 14 15 16	17 18 19 20 21 22 23	24 25 26 27 28 29 30	31
DECEMBER / Week	48	49	50	51	52	53

(17)

$$P(B) = \frac{|B|}{|S|} = \frac{5}{36}$$

$$A \cap B = \{(2,4), (4,4), (6,2)\}$$

$$|A \cap B| = 3$$

$$P(A \cap B) = \frac{|A \cap B|}{|S|} = \frac{3}{36}$$

W.K.T Addition theorem

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$= \frac{18}{36} + \frac{5}{36} - \frac{3}{36}$$

$$= \frac{18 + 5 - 3}{36} = \frac{20}{36} = \frac{5}{9}$$

$$P(A \cup B) = \frac{5}{9}$$