

# ASSIGNMENT 5

SOWMYA BANDI

Download all python codes from

```
\begin{lstlisting}
https://github.com/Sowmyabandi99/Assignment5/
blob/main/Ass5/assignment5.py
```

Latex-tikz codes from

```
https://github.com/Sowmyabandi99/Assignment5/
blob/main/Ass5/main.tex
```

## 1 QUESTION No 2.51

Find the intervals in which the function

$$f(x) = x^2 - 4x + 6 \quad (1.0.1)$$

is

- 1) increasing
- 2) decreasing

## 2 SOLUTION

Given equation can be written as

$$y = x^2 - 4x + 6 \quad (2.0.1)$$

$$\Rightarrow x^2 - 4x - y + 6 = 0 \quad (2.0.2)$$

$$\mathbf{x}^T \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \mathbf{x} + 2 \begin{pmatrix} 2 \\ -1 \end{pmatrix} \mathbf{x} + f = 0 \quad (2.0.3)$$

Here,

$$\mathbf{V} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \mathbf{u} = \begin{pmatrix} 2 \\ -1 \end{pmatrix}, f = 6 \quad (2.0.4)$$

Using eigen decomposition on  $\mathbf{V}$ ,

$$\mathbf{V} = \mathbf{PDP}^T \quad (2.0.5)$$

$$\text{where, } \mathbf{D} = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \quad (2.0.6)$$

$$\mathbf{P} = (\mathbf{p}_1 \quad \mathbf{p}_2) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad (2.0.7)$$

The vertex of parabola  $\mathbf{c}$  is given by

$$\begin{pmatrix} \mathbf{u}^T + \eta \mathbf{p}_1^T \\ \mathbf{V} \end{pmatrix} \mathbf{c} = \begin{pmatrix} -f \\ \eta \mathbf{p}_1 - \mathbf{u} \end{pmatrix} \quad (2.0.8)$$

$$\text{where, } \eta = \mathbf{u}^T \mathbf{p}_1 = \frac{-1}{2} \quad (2.0.9)$$

Substituting values from (2.0.4), (2.0.7) and (2.0.9) in (2.0.8)

$$\begin{pmatrix} -2 & -1 \\ 1 & 0 \\ 0 & 0 \end{pmatrix} \mathbf{c} = \begin{pmatrix} -6 \\ 2 \\ 0 \end{pmatrix} \quad (2.0.10)$$

Removing last row and representing (2.0.10) as augmented matrix and then converting the matrix to echelon form,

$$\begin{pmatrix} -2 & -1 & -6 \\ 1 & 0 & 2 \end{pmatrix} \xrightarrow{R_1 \leftarrow \frac{R_1}{-2}} \begin{pmatrix} 1 & \frac{1}{2} & 3 \\ 1 & 0 & 2 \end{pmatrix} \quad (2.0.11)$$

$$\xrightarrow{R_2 \leftarrow R_2 - R_1} \begin{pmatrix} 1 & \frac{1}{2} & 3 \\ 0 & -\frac{1}{2} & -1 \end{pmatrix} \quad (2.0.12)$$

$$\xrightarrow{R_2 \leftarrow (-2R_2)} \begin{pmatrix} 1 & \frac{1}{2} & 3 \\ 0 & 1 & 2 \end{pmatrix} \quad (2.0.13)$$

$$\xrightarrow{R_1 \leftarrow R_1 - \frac{R_2}{2}} \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 2 \end{pmatrix} \quad (2.0.14)$$

From (2.0.14) it can be observed that,

$$\mathbf{c} = \begin{pmatrix} 2 \\ 2 \end{pmatrix} \quad (2.0.15)$$

$\therefore$  From  $-\infty$  to  $\mathbf{c}$  the function is decreasing and from  $\mathbf{c}$  to  $\infty$  the function is increasing.

- 1)  $f$  is increasing in interval  $(2, \infty)$
- 2)  $f$  is decreasing in interval  $(-\infty, 2)$

Plot of Parabola-




Fig. 2.1: Parabola