

MSU Graduate Spatial Ecology Lab 6

Phoebe Zarnetske: plz@msu.edu; Lala Kounta: kountala@msu.edu; Kelly Kapsar: kapsarke@msu.edu

Last rev. Oct, 2025; created Sep, 2020

Lab 6: Connectivity, Interpolation

RMarkdown version of this Lab

This lab covers connectivity analysis with the **grainscape** R package. We have also included some extra OPTIONAL exercises at the end of the lab on interpolation with the **gstat** R package used in lab 5.

For the connectivity analysis, you need to hand in a **single PDF**, including output plots, any statistics, code, and text for answers to the questions.

Connectivity with **grainscape** R package

Landscape connectivity reflects how well patches on a landscape are connected. Applications in conservation biology include corridor analysis and resilience of networks of reserves. Landscape connectivity can tell us how difficult or easy it is to travel between patches across a landscape, depending on the organism or phenomena of interest. For a great example of some applications of Corridors for Conservation and examples of Corridor projects and landscape-scale experiments, check out: <https://conservationcorridor.org/>, which is co-led by Nick Haddad, Professor at Michigan State University and Director, Kellogg Biological Station Long-Term Ecological Research (LTER).

Lake Tekapo and surrounding landscape, New Zealand. Phoebe L. Zarnetske 2010.

Connectivity analysis takes into account barriers to dispersal and movement for a given organism or phenomena of interest. These include natural barriers (e.g., terrestrial vs. aquatic systems, mountain ranges, certain land cover types) and manmade (e.g., dams, roads, developed land, conversion to agriculture).

To learn about how to quantify connectivity across a landscape, you will work through the **grainscape vignette**. Here is the *RMarkdown version of the **grainscape** Connectivity Vignette*

Work through the **grainscape** vignette at least through *Modeling with Grainscape* - you can continue on if you'd like (it is a long vignette). Read the first few pages of *Landscape networks with 1D and 2D nodes* to be able to answer one of the questions below. Turn in your work on the vignette to show how far you completed, plus the questions below. Even if you do not finish the entire vignette, answer all of the following questions:

grainscape: QUESTION 1

Answer the following questions in a few sentences each: What is a minimum planar graph? What is a Voronoi tessellation? What is a 2D node and how does it compare with a 1D node?

grainscape: QUESTION 2

How do the analyses presented in this vignette relate to analyses you would perform with patch metrics from the landscapemetrics R package or Fragstats (see Lab 3 Question 1 for links to definitions on patch metrics)? How are they similar and how are they different?

grainscape: QUESTION 3

For a given species or phenomena moving across a landscape, how would you use ecological theory, and knowledge of the species natural history or phenomena, to assign resistance values and create a minimum planar graph? Use an example to explain your answer.

grainscape: QUESTION 4

How does spatial scale of land cover or elevation data affect the interpretation of connectivity?

OPTIONAL EXERCISES: Interpolation with gstat R package

If you would like more practice fitting variograms and applying a variogram fit to interpolate between points in a different R package to create a smooth surface, complete the [gstat vignette](#). After completing the vignette with the meuse data, complete the vignette again to construct variograms and interpolate using a dataset of your choice with a continuous variable that is associated with geographic coordinates. If you don't have a dataset of your own, you could use the Bryce Canyon species and site data from Lab 4 or the WEF data from Lab 5. For any dataset you choose, select just 1 response variable of interest (e.g., as is done with “zinc” with the meuse data).

Extra Resource: You may also find this exercise helpful in explaining some additional aspects of interpolation: <https://rspatial.org/raster/analysis/4-interpolation.html>. It uses the **gstat** package as well, but with precipitation data in California from the **rspatial** R package.

When working through the [gstat vignette](#) with the data you chose:

1. Fit 2 types of variograms. The example with the meuse data fits a Spherical and an Exponential model. See `?fit.variogram` and `?vgm` for options.
2. Use 2 grid resolutions to evaluate the influence of resolution on the interpolation. see below for the `cellsize()` option.
3. Consider the questions at the end of this section below.

Below is some example code for starting to use `bryceveg` and `brycesite` with **gstat**. You could modify it for your own data or use it to start the **gstat** vignette with `bryceveg` and `brycesite`.

Start working in R:

```
##### STARTING UP R
# Clear all existing data (or don't and just make sure you start with a newly opened RStudio session; s
rm(list=ls())

# Close graphics devices
graphics.off()

# Set the paths for your work
```

```

output_path<-"output"
# if this folder doesn't exist, create it
if(!dir.exists(output_path)){
  dir.create(output_path)
}

# Create the folders (directories) "data" and "lab6" - If they exist already, this command won't over-w
data_path<-(file.path("data","lab6"))
if(!dir.exists(data_path)){
  dir.create(data_path,recursive = TRUE)
}

```

If you previously saved your workspace and want to load it here, do so this way:

```
load(file.path(output_path,"lab6.RData"))
```

NOTES on R packages used in this lab:

- (1) With R Markdown, it is helpful to install packages locally before knitting (copy this into your R console before you knit). for (package in c("gstat","sp","labdsv")) { if (!require(package, character.only=T, quietly=T)) {

```

install.packages(package)
library(package, character.only=T)
}
}

```

Load the packages:

```

library(gstat)
library(sp)
library(labdsv)

```

Example code for starting with the bryceveg and brycesite data. Merge it into one dataframe, containing only points with valid coordinates.

```

data(bryceveg)
data(brycesite)
veg<-bryceveg
site<-brycesite
# Calculate species richness by plot
spc.rich <- apply(veg,1,sum)
spc.rich <- spc.rich[-109] # spc.rich[109] has NA values in coordinates, we will consider only columns
rich<-data.frame(spc.rich)

east <- na.omit(site$east)
north <- na.omit(site$north)

rich <- cbind(Row.Names = rownames(rich), rich)
site <- cbind(Row.Names = rownames(site), site)
row.names(rich)<-NULL
row.names(site)<-NULL

```

```


# Merge these data together by the Row.Names
rs<-merge(rich,site,by=c("Row.Names"))

# Make points a matrix for coordinates by referencing "east" and "north" columns.
names(rs)
rsp <- as.matrix(rs[c(8,11)])

# Make spatial points dataframe and a grid to interpolate over. This code makes the cellsize (resolution)
rs1<-SpatialPointsDataFrame(rsp,rs)
rs_grid = spsample(rs1, type = "regular", cellsize = c(100,100))
gridded(rs_grid) = TRUE
# Assign coordinates to the data
coordinates(rs)<- ~east+north

# Visualize the grid cell size relative to the point locations. Adjust the cell size if needed - you may want to
plot(geometry(rs_grid))
plot(rs, add=TRUE, col="blue")

```



The code above leaves off at the `gstat` vignette 3 *Spatial data frames*. Complete the `gstat` vignette with the data you chose to use. Consider the questions below.

gstat: QUESTION 1

How does the choice of the variogram models you chose (e.g., “Exp”, “Sph”, “Gau”, “Mat”), affect the interpolation outcome? Make 2 interpolation plots, one from each model and explain why they differ. If

they are similar, why are they are similar?

gstat: QUESTION 2

How does grid resolution alter your interpolation? Compare 2 or more grid sizes by adjusting the `cellsize()` option. What resolution is most appropriate, given your data?

gstat: QUESTION 3

What could be causing the interpolated pattern in the variable that you are observing? What are some potential explanatory variables that influence the interpolated response variable? Are there some prior studies that provide evidence of some explanatory variables?

This work is licensed under a Licensed under CC-BY 4.0 2025 by Phoebe Zarnetske.