

# Biomass Plots

Kara Dobson

3/14/2022

**TITLE:** Biomass plots

**AUTHORS:** Kara Dobson

**COLLABORATORS:** Phoebe Zarnetske, Moriah Young, Mark Hammond, Pat Bills

**DATA INPUT:** Data imported as csv files from shared Google drive L0 folder

**DATA OUTPUT:** Visualizations of biomass data

**PROJECT:** warmXtrophic

**DATE:** March 2022

```
# Clear all existing data
rm(list=ls())

#Load packages
library(tidyverse)

## -- Attaching packages ----- tidyverse 1.3.0 --
## v ggplot2 3.3.2      v purrr   0.3.4
## v tibble  3.0.3      v dplyr  1.0.2
## v tidyr   1.1.2      v stringr 1.4.0
## v readr   1.3.1      v forcats 0.5.0

## -- Conflicts ----- tidyverse_conflicts() --
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()     masks stats::lag()

library(plotrix)
library(ggpubr)
library(emmeans)

# Get data
Sys.getenv("L1DIR")
```

```

## [1] "/Volumes/GoogleDrive-117472253521803368412/Shared drives/SpaCE_Lab_warmXtrophic/data/L1/"
L1_dir<-Sys.getenv("L1DIR")
list.files(L1_dir)

## [1] "ANPP"           "climate_data"    "CN"
## [4] "Greenness"      "herbivory"       "HOB0_data"
## [7] "PAR"           "phenology"       "plant_composition"
## [10] "SLA"

kbs_biomass_21 <- read.csv(file.path(L1_dir, "ANPP/kbs_biomass_2021_L1.csv"))
umbs_biomass_21 <- read.csv(file.path(L1_dir, "ANPP/umbs_biomass_2021_L1.csv"))
kbs_biomass_21 <- kbs_biomass_21 %>% dplyr::select(-X) # get rid of "X" column that shows up (could fix
umbs_biomass_21 <- umbs_biomass_21 %>% dplyr::select(-X)

# making separate dataframe for biomass - easier in plots
kbs_biomass_only <- kbs_biomass_21 %>%
  select(-cover) %>%
  drop_na(weight_g)
umbs_biomass_only <- umbs_biomass_21 %>%
  select(-cover) %>%
  drop_na(weight_g)

# remove uninformative species
kbs_biomass_live <- kbs_biomass_only[!grepl("Litter", kbs_biomass_only$species),]
umbs_biomass_live <- umbs_biomass_only[!grepl("Litter", umbs_biomass_only$species),]
umbs_biomass_live <- umbs_biomass_live[!grepl("Standing_Dead", umbs_biomass_live$species),]
umbs_biomass_live <- umbs_biomass_live[!grepl("Surface_Litter", umbs_biomass_live$species),]


# keeping species found only in w and a
kbs_biomass_live2 <- kbs_biomass_live %>%
  group_by(species) %>%
  filter(all(c('warmed', 'ambient') %in% state))
umbs_biomass_live2 <- umbs_biomass_live %>%
  group_by(species) %>%
  filter(all(c('warmed', 'ambient') %in% state))

### Overall averages btwn treatments - barplot
biomass_sum_k <- kbs_biomass_live %>%
  group_by(state) %>%
  summarize(avg_weight = mean(weight_g, na.rm = TRUE),
            se = std.error(weight_g, na.rm = TRUE))

## `summarise()` ungrouping output (override with `.groups` argument)

ggplot(biomass_sum_k, aes(x = state, y = avg_weight, fill = state)) +
  geom_bar(position = "identity", stat = "identity", col = "black") +
  geom_errorbar(aes(ymin = avg_weight - se, ymax = avg_weight + se), width = 0.2,
               position = "identity") +
  labs(title="KBS") +
  scale_fill_manual(values = c("#a6bddb", "#fb6a4a")) +
  scale_x_discrete(labels=c("ambient" = "A", "warmed" = "W")) +
  theme_classic()


```




```
biomass_sum_u <- umbs_biomass_live %>%
  group_by(state) %>%
  summarize(avg_weight = mean(weight_g, na.rm = TRUE),
            se = std.error(weight_g, na.rm = TRUE))

## `summarise()` ungrouping output (override with `.groups` argument)


ggplot(biomass_sum_u, aes(x = state, y = avg_weight, fill = state)) +
  geom_bar(position = "identity", stat = "identity", col = "black") +
  geom_errorbar(aes(ymin = avg_weight - se, ymax = avg_weight + se), width = 0.2,
               position = "identity") +
  labs(title="UMBS")+
  scale_fill_manual(values = c("#a6bddb", "#fb6a4a")) +
  scale_x_discrete(labels=c("ambient" = "A", "warmed" = "W")) +
  theme_classic()
```




```
### Overall averages btwn treatments - boxplot
ggplot(kbs_biomass_live, aes(x = state, y = weight_g, fill=state)) +
  geom_boxplot(outlier.shape=NA, alpha=0.7) +
  geom_jitter(aes(alpha=0.6, color=state, fill=state), shape=16, size=2) +
  scale_color_manual(values = c("#a6bddb", "#fb6a4a")) +
  scale_fill_manual(values = c("#a6bddb", "#fb6a4a")) +
  labs(title="KBS")+
  scale_x_discrete(labels=c("ambient" = "Ambient", "warmed" = "Warmed")) +
  theme_classic()
```




```
ggplot(umbs_biomass_live, aes(x = state, y = weight_g, fill=state)) +
  geom_boxplot(outlier.shape=NA, alpha=0.7) +
  geom_jitter(aes(alpha=0.6, color=state, fill=state), shape=16, size=2) +
  scale_color_manual(values = c("#a6bddb", "#fb6a4a")) +
  scale_fill_manual(values = c("#a6bddb", "#fb6a4a")) +
  labs(title="UMBS")+
  scale_x_discrete(labels=c("ambient" = "Ambient", "warmed" = "Warmed")) +
  theme_classic()
```



```
### Line plot of species responses to warming
mod11_k <- lm(log(weight_g) ~ state * species, kbs_biomass_live2)
emmip(mod11_k, species~state) +
  labs(title="KBS")
```



```
mod11_u <- lm(log(weight_g) ~ state * species, umbs_biomass_live2)
emmip(mod11_u, species~state)+
  labs(title="UMBS")
```




### Differences between warmed and ambient at a species level

```
biomass_spp_k <- kbs_biomass_live2 %>%
  group_by(species, state) %>%
  summarize(avg_weight = mean(weight_g, na.rm = TRUE),
            se = std.error(weight_g, na.rm = TRUE))
```

## `summarise()` regrouping output by 'species' (override with `.groups` argument)

```
ggplot(biomass_spp_k, aes(x = species, y = avg_weight, fill = state)) +
  geom_bar(position = "dodge", stat = "identity", col = "black") +
  geom_errorbar(aes(ymin = avg_weight - se, ymax = avg_weight + se), width = 0.2,
               position = position_dodge(0.9)) +
  labs(title="KBS") +
  scale_fill_manual(values = c("#a6bddb", "#fb6a4a")) +
  scale_x_discrete(labels=c("ambient" = "A", "warmed" = "W")) +
  theme_classic()
```






```
biomass_spp_u <- umbs_biomass_live2 %>%
  group_by(species, state) %>%
  summarize(avg_weight = mean(weight_g, na.rm = TRUE),
            se = std.error(weight_g, na.rm = TRUE))

## `summarise()` regrouping output by 'species' (override with `.groups` argument)

ggplot(biomass_spp_u, aes(x = species, y = avg_weight, fill = state)) +
  geom_bar(position = "dodge", stat = "identity", col = "black") +
  geom_errorbar(aes(ymin = avg_weight - se, ymax = avg_weight + se), width = 0.2,
               position = position_dodge(0.9)) +
  labs(title="UMBS") +
  scale_fill_manual(values = c("#a6bddb", "#fb6a4a")) +
  scale_x_discrete(labels=c("ambient" = "A", "warmed" = "W")) +
  theme_classic()
```



```
## Regression between cover and biomass
# making a dataframe for regression between cover and biomass - removing uninformative species first
kbs_biomass_reg <- kbs_biomass_21[!grepl("Litter", kbs_biomass_21$species),]
kbs_biomass_reg <- kbs_biomass_reg[!grepl("Total Live", kbs_biomass_reg$species),]
kbs_biomass_reg <- kbs_biomass_reg[!grepl("Unknown", kbs_biomass_reg$species),]
kbs_biomass_reg <- kbs_biomass_reg[!grepl("Unsorted", kbs_biomass_reg$species),]
umbs_biomass_reg <- umbs_biomass_21[!grepl("Bare_Ground", umbs_biomass_21$species),]
umbs_biomass_reg <- umbs_biomass_reg[!grepl("Groundhog", umbs_biomass_reg$species),]
umbs_biomass_reg <- umbs_biomass_reg[!grepl("Litter", umbs_biomass_reg$species),]
umbs_biomass_reg <- umbs_biomass_reg[!grepl("Standing_Dead", umbs_biomass_reg$species),]
umbs_biomass_reg <- umbs_biomass_reg[!grepl("Surface_Litter", umbs_biomass_reg$species),]

# setting NA to 0 for cover or biomass for the regression
kbs_biomass_reg$cover[is.na(kbs_biomass_reg$cover)] <- 0
kbs_biomass_reg$weight_g[is.na(kbs_biomass_reg$weight_g)] <- 0
umbs_biomass_reg$cover[is.na(umbs_biomass_reg$cover)] <- 0
umbs_biomass_reg$weight_g[is.na(umbs_biomass_reg$weight_g)] <- 0


# fixing values listed as <1 for umbs
umbs_biomass_reg[umbs_biomass_reg=="<1"] <- 0
# making cover column numeric for umbs
umbs_biomass_reg$cover <- as.numeric(umbs_biomass_reg$cover)

# regression #
cor.test(kbs_biomass_reg$cover, kbs_biomass_reg$weight_g, method="pearson")

##
```

```
## Pearson's product-moment correlation
##
## data: kbs_biomass_reg$cover and kbs_biomass_reg$weight_g
## t = 17.401, df = 186, p-value < 2.2e-16
## alternative hypothesis: true correlation is not equal to 0
## 95 percent confidence interval:
## 0.7257042 0.8360155
## sample estimates:
## cor
## 0.7870724
```

```
lm1 <- lm(cover ~ weight_g, data = kbs_biomass_reg)
plot(cover ~ weight_g, data = kbs_biomass_reg)
abline(lm1)
```



```
cor.test(umbs_biomass_reg$cover, umbs_biomass_reg$weight_g, method="pearson")
```

```
##
## Pearson's product-moment correlation
##
## data: umbs_biomass_reg$cover and umbs_biomass_reg$weight_g
## t = 17.206, df = 172, p-value < 2.2e-16
## alternative hypothesis: true correlation is not equal to 0
## 95 percent confidence interval:
## 0.7332977 0.8441927
## sample estimates:
## cor
## 0.7953056
```

```
lm1 <- lm(cover ~ weight_g, data = umbs_biomass_reg)
plot(cover ~ weight_g, data = umbs_biomass_reg)
abline(lm1)
```

